

This work is licensed under a Creative Commons Attribution License (CC BY 4.0).

Research article

urn:lsid:zoobank.org:pub:F539B8D7-E174-4345-8EF7-F0207A104CDE

Five new species of the planthopper genus *Atracis* Stål (Hemiptera, Fulgoromorpha, Flatidae) from China

Deqiang AI¹, Lingfei PENG² & Yalin ZHANG^{3,*}

^{1,2,3}Key Laboratory of Plant Protection Resources and Pest Management, Ministry of Education, Entomological Museum, Northwest A&F University, Yangling, Shaanxi, 712100, P.R. China.

²Key Laboratory of Integrated Pest Management for Fujian-Taiwan Crops, Ministry of Agriculture, Fujian Agriculture and Forestry University, Fuzhou, Fujian, 350002, P.R. China.

*Corresponding author: yalinzh@nwsuaf.edu.cn

¹Email: aideqiang@nwafu.edu.cn

²Email: lingfeipeng@fafu.edu.cn

¹urn:lsid:zoobank.org:author:B8603913-50C3-479C-877F-7DF322067B08

²urn:lsid:zoobank.org:author:5FB1448D-3BED-4DC7-8CD8-F0D62DE0D7C7

³urn:lsid:zoobank.org:author:918B4F5D-D0A7-4E64-B866-5A2D94CA96B8

Abstract. This paper treats the planthopper genus, *Atracis* Stål, 1866 (Flatidae: Flatoidinae) in China. Five new species, *A. ocularia* sp. nov., *A. patefacta* sp. nov., *A. punctulata* sp. nov., *A. sphaerica* sp. nov. and *A. unguilata* sp. nov., are described and a checklist of all its known species from China is provided.

Keywords. Diversity, Fulgoroidea, morphology, taxonomy, Oriental region.

Ai D., Peng L. & Zhang Y. 2020. Five new species of the planthopper genus *Atracis* Stål (Hemiptera, Fulgoromorpha, Flatidae) from China. *European Journal of Taxonomy* 717: 108–128. <https://doi.org/10.5852/ejt.2020.717.1105>

Introduction

The planthopper subfamily Flatoidinae Melichar, 1901 belongs together with the subfamily Flatinae Melichar, 1901 to the family Flatidae Spinola, 1839. It comprises 31 previously described genera distributed worldwide (Metcalf 1957; Bourgoin 2020). Its members are distinguished externally mainly by the tegmina positioned more or less flat, not vertically, usually more than two times as long as wide, the costal margin often undulate and the costal membrane obviously wider than the costal cell.

Currently, in the Oriental region, the subfamily Flatoidinae comprises eight genera (*Atracis* Stål, 1866; *Bochara* Distant, 1906; *Cerfennia* Stål, 1870; *Cisatra* Melichar, 1923; *Gaja* Distant, 1906; *Malleja* Medler, 1990; *Ortracis* Medler, 1996; *Uxantis* Stål, 1870), of which one is recorded from China (Fang 1989; Bourgoin 2020).

The Flatoidinae genus *Atracis* was established by Stål (1866) without designation of a type species. Distant (1906) studied the genus *Atracis* and designated *Flata pyralis* Guérin-Méneville, 1831 as its type species. Stål established the genus *Uxantis* in 1870 and did not designate a type species either (Stål

1870). Banks (1910) considered it as a subgenus of *Atracis* and designated *Uxantis consputa* Stål, 1870 as its type species. Metcalf (1957) synonymized *Franciscus* Distant, 1910 and *Grapaldus* Distant, 1914 with the genus *Atracis*. Medler (1986) proposed the new name *Staliana* to replace *Atracis* and moved all the species of *Atracis* to it, but it was an invalid treatment (Fletcher 2009 and updates). Medler (1988) excluded from *Atracis* all Afrotropical species and treated *Atracis* as an Oriental genus.

In this paper, five new species are described and a checklist of all known species in China is provided.

Material and methods

The venation terminology follows Bourgoin *et al.* (2015), the male genitalia terminology follows Bourgoin & Huang (1990), and the female genitalia terminology follows Bourgoin (1993). All measurements are in millimeters (mm). The external morphology was observed under a Leica ZOOM 2000 stereo microscope. The genital segments of the examined specimens were treated with 10% NaOH solution at 100°C for several minutes, rinsed with water, immersed in 2–3 droplets of glycerol and dissected following standard procedures. They were observed under a Leica ZOOM 2000 stereo microscope. Photographs of the specimens were made using a Leica M205A microscope with a Leica DFC Camera. Images were produced using the software LAS (Leica Application Suite) ver. 3.7. Photographs were treated with Adobe Photoshop CS4. One female of *A. sphaerica* sp. nov. is deposited in the Beijing Natural History Museum, China (BMNHC), and two males of *A. unguilata* sp. nov. are deposited in the Chinese Academy of Forestry (CAF), Beijing, China or the Sun Yat-sen University (SYSU), Guangzhou, China. All other specimens are deposited in the Entomological Museum, Northwest A&F University (NWAFU), Yangling, China.

The following measurements were used in this study:

Total length = length of specimen from head apex to tegmina apex (in dorsal view)

Length of tegmen = length from the base to the posterior margin of tegmen

Width of tegmen = width at the widest part of tegmen

Width of costal membrane versus costal cell = width of costal membrane at the widest part versus width of costal cell at the widest part

Results

Class Insecta Linnaeus, 1758
 Order Hemiptera Linnaeus, 1758
 Suborder Fulgoromorpha Evans, 1946
 Superfamily Fulgoroidea Latreille, 1807
 Family Flatidae Spinola, 1839
 Subfamily Flatoidinae Melichar, 1901

Genus *Atracis* Stål, 1866

Atracis Stål, 1866: 237 (type species: *Flata pyralis* Guérin-Méneville, 1831, by subsequent designation in Distant 1906: 450).

Uxantis Stål, 1870: 775.

Franciscus Distant, 1910: 337 (type species: *Flatoides fasciatus* Walker, 1870, by original designation).

Grapaldus Distant, 1914: 335 (type species: *Grapaldus corticinus* Distant, 1914, by original designation).

Staliana Medler, 1986: 18, new name for *Atracis* Stål, 1866 (type species: *Elidiptera inaequalis* Walker, 1858, by original designation).

Atracis – Metcalf 1957: 478.

Uxantis – Melichar 1902: 160, subgenus of *Atracis* Stål, 1866. — Banks 1910: 47, valid genus, designated type species: *Uxantis consputa* Stål, 1870 — Medler 1986: 18, synonymized.

Franciscus – Metcalf 1957: 513, synonymized.

Grapaldus – Metcalf 1957: 455, synonymized.

Staliana – Fletcher 2009 and updates, invalid new name.

Distribution

Sri Lanka, Malaysia, China (Zhejiang, Fujian, Taiwan, Guangxi and Hainan Province), India, Japan, Indonesia, Bangladesh.

Checklist of species from China

A. fimbria (Walker, 1851)

Elidiptera fimbria Walker, 1851: 331.

Flatoides fimbria (Walker) – Stål 1862: 489.

Atracis fimbria (Walker) – Melichar 1902: 199.

Distribution: China (Yunnan), India, Sri Lanka.

A. formosana Jacobi, 1915

Atracis formosana Jacobi, 1915: 177.

Distribution: China (Taiwan), Japan (Honshu, Kyushu).

A. hainanensis Distant, 1912

Atracis hainanensis Distant, 1912: 470.

Distribution: China (Hainan).

A. koshunensis Kato, 1933

Atracis koshunensis Kato, 1933: 466, fig. 2.

Distribution: China (Taiwan).

A. kotoshonis Matsumura, 1940

Atracis kotoshonis Matsumura, 1940: 48.

Distribution: China (Taiwan).

A. mucida Jacobi, 1915

Atracis mucida Jacobi, 1915: 177.

Distribution: China (Fujian, Taiwan, Hainan).

A. obscura Zia, 1935

Atracis obscura Zia, 1935: 534, fig. 5.

Distribution: China (Guangxi), India.

A. ocularia sp. nov.

Distribution: China (Hainan).

A. patefacta sp. nov.

Distribution: China (Yunnan).

A. pruinosa (Walker, 1858)

Elidiptera pruinosa Walker, 1858: 75.

Flatoides pruinosa (Walker) – Stål 1862: 489.

Atracis pruinosa – Stål 1866: 250.

Distribution: Northern China.

A. punctulata sp. nov.

Distribution: China (Guangdong).

A. sphaerica sp. nov.

Distribution: China (Guangxi, Zhejiang).

A. unguis sp. nov.

Distribution: China (Hainan).

Key to species of *Atracis* in China

1. Pronotum with three carinae *A. koshunensis* Kato, 1933
- Pronotum without carina 2

2. Small size, body length less than 7 mm *A. kotoshonis* Matsumura, 1940
- Medium size, body length more than 10 mm 3

3. Tegminal costal margin evenly convex 4
- Tegminal costal margin sinuate 5

4. Vertex 1.9 times wider than long *A. formosana* Jacobi, 1915
- Vertex as long as wide *A. obscura* Zia, 1935

5. Tegmina marked with large spot (Fig. 1B) *A. ocularia* sp. nov.
- Tegmina variously marked but lacking large spot 6

6. Vertex wider than long 7
- Vertex much longer than wide 8

7. Vertex 1.1 times as wide as long; frons with one short carina (Fig. 8D–E) *A. sphaerica* sp. nov.
- Vertex obviously wider than long; frons with two weak carinae *A. pruinosa* (Walker, 1858)

8. Frons without longitudinal median carina *A. mucida* Jacobi, 1915
- Frons with short longitudinal median carina at dorsal portion 9

9. Posterior lateral margins of vertex convex or humped, higher than anterior portion 10
- Posterior lateral margins of vertex not humped, at same level as anterior portion (Fig. 3D) *A. patefacta* sp. nov.

10. Head and thorax slightly mottled with dark color *A. hainanensis* Distant, 1912
- Head and thorax heavily marked with fuscous patches 11

11. Periandrium with pair of processes, apex of processes forked (Fig. 12C) *A. unguis* sp. nov.
- Periandrium with pair of processes, but apex of processes not forked 12

12. Processes of periandrium extending ventrad and crossed *A. fimbria* (Walker, 1851)
- Processes of periandrium extending cephaloventrad and parallel (Fig. 7A) ... *A. punctulata* sp. nov.

Atracis ocularia sp. nov.
urn:lsid:zoobank.org:act:4A7434A6-B277-4254-B2DE-7E807E2140A8
Figs 1–2

Diagnosis

This new species is similar to *A. punctulata* sp. nov., but differs from the latter by the following characters: tegmina with a large black patch (without a large patch in *A. punctulata* sp. nov.); costal membrane 3.2 times as wide as costal cell (4 times as wide as costal cell in *A. punctulata* sp. nov.); periandrium with a pair of processes, apex of processes forked (periandrium with paired processes, apex of processes not forked in *A. punctulata* sp. nov.).




Fig. 1. *Atracis ocularia* sp. nov., holotype, ♂ (NWAFU). **A.** Habitus, dorsal view. **B.** Tegmen. **C.** Hind leg apex. **D.** Head and thorax, dorsal view. **E.** Head and thorax, left lateral view. **F.** Face. **G.** Male genitalia (phallic complex removed), right lateral view. **H.** Male anal tube, dorsal view. **I.** Phallic complex, right lateral view.

Etymology

The specific name is a feminine adjective ‘*ocularia*’ to emphasize a big eye shaped spot on the tegmina.

Type material

Holotype

CHINA • ♂; Hainan Province, Mt Jianfengling; 7 May 1985; Li Weihua and Zhang Jinghong leg.; NWAFU.

Description

MEASUREMENTS. Male length (N = 1) (including tegmen): 13.5 mm, length of tegmen: 12.1 mm.

COLORATION. Body and tegmina yellowish brown with black markings; vertex yellowish brown marked with dark patches; frons light brown with dorsal apex and longitudinal carina black; gena yellowish




Fig. 2. *Atracis ocularia* sp. nov., holotype, ♂ (NWAFU). **A.** Male genitalia, right lateral view. **B.** Phallic complex, right lateral view. **C.** Phallic complex, dorsal view. **D.** Phallic complex, ventral view. **E.** Periandrium, right lateral view. **F.** Periandrium, dorsal view. **G.** Periandrium, ventral view. **H.** Aedeagus, right lateral view. **I.** Aedeagus, dorsal view. **J.** Aedeagus, ventral view. Scale bars = 0.5 mm.

brown, dorsal portion and anterior portion of eye marked with darker brown color; eyes brown, ocelli milk white; dorsal apex of antennal segment II black marked; pronotum and mesonotum yellowish brown, lateral carinae black, disc mottled; tegmina yellowish brown, veins brown or black, a large black patch positioned after fork of vein MP, 1.3 mm in diameter (Fig. 1A–B, D–F).

HEAD AND THORAX. Head anterior margin convex; vertex pentagonal; disc pressed; longer than wide; lateral margins parallel, highly raised at posterior portion; posterior margin raised, median groove full length of vertex, apex T-forked (Fig. 1D–F). Frons disc flat, 1.3 times as long as wide, median longitudinal carina almost half length of frons, lateral margins moderately raised (Fig. 1F). Frontoclypeal suture pressed, truncate; clypeus convex. Rostrum extending beyond metatrochanter (Fig. 1F). Antennae short. Pronotum with anterior margin truncate, lateral side thick and tended up, median carina obscure, ventral margin rolled upwards, postocular eminence evenly convex (Fig. 1D–E). Mesonotum slightly humped, anterior margin convex, disc flat, without carina but lines. Metatibia with six spines apically, metatarsal basal segment with nine spines apically (Fig. 1C). Tegmina elongate, 13 mm in length, 5 mm in width; costal membrane 3.2 times as wide as costal cell at the level of bulla; costal margin undulate, apical margin convex and sinuate, sutural margin truncate, apical and sutural angles convex; vein ScP+RA elevated ridgelike above plane of vein P_c+CA and crossing pustulate bulla, CuA forked, clavus and costal membrane with reticulated crossveins and pustulate; one subapical line present (Fig. 1B).

MALE GENITALIA. Pygofer ring-like, anterior margin concave, posterior margin convex, dorsal and ventral margins more or less truncate (Figs 1G, 2A). Genital style ventral and apical margins convex, dorsal margin concave, with a process (Figs 1G, 2A). Male anal tube elongated, evenly bent down (Figs 1G–H, 2A). Phallic complex arched. Periandrium tubular. Dorsal part of periandrium slightly shorter than ventral part in lateral view, median lobe narrow and acute. Each side of lateral part of periandrium with forked process, recurved. Ventral part of periandrium distinctly tapering apicad in ventral view (Figs 1I, 2B–D). Aedeagus bipartite; each side of apex without process; apex of lateral portion with even prominence.

Distribution

Hainan Province, China.

Atracis patefacta sp. nov.

urn:lsid:zoobank.org:act:F084DFB5-F36E-4964-A9C2-B24260B2ECE3

Figs 3–5

Diagnosis

This new species looks similar to *A. unguilata* sp. nov., but differs from the latter by: pronotum anterior margin concave in the middle (truncate in *A. unguilata* sp. nov.); process of lateral lobe of periandrium not forked (forked in *A. unguilata* sp. nov.).

Etymology

The specific name is derived from the Latin word ‘*patefactus*’ which refers to the flat tegmina.

Type material

Holotype

CHINA • ♂; Yunnan Province, Mengyang, Wild elephants Valley; 800 m a.s.l.; 5 May 2009; Zhang Lei leg.; NWAFU.

Paratype

CHINA • 1 ♀; Yunnan Province, Banna, Menglong; 28 Apr. 1958; Meng Song County leg.; NWAFU.

Description

MEASUREMENTS. Male length (N = 1) (including tegmen): 13.3 mm, length of tegmen: 12.0 mm; female length (N = 1) (including tegmen): 15.5 mm, length of tegmen: 13.1 mm.




Fig. 3. *Atracis patefacta* sp. nov. **A.** Habitus, dorsal view. **B.** Tegmen. **C.** Hind leg apex. **D.** Head and thorax, dorsal view. **E.** Head and thorax, left lateral view. **F.** Face. **G.** Male anal tube, dorsal view. **H.** Male genitalia, right lateral view. **I.** Phallic complex, right lateral view. **J.** Phallic complex, dorsal view. **K.** Phallic complex, ventral view.

COLORATION. Body and tegmina green with black markings; vertex green marked with dark patches; frons light green with dorsal apex and longitudinal carina fuscous; gena yellowish green, dorsal portion and anterior portion of eye marked with darker brown color; eyes black, ocelli milk white; pronotum dull green, median carina fuscous; mesonotum dull green, heavily mottled, each posterior margin with triangular black spot; tegmina dull green, cross vein of costal area and veins of apical cell brown or black (Fig. 3A–B, D–F).

HEAD AND THORAX. Head anterior margin pointed, about 80°; vertex pentagonal; disc pressed; longer than wide; lateral margins slightly raised; posterior margin raised, pressed in middle; median groove full length of vertex, apex Y-forked (Fig. 3D–F). Frons disc flat, 1.4 times as wide as wide, median longitudinal carina almost half length of frons, lateral margins moderately raised (Fig. 3F). Frontoclypeal suture pressed, evenly convex; clypeus convex. Rostrum extending beyond metatrochanter (Fig. 3F). Antennae short. Pronotum with anterior margin concave in middle, disc pressed, lateral side thick and tended up, median carina obscure, ventral margin rolled upwards, postocular eminence evenly convex (Fig. 3D–E). Mesonotum humped, anterior margin convex, disc flat, without carina but lines (Fig. 3D–E). Metatibia with six spines apically, metatarsal basal segment with two large spines and nine small spines apically (Fig. 3C). Tegmina elongate, 11.5 mm in length, 4.5 mm in width, costal membrane 3.8 times as wide as costal cell at level of bulla; costal margin undulate, apical margin convex and sinuate, sutural margin truncate, apical and sutural angles convex; vein ScP+RA elevated ridgelike above plane of vein $Pc+CA$ and crossing pustulate bulla, CuA forked, clavus with few crossveins; one subapical line present (Fig. 3B).

MALE GENITALIA. Pygofer ring-like, anterior margin strongly concave, posterior margin convex, ventral margin evenly convex, dorsal margin concaved (Fig. 3H). Genital style ventral and apical margins




Fig. 4. *Atracis patefacta* sp. nov. **A.** Periandrium, right lateral view. **B.** Periandrium, dorsal view. **C.** Periandrium, ventral view. **D.** Aedeagus, right lateral view. **E.** Aedeagus, dorsal view. **F.** Aedeagus, ventral view. **G.** Gonapophysis VIII, right lateral outside view. **H.** Gonapophysis IX, right lateral view. **I.** Gonapophysis IX, dorsal view. **J.** Gonoplac, right lateral outside view. Scale bars = 0.5 mm.




Fig. 5. *Atracis patefacta* sp. nov., paratype, ♀ (NWAFU). **A.** Female genitalia, right lateral view. **B.** Anal segment, dorsal view.

convex, dorsal margin concave, with a process apically (Fig. 3H). Male anal tube elongated, apex bent down (Fig. 3G–H). Phallic complex slightly arched (Fig. 3I–K). Periandrium tubular. Dorsal part of periandrium distinctly shorter than ventral part in lateral view, median lobe acute. Each side of lateral part of periandrium with recurved process, not forked. Ventral part of periandrium distinctly tapering apicad in ventral view (Fig. 4A–C). Aedeagus bipartite; each side of apex without process; apex of lateral portion with obviously raised prominence (Fig. 4D–F).

FEMALE GENITALIA. Female anal segment small, oval in dorsal view, apex slightly concave (Fig. 5B). Gonapophysis VIII slender, triangular, apex with 8 teeth, dorsal margin evenly concave, ventral margin convex (Fig. 4G); gonapophysis IX short and triangular, apex acute (Fig. 4H, I); gonoplac with arrays of strong marginal teeth (Figs 4J, 5A).

Distribution

Yunnan Province, China.

Atracis punctulata sp. nov.

urn:lsid:zoobank.org:act:E27913C7-0D22-47D6-9843-7AB43A88E75A

Figs 6–7

Diagnosis

Atracis punctulata sp. nov. is similar to *A. fimbria*, but differs from the latter by: body and tegmina greenish and yellow (brownish in *A. fimbria*); disc of vertex pressed (flat in *A. fimbria*); precesses of periandrium extending cephaloventrad and parallel (ventrad and crossed in *A. fimbria*).

Etymology

The specific name is derived from the Latin word ‘*punctum*’, which refers to the mottled color of the tegmina.

Type material

Holotype

CHINA • ♂; Guangdong Province, Mt Nanling; 8 May 2009; NWAFU.

Description

MEASUREMENTS. Male length (N = 1) (including tegmen): 15.1 mm, length of tegmen: 13.1 mm.

COLORATION. Body and tegmina greenish and yellow with black markings; vertex green heavily marked with dark color; frons light greenish yellow and mottled fuscous; gena yellowish green, dorsal portion

and anterior portion of eye marked with darker brown color; eyes black, ocelli milk white; pronotum dull green and yellow, heavily fuscous; mesonotum dark brown, marked with black spots, heavily mottled; tegmina light greenish with black markings (Fig. 6A–B, D–F).




Fig. 6. *Atracis punctulata* sp. nov., holotype, ♂ (NWAFU). **A.** Habitus, dorsal view. **B.** Tegmen. **C.** Hind leg apex. **D.** Head and thorax, dorsal view. **E.** Head and thorax, left lateral view. **F.** Face. **G.** Male anal tube, dorsal view. **H.** Male genitalia, right lateral view. **I.** Phallic complex, right lateral view. **J.** Phallic complex, dorsal view. **K.** Phallic complex, ventral view.

HEAD AND THORAX. Head anterior margin convex, vertex pentagonal; disc pressed; as long as wide, lateral margins highly raised at posterior portion; posterior raised, pressed in the middle; median groove full length of vertex, apex Y-forked (Fig. 6D–E). Frons disc flat, 1.3 times longer than wide, median longitudinal carina no more than half length of frons, lateral carinae positioned at dorsal portion no longer than median longitudinal carina; lateral margins sinuate, highly raised (Fig. 6F). Frontoclypeal suture pressed, truncate; clypeus convex. Rostrum extending beyond metatrochanter (Fig. 6F). Antennae short. Pronotum with anterior margin truncate, slightly concave in middle, disc pressed, lateral side thick and tended up, median carina weakly raised, ventral margin rolled upwards, postocular eminence evenly convex (Fig. 6D–E). Mesonotum humped, anterior margin convex, disc flat, tricarinate (Fig. 6D–E). Metatibia with six spines apically, metatarsal basal segment with two large spines and eight small spines apically (Fig. 6C). Tegmina elongate, 13.8 mm in length, 6 mm in width, costal membrane 4 times as wide as costal cell at level of bulla; costal margin undulate, apical margin convex and sinuate, sutural margin truncate, apical and sutural angles convex; vein ScP+RA elevated ridgelike above plane of vein $Pc+CA$ and crossing pustulate bulla, CuA forked, clavus with few crossveins; one subapical line present (Fig. 6B).

MALE GENITALIA. Pygofer ring-like, anterior margin strongly concave, posterior margin convex and slightly sinuate, ventral margin truncate, dorsal margin concave (Fig. 6H). Genital style ventral and apical margins convex, dorsal margin concave, with a process apically (Fig. 6H). Male anal tube elongated, apex bent down (Fig. 6G–H). Phallic complex slightly arched. Periandrium tubular. Dorsal part of periandrium distinctly shorter than ventral part in lateral view, median lobe short and acute; each side of apex with three or four small teeth. Each side of lateral part of periandrium with recurved process, not forked, extending to half length of periandrium. Ventral part of periandrium distinctly slender and tapering apicad in ventral view (Figs 6I–K, 7A–C). Aedeagus bipartite; each side of apex without process; apex of lateral portion with an evenly pointed prominence (Fig. 7D–F).

Distribution

Guangdong Province, China.




Fig. 7. *Atracis punctulata* sp. nov. **A.** Periandrium, right lateral view. **B.** Periandrium, dorsal view. **C.** Periandrium, ventral view. **D.** Aedeagus, right lateral view. **E.** Aedeagus, dorsal view. **F.** Aedeagus, ventral view. Scale bars = 0.5 mm.

Atracis sphaerica sp. nov.
urn:lsid:zoobank.org:act:64303802-253F-42FC-B7F9-DD1BFFB0F3AE
Figs 8–10

Diagnosis

This new species looks similar to *A. mucida*, but differs by: head with anterior margin angulated at about 90° (slightly acute angled in *A. mucida*); vertex 1.1 times as wide as long (1.1 times as long as wide in *A. mucida*); frons with one obviously short median carina (median carina almost absent in *A. mucida*).

Etymology

The specific name is a Latin feminine adjective ‘*sphaerica*’ to refer to the globular shape of the head.




Fig. 8. *Atracis sphaerica* sp. nov. **A.** Habitus, dorsal view. **B.** Tegmen. **C.** Hind leg apex. **D.** Head and thorax, dorsal view. **E.** Head and thorax, left lateral view. **F.** Face. **G.** Male anal tube, dorsal view. **H.** Male genitalia (phallic complex removed), right lateral view. **I.** Phallic complex, right lateral view.

Type material**Holotype**

CHINA • ♂; Guangxi Province, Longzhou, Nonggang; 19 May 1985; Li Weihua and Zhang Jinghong leg.; NWAFU.

Paratypes

CHINA • 1 ♀; Guangxi Province, Mt Daqingshan; 15 Jun. 1963; Liu Sikong leg.; BMNH • 1 ♀; Zhejiang Province, Mt Wuyanling; 20 Jul. 1983; Zhen Bin leg.; NWAFU.

Description

MEASUREMENTS. Male length (N = 1) (including tegmen): 14.2 mm, length of tegmen: 11.9 mm; female length (N = 2) (including tegmen): 16.1–16.5 mm, length of tegmen: 13.0–13.5 mm.

COLORATION. Body and tegmina brown with dark brown marks; vertex, pronotum and mesonotum mottled, median carina of pronotum dark brown; frons light brown with apex mottled fuscous; gena yellowish brown, dorsal portion and anterior portion of eye marked with darker brown color; eyes black,




Fig. 9. *Atracis sphaerica* sp. nov. **A.** Male genitalia (phallic complex removed), right lateral view. **B.** Phallic complex, right lateral view. **C.** Phallic complex, dorsal view. **D.** Phallic complex, ventral view. **E.** Periandrium, right lateral view. **F.** Periandrium, dorsal view. **G.** Periandrium, ventral view. **H.** Aedeagus, right lateral view. **I.** Aedeagus, dorsal view. **J.** Aedeagus, ventral view. **K.** Gonapophysis VIII, right lateral outside view. **L.** Gonapophysis IX, lateral view. **M.** Gonapophysis IX, dorsal view. **N.** Gonoplac, right lateral outside view. Scale bars = 0.5 mm.

ocelli milk white; tegmina light brown, slightly transparent, marked with brown dots, apical portion and costal area dark brown (Fig. 8A–B, D–F).

HEAD AND THORAX. Head anterior margin angulated, about 90°; vertex pentagonal; disc pressed; 1.1 times as wide as long, lateral margins highly raised; posterior margin raised, pressed in the middle; median groove full length of vertex, apex Y-forked (Fig. 8D). Frons disc flat, 1.1 times as long as wide, median longitudinal carina very short positioned at dorsal portion, lateral margins moderately raised (Fig. 8F). Frontoclypeal suture pressed, truncate; clypeus convex. Rostrum extending to metatrochanter (Fig. 8F). Antennae short. Pronotum with anterior margin truncate, slightly concave in the middle, disc pressed, lateral side thick and tended up, median carina weakly raised, ventral margin rolled upwards, postocular eminence evenly convex (Fig. 8D–E). Mesonotum humped, anterior margin convex, disc flat, without carina (Fig. 8D–E). Metatibia with six spines apically, metatarsal basal segment with two large spines and six small spines apically (Fig. 8C). Tegmina elongate, 11.6 mm in length, 5 mm in width, costal membrane 3.5 times as wide as costal cell at level of bulla; costal margin slightly undulate, apical margin convex, sutural margin truncate, apical and sutural angles convex; vein ScP+RA elevated ridgelike above plane of vein $Pc+CA$ and crossing pustulate bulla, CuA forked, clavus with few crossveins; one subapical line present (Fig. 8B).

MALE GENITALIA. Pygofer ring-like, anterior margin concave, posterior margin convex, ventral margin truncate, dorsal margin concave (Figs 8H, 9A). Genital style ventral, dorsal and apical margins truncate, dorsal apex with a process apically, ventral apex concave (Figs 8H, 9A). Male anal tube elongated and arched (Fig. 8G–H). Phallic complex arched (Figs 8I, 9B–D). Periandrium tubular. Dorsal part of periandrium distinctly shorter than ventral part in lateral view, median lobe narrow and acute. Each side of lateral part of periandrium with a recurved process, not forked. Ventral part of periandrium distinctly slender and tapering in ventral view (Fig. 9E–G). Aedeagus bipartite; fourth of dorsal portion apparently elevated; each side of apex without process; apex of lateral portion with a sharply short prominence (Fig. 9H–J).

FEMALE GENITALIA. Female anal segment small, oval in dorsal view, apex slightly concave (Fig. 10B). Gonapophysis VIII slender, triangular, apex with 7 teeth, dorsal margin evenly concave, ventral margin convex (Fig. 9K); gonapophysis IX short and slender, apex acute (Fig. 9L–M); gonoplac with arrays of strong marginal teeth (Figs 9N, 10A).

Distribution

Zhejiang and Guangxi Province, China.




Fig. 10. *Atracis sphaerica* sp. nov. **A.** Female terminalia, right lateral view. **B.** Anal segment, dorsal view.

Atracis ungulata sp. nov.
 urn:lsid:zoobank.org:act:1F84C668-54FD-4A89-8F15-F0898D4654B8
 Figs 11–12

Diagnosis

This new species is similar to *A. hainanensis*, but differs from the latter by: vertex pentagonal (triangular in *A. hainanensis*); disc of vertex flat (evenly elevated in *A. hainanensis*); mesonotum without carina (tricarinate in *A. hainanensis*); dorsal lobe of periandrium short (long in *A. hainanensis*).

Etymology

The specific name is a Latin feminine adjective ‘*ungulata*’, which means a periandrium with one pair of forked processes at the apex.

Type material

Holotype

CHINA • ♂; Hainan Province, Mt Jianfengling; 4 Apr. 1984; Lin Youdong leg.; CAF.

Paratypes

CHINA • 1 ♂; Hainan Province; 12 May 1936; NWAFU • 1 ♂; Hainan Province, Mt Jianfengling, Tianchi; 30 Mar. 1982; Chen Huanqiang leg.; SYSU.




Fig. 11. *Atracis ungulata* sp. nov. **A.** Habitus, dorsal view. **B.** Head and thorax, dorsal view. **C.** Head and thorax, left lateral view. **D.** Face. **E.** Hind leg apex. **F.** Male genitalia (male anal tube and phallic complex removed), right lateral view. **G.** Phallic complex, right lateral view. **H.** Male anal tube, dorsal view.

Description

MEASUREMENTS. Male length (N = 3) (including tegmen): 14.7–14.9 mm, length of tegmen: 11.9–12.1 mm.

COLORATION. Body and tegmina brown with dark brown marks; vertex, pronotum and mesonotum mottled, median carina of pronotum dark brown; frons light brown with apex mottled fuscous; gena yellowish brown, dorsal portion and anterior portion of eye marked with darker brown color; eyes black, ocelli milk white; tegmina light brown, marked with darker marks, apical portion and costal area dark brown (Fig. 11A–D).

HEAD AND THORAX. Head anterior margin angulated, about 80°; vertex pentagonal; disc flat; 1.1 times as long as wide, lateral margins highly raised; posterior margin raised; median groove full length of vertex, apex Y-forked (Fig. 11B–C). Frons disc flat, 1.4 times as long as wide, median longitudinal carina very short positioned at dorsal portion, lateral margins slightly sinuate, moderately raised (Fig. 11D).




Fig. 12. *Atracis ungulata* sp. nov. **A.** Tegmen. **B.** Male genitalia, right lateral view. **C.** Phallic complex, right lateral view. **D.** Phallic complex, dorsal view. **E.** Phallic complex, ventral view. **F.** Periandrium, right lateral view. **G.** Periandrium, dorsal view. **H.** Periandrium, ventral view. **I.** Aedeagus, right lateral view. **J.** Aedeagus, dorsal view. **K.** Aedeagus, ventral view. Scale bars = 0.5 mm.

Frontoclypeal suture pressed, convex, clypeus convex. Rostrum extending beyond metatrochanter (Fig. 11D). Antennae short. Pronotum anterior margin truncate, disc pressed, lateral side thick and tended up, median carina weakly raised, ventral margin rolled upwards, postocular eminence evenly convex (Fig. 11B–C). Mesonotum humped, anterior margin convex, disc flat, without carina (Fig. 11B–C). Metatibia with six spines apically, metatarsal basal segment with two large spines and seven or eight small spines apically (Fig. 11E). Tegmina elongate, 12.9 mm in length, 5.4 mm in width, costal membrane 3.5 times as wide as costal cell at level of bulla; costal margin evenly convex, very weak undulated, apical margin convex, sutural margin truncate, apical and sutural angles convex; vein ScP+RA elevated ridgelike above plane of vein P_c+CA and crossing pustulate bulla, CuA forked, clavus with few crossveins; one subapical line present (Fig. 12A).

MALE GENITALIA. Pygofer ring-like, anterior and posterior margins sinuate, ventral margin truncate, dorsal margin sinuate (Figs 11F, 12B). Genital style ventral margin convex, dorsal margin weakly concave with process apically (Figs 11F, 12B). Male anal tube elongated and arched (Figs 11H, 12B). Phallic complex arched (Figs 11G, 12C–E). Periandrium tubular. Dorsal part of periandrium distinctly shorter than ventral part in lateral view, median lobe short and acute. Each side of lateral part of periandrium with recurved process which apex forked. Ventral part of periandrium distinctly slender and tapering in ventral view (Fig. 12F–H). Aedeagus bipartite; each side of apex without process; apex of lateral portion with ear-like prominence (Fig. 12I–K).

Distribution

Hainan Province, China.

Discussion

Male genitalia are usually regarded as the most important morphological characters in the taxonomical study of Flatidae. Based on the morphological characters, Medler (1990, 1993, 1996, 1999) and Fang (1989) illustrated the male genitalia for species of *Atracis*, as did for the five new species described in this paper and the specimens examined in China. It is clear that species of *Atracis* from China, India, Sri Lanka, Indonesia, Brunei and Malaysia are several species-groups that can be separated. Specifically, based on the characters of the phallic complex, the first species-group has more than one pair of processes of the phallic complex; the apex of the phallic complex has acute processes and the length of the process of the lateral part of the periandrium is longer than half the length of the phallic complex (*A. jangis* Medler, 1999, *A. servis* Medler, 1999, *A. taenia* Schmidt, 1904). The second species-group has one pair of processes of the phallic complex; the apex of the phallic complex is evenly rounded without process; the length of the process of lateral part of periandrium is slightly shorter than half length of the phallic complex (*A. fimbria*, *A. hainanensis*, *A. nodosa* Gerstaecker, 1895, *A. ocularia* sp. nov., *A. patefacta* sp. nov., *A. sphaerica* sp. nov., *A. ungulata* sp. nov.). The third species-group has one pair of processes of phallic complex; the apex of the phallic complex is evenly rounded without a process and the length of the process of the lateral part of the periandrium is longer than half the length of the phallic complex (*A. lurida* Melichar, 1902, *A. mucida*, *A. obscura*, *A. punctulata* sp. nov.). The first species-group is distributed in Indonesia, while the other two species-groups are distributed in China, India, Sri Lanka and Indonesia. Meanwhile, the pygofer of *A. taenia* in Indonesia is the most specialized one, approximately triangular; the pygofer of the other 13 species is nearly quadrilateral. Hence, we infer that the species of *Atracis* distributed in Indonesia maybe the most specialized one and the species of *Atracis* indicate a complex diversity. However, the phylogenetical relationships within *Atracis* are still not quite clear. In the future, it will be necessary to address the systematic analysis of *Atracis* from morphological, molecular and biogeographical approaches, and to clarify interspecific relationships within the genus.

Acknowledgements

We are sincerely grateful to Prof. J.R. Schrock (Department of Biological Sciences, Emporia State University, USA) for revising the manuscript. We would like to thank Prof. Masami Hayashi (Laboratory of Entomology, Tokyo University of Agriculture, Atsugi, Japan) for private communication and providing references. We also appreciate Senior Researcher Liu Sikong (Beijing Natural History Museum) and Prof. Cai Wanzhi (China Agricultural University, Beijing) for the loan of specimens. Special thanks go to Prof. Wang Yinglun for valuable comments and suggestion on the revision of the manuscript. We thank the anonymous reviewers for their time and helpful comments on an earlier draft of this paper. This study is supported by the National Natural Science Foundation of China (31420103911) and the Ministry of Science and Technology of China (2015FY210300, 2005DKA21402).

References

- Banks C.S. 1910. Rhynchota Palawanica. Part 2. Homoptera. *Philippine Journal of Science* 5: 33–55.
- Bourgoin T. 1993. Female genitalia in Hemiptera Fulgoromorpha, morphological and phylogenetic data. *Annales de la Société entomologique de France* 29 (3): 225–244.
- Bourgoin T. 2020. FLOW (Fulgoromorpha Lists on The Web): a world knowledge base dedicated to Fulgoromorpha. Version 8, updated [17 Feb. 2020]. Available from <https://hemiptera-databases.org/flow/> [accessed 17 Feb. 2020].
- Bourgoin T. & Huang J. 1990. Morphologie comparée des genitalia mâles des Trypetimorphini et remarques phylogénétiques (Hemiptera: Fulgoromorpha: Tropiduchidae). *Annales de la Société entomologique de France* 26 (4): 555–564.
- Bourgoin T., Wang R.R., Asche M., Hoch, H., Soulier-Perkins A., Stroiński A., Yap S. & Szwedo J. 2015. From micropterism to hyperpterism: recognition strategy and standardized homology-driven terminology of the forewing venation patterns in planthoppers (Hemiptera: Fulgoromorpha). *Zoomorphology* 134 (1): 63–77. <https://doi.org/10.1007/s00435-014-0243-6>
- Distant W.L. 1906. *The Fauna of British India, including Ceylon and Burma. Rhynchota. – Vol. III. (Heteroptera-Homoptera)*. Published under the authority of the Secretary of State for India in Council, London.
- Distant W.L. 1910. Rhynchota Malayana. Part III. *Records of the Indian Museum* 5: 313–338. <https://doi.org/10.5962/bhl.part.10506>
- Distant W.L. 1912. XLVIII.– New genera and species of Oriental Homoptera. *Annals and Magazine of Natural History, Series 8* 9 (52): 459–471. <https://doi.org/10.1080/00222931208693156>
- Distant W.L. 1914. XLVII.– Some additions to the genera and species in the Homopterous family Fulgoridae. *Annals and Magazine of Natural History, Series 8* 13 (76): 409–424. <https://doi.org/10.1080/00222931408693503>
- Fang S.J. 1989. Flatidae of Taiwan (Homoptera, Fulgoroidea). *Collected Papers on Homoptera of Taiwan, Taiwan Museum Special Publication* 8: 117–152.
- Fletcher M.J. 2009 and updates. Identification keys and checklists for the leafhoppers, planthoppers and their relatives occurring in Australia and neighbouring areas (Hemiptera: Auchenorrhyncha). Available from <https://idtools.dpi.nsw.gov.au/keys/auch/index.html> [accessed 8 Sep. 2020].
- Jacobi A. 1915. Kritische Bemerkungen über die Flatinae (Rhynchota Homoptera). *Berliner entomologische Zeitschrift* 1915 (2): 157–178. <https://doi.org/10.1002/mmnd.191519150206>
- Kato M. 1933. Notes on Japanese Homoptera, with descriptions of one new genus and some new species. *The Entomological World* 1: 452–471.

- Matsumura S. 1940. Homopterous insects collected at Kotosho (Botel Tobago) Formosa, by Mr. Tadao Kano. *Insecta Matsumurana* 15 (1–2): 34–51.
- Medler J.T. 1986. Types of Flatidae (Homoptera). XII. Taxonomic notes on Guerin-Meneville types in the Naples Museum, with illustrations of male genitalia of plesiotypes for the respective species. *Bollettino del Laboratorio di entomologia agraria Filippo Silvestri Portici* 43: 11–20.
- Medler J.T. 1988. Flatidae from the Tai Forest, Cote d'Ivoire, and taxonomic notes on the family in west Africa (Homoptera, Auchenorrhyncha, Fulgoroidea). *Revue Française d'entomologie* 10 (2): 117–148.
- Medler J.T. 1990. Review of *Jamella* Kirkaldy and *Malleja*, gen. nov. in Australia and New Guinea, with descriptions of new species (Homoptera: Flatidae). *Invertebrate Taxonomy* 3 (7): 995–1004. <https://doi.org/10.1071/IT9890995>
- Medler J.T. 1993. Types of Flatidae: 20. Lectotype designations and taxonomic notes on species in the MNHN Paris: Part 2. (Homoptera, Fulgoroidea). *Revue Française d'entomologie (Nouvelle Série)* 15 (2): 49–60.
- Medler J.T. 1996. Flatidae of Borneo, with descriptions of new genera and species (Homoptera: Fulgoroidea). *Oriental Insects* 30 (1): 11–96. <https://doi.org/10.1080/00305316.1996.10433829>
- Medler J.T. 1999. Flatidae (Homoptera: Fulgoroidea) of Indonesia, exclusive of Irian Jaya. *Zoologische Verhandelingen Leiden* 324: 1–88.
- Melichar L. 1902. Monographie der Acanaloniiden und Flatiden (Homoptera) (Fortsetzung). *Annalen des K.K. Naturhistorischen Hofmuseums* 17: 1–253.
- Metcalf Z.P. 1957. *General Catalogue of the Homoptera, Fascicle IV. Fulgoroidea, Part 13 Flatidae and Hypochthonellidae*. North Carolina State College, Raleigh.
- Stål C. 1862. Synonymiska och systematiska anteckningar öfver Hemiptera. *Öfversigt af Kongliga Vetenskaps-Akademiens Förfärlingar* 19: 479–504.
- Stål C. 1866. Hemiptera Homoptera Latr. *Hemiptera Africana* 4: 1–276.
- Stål C. 1870. Hemiptera insularum Philippinarum. Bidrag till Phillipinska oarues Hemiptera fauna. *Öfversigt af Kongliga Vetenskaps-Akademiens Förfärlingar* 27: 607–776.
- Walker F. 1851. *List of the Specimens of Homopterous Insects in the Collection of the British Museum, Part II*. British Museum Trustees, London. <https://doi.org/10.5962/bhl.title.9063>
- Walker F. 1858. *List of the Specimens of Homopterous Insects in the Collection of the British Museum, Supplement*. British Museum Trustees, London. <https://doi.org/10.5962/bhl.title.9063>
- Zia Y. 1935. Note sur les Flatinae et les Ricaniae de la Chine du sud et du Tonkin. *Sinensis* 6: 525–535.

Manuscript received: 17 November 2019

Manuscript accepted: 27 July 2020

Published on: 29 September 2020

Topic editor: Nesrine Akkari

Desk editor: Pepe Fernández

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the EJT consortium: Muséum national d'histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences,

Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Real Jardín Botánico de Madrid CSIC, Spain; Zoological Research Museum Alexander Koenig, Bonn, Germany; National Museum, Prague, Czech Republic.