

Research article

[urn:lsid:zoobank.org:pub:08112A58-AF64-4B0B-B603-41B94E88FA4B](https://zoobank.org/pub:08112A58-AF64-4B0B-B603-41B94E88FA4B)**Two new species of *Sarinda* Peckham & Peckham, 1892,
with an update on *Sarindini* in Uruguay (Araneae: Salticidae)**Damián HAGOPIÁN^{1,*}, Abel A. BUSTAMANTE², Álvaro LABORDA³ & Miguel SIMÓ⁴^{1,3,4}Sección Entomología, Facultad de Ciencias, Universidad de la República,
Iguá 4225, PC 11400, Montevideo, Uruguay.²Museu Paraense Emílio Goeldi, Coordenação de Zoologia, Laboratório de Aracnologia,
Av. Perimetral, 1901, Terra Firme, CEP 66077–830, Belém, Pará, Brazil.* Corresponding autor: dhagopian@fcien.edu.uy²Email: a.bustamanteferrada@gmail.com³Email: alaborda@fcien.edu.uy⁴Email: simo@fcien.edu.uy¹[urn:lsid:zoobank.org:author:B93E6659-7528-4230-B272-D3EBDAAFD5BB](https://zoobank.org/author:B93E6659-7528-4230-B272-D3EBDAAFD5BB)²[urn:lsid:zoobank.org:author:AB7BF4CC-3277-4376-99B9-8F12D9C00E6E](https://zoobank.org/author:AB7BF4CC-3277-4376-99B9-8F12D9C00E6E)³[urn:lsid:zoobank.org:author:DA4C0AC7-9EDB-497A-9BEC-197DA843ED9C](https://zoobank.org/author:DA4C0AC7-9EDB-497A-9BEC-197DA843ED9C)⁴[urn:lsid:zoobank.org:author:8967ABC7-0EF3-48F0-AFC4-4F29E8D8BCD4](https://zoobank.org/author:8967ABC7-0EF3-48F0-AFC4-4F29E8D8BCD4)

Abstract. *Sarinda sombraluminosa* Hagopían, Laborda & Simó sp. nov. and *Sarinda contraluz* Hagopían & Bustamante sp. nov. are described from Uruguay based on males and females. New records of *Parafтуда banksi* Chickering, 1946 and *Sarinda marcosi* Piza, 1937 for the country are provided. Illustrations and electron micrographs of sexual characters, photographs of alive specimens, natural history data and a distribution map of species of *Sarandini* from Uruguay are provided.

Keywords. Ant-like, Salticinae, Amycoidea, *Parafтуда*.

Hagopían D., Bustamante A., Laborda Á. & Simó M. 2024. Two new species of *Sarinda* Peckham & Peckham, 1892, with an update on *Sarindini* in Uruguay (Araneae: Salticidae). *European Journal of Taxonomy* 925: 100–134. <https://doi.org/10.5852/ejt.2024.925.2455>

Introduction

The species of the jumping spider tribe *Sarindini* Simon, 1901 are relatively poorly known. These jumping spiders of the Americas are characterized by mimicking ants in behavior and morphology (Galiano 1965, 1967, 1971), likely representatives of the genera *Formica* Linnaeus, 1758 and *Camponotus* Mayr, 1861 (Maddison 2015). A recent study on *Sarinda marcosi* Piza, 1937 revealed that the ant-like model of this species is *Camponotus mus* Roger, 1863 based on the defensive mimicry (Hagopían *et al.* 2021).

Currently, the *Sarindini* is composed of 45 species divided into seven genera: *Corcovetella* Galiano, 1975; *Martella* Peckham & Peckham, 1892; *Parafтуда* Chickering, 1946; *Sarinda* Peckham & Peckham, 1892; *Simprulla* Simon, 1901; *Tanybelus* Simon, 1902 and *Zuniga* Peckham & Peckham,

1892 (Maddison 2015; World Spider Catalog 2023). The richest genus in this tribe is *Sarinda*, with 16 described species, followed by *Martella* (12 species) and *Corcovetella*, *Simprulla* and *Zuniga* (two species each). *Parafluda* and *Tanybelus* are monotypic (World Spider Catalog 2023).

According to Galiano (1965, 1996), the genus *Sarinda* resembles *Martella*, but males of *Sarinda* do not have the proximal ectal apophysis on the cymbium, which is present in males of *Martella*. Additionally, females of *Sarinda* present two main groups, based on genitalic patterns. One of these groups (“group 1”) presents two pairs of spermathecae and the spiraled median copulatory ducts (common also in “group 2”), while those of *Martella* always have one pair of spermathecae and a membranous compartment from which spermathecal and glandular ducts originate (see Galiano 1964). On the other hand, Galiano’s “group 2” of *Sarinda* presents the copulatory duct slightly dilated, not forming a second pair of spermathecae (Galiano 1964).

From recent surveys in Uruguay, we found specimens that agree with the diagnosis proposed for “group 1” of *Sarinda* (see Galiano 1965: 268) and do not correspond with the currently known species of this genus. Also, we found new records of other species for its tribe. Therefore, the aim of this study is to describe two new species of *Sarinda* from Uruguay, as well as to provide new data on the distribution and natural history of the species of Sarandini in the country.

Material and methods

Specimens are deposited in the arachnological collection of the Facultad de Ciencias, Universidad de la República (UdelaR), Montevideo, Uruguay (FCE-Ar, M. Simó) and the Spencer Entomological Collection, Beaty Biodiversity Museum, Vancouver, Canada (UBCZ, W. Maddison). Measurements are in millimeters. Total length includes anterior median eyes and anal tubercle (Edwards 2004). Color images were obtained using a Leica M205 A stereo microscope, attached to a Leica DMC 2900 camera enabled with the Leica LAS-X-Z and SW software. Photographs of female genitalia were taken with a Nikon D3500 digital camera attached to a microscope and images were stacked using Helicon Focus 7 ver. 7.6.4 Lite software. Also, we took electron micrographs of somatic and genital features with a JEOL 5900 Scanning Electron Microscope from the Servicio de Microscopía Electrónica de Barrido, Facultad de Ciencias, Universidad de la República (UdelaR). Drawings were made using a Wacom Intuos Pro pen tablet and SketchBook ver. 8.7.1 illustration software (<https://sketchbook.com>) following Cala-Riquelme (2021). Female genitalia were cleaned in a solution of trypsin for the digestion of soft tissues and then cleared using clove oil (Levi 1965). Length of embolus is approximated in degrees, starting from the base of the embolus and ending at the part of the embolus where the terminal coil starts (this coil is often at the tip of the embolus); the letter T (turn) in embolus description equals 360°; the length of the embolus was not estimated when the palp was distended (Bustamante & Ruiz 2017), the position of the embolus base is expressed using the position of the hours of the clock (Bustamante & Ruiz 2020). Leg spines are described as in Petrunkevitch (1925), with the modifications of Bustamante & Ruiz (2017); only for femora, patellae, tibiae and metatarsi. In general, ventral spines of tibia are equal size; when not, this is marked with an asterisk (*). In vivo pictures and videos were taken with an Olympus Tough Tg-4 digital camera. Geographic coordinates were taken directly from the labels. In cases when the labels did not give information about geographic coordinates, these were approximated with Google Earth® and marked with square brackets; in cases when an error was detected in the name of a locality, it was corrected and marked with square brackets. The distribution map was created using Simplemappr (Shorthouse 2010).

Abbreviations

Depository institution and curator

- FCE-Ar = Arachnological Collection of Facultad de Ciencias, Universidad de la República, Montevideo, Uruguay (M. Simó)
 UBCZ = Spencer Entomological Collection, Beaty Biodiversity Museum, Vancouver, Canada (W. Maddison)

Morphology

ac	=	aciniiform gland
alS	=	anterior lateral spinneret
ap	=	apical
CD	=	copulatory duct
CO	=	copulatory opening
d	=	dorsal
E	=	embolus
FD	=	fertilization duct
gl	=	gland
ma	=	mastidion
mAP	=	minor ampullate spigot
MAP	=	major ampullate spigot
p	=	prolateral
pi	=	piriform gland
plE	=	posterior lateral eyes
plS	=	posterior lateral spinneret
pmS	=	posterior median spinneret
PS	=	primary spermatheca
r	=	retrolateral
RTA	=	retrolateral tibial apophysis
RvTA	=	retroventral tibial apophysis
S	=	spermatheca
SS	=	secondary spermatheca
v	=	ventral

Results

Class Arachnida Cuvier, 1812
Order Araneae Clerck, 1757
Family Salticidae Blackwall, 1841
Subfamily Salticinae Blackwall, 1841
Tribe Sarindini Simon, 1901

Genus *Sarinda* Peckham & Peckham, 1892

Type species

Sarinda nigra Peckham & Peckham, 1892.

Sarinda sombraluminosa Hagopían, Laborda & Simó sp. nov.
[urn:lsid:zoobank.org:act:F9B935EA-2DCE-4E3C-92C5-A815FCA8B9CB](https://zoobank.org/urn:lsid:zoobank.org:act:F9B935EA-2DCE-4E3C-92C5-A815FCA8B9CB)
Figs 1–12, 33

Diagnosis

Among *Sarinda* with an embolus longer than 2T, *S. sombraluminosa* Hagopían, Laborda & Simó sp. nov. can be distinguished from *S. nigra*, *S. capibarae* Galiano, 1967, *S. silvatica* Chickering, 1946, *S. marcosi*, and *S. panamae* Galiano, 1965 by having a dorsally curved RvTA and a shorter embolus ($3T+90^\circ$) (9T or 10T in *S. nigra*, see Galiano 1965: 301, fig. 1; 8T or 9T in *S. capibarae*, see Galiano 1967: 32, fig. 21; 6T or 7T in *S. silvatica*, see Galiano 1965: 301, fig. 4; 4T or 5T in *S. marcosi*, see Galiano 1965: 304, fig. 1 and Hagopían *et al.* 2021: 265, fig. 1e; 4T in *S. panamae*, see Galiano 1965: 304, fig. 2); males of

S. sombraluminosa can be distinguished of *S. ruficeps* (Simon, 1901) by having a curved RvTA pointing to the bulb and longer embolus ($2T+90^\circ$ in *S. ruficeps*, see Müller & Cutler 1989: 75, fig. 5) (Figs 4A–B, 5A–B, 6A–D). Among *Sarinda* with two pairs of spermathecae, *S. sombraluminosa* resemble those of *S. capibarae*, but can be distinguished from those of *S. capibarae* by having shorter copulatory ducts and an internal duct that connects the secondary spermatheca with the primary having four turns around the gland (16 in *S. capibarae*, see Galiano 1967: 32, fig. 23) (Figs 4E–F, 5E–F, 7A–B).

Etymology

The specific epithet is a compound noun in apposition that means ‘luminous shadow’ in Spanish, because of the contrasting colors of white scales on the chelicerae of the male and on the dark body of both sexes of the species.

Type material

Holotype

URUGUAY • ♂; Montevideo, Melilla; 34.73194° S, 56.32218° W; 18 Apr. 2020; D. Hagopían leg.; in a pile of grass cut with a tractor rotary cutter in grassland; FCE-Ar 13551.

Paratypes

URUGUAY • 1 ♀; same collection data as for holotype; FCE-Ar 5896 • 5 ♂♂, 7 ♀♀; same collection data as for holotype; FCE-Ar 5896.

Other material examined

URUGUAY – **Durazno** • 1 ♂, 3 ♀♀; San Eduardo; 32.59222° S, 55.71194° W; 29 Jan. 2019; G. Pompozzi leg.; FCE-Ar 11400 • 2 ♂♂, 2 ♀♀; same locality as for preceding; 12 Aug. 2019; G. Pompozzi leg.; FCE-Ar 11259. – **Flores** • 1 ♂, 1 ♀; Rincón de Piedra; 33.87750° S, 56.98833° W; 2 May 2019; G. Pompozzi leg.; in grassland; FCE-Ar 11393 • 1 ♀; same locality as for preceding; 21 Oct. 2019; G. Pompozzi leg.; FCE-Ar 11241. – **Montevideo** • 1 ♀; Melilla; 34.73261° S, 56.32110° W; 18 Feb. 2018; D Hagopían leg.; walking on outside wall; FCE-Ar 11214 • 1 ♂, same locality as for preceding; 8 Apr. 2018; D. Hagopían leg.; FCE-Ar 9010 • 1 ♂; same locality as for preceding; 15 May 2018; D. Hagopían leg.; FCE-Ar 9462 • 1 ♂; same locality as for preceding; 9 Dec. 2022; D. Hagopían leg.; FCE-Ar 14008. – **Río Negro** • 1 ♂; Ruta 24 Km 85, Estancia “Las Cadenas”; 32.52742° S, 58.03322° W; 30 Abr. 2020; A. Mailhos leg.; walking on fence (in grassland); FCE-Ar 13561 • 1 ♂; same locality as for preceding; 32.52706° S, 58.03436° W; 7 May 2020; A. Mailhos leg.; walking on fence (in grassland); FCE-Ar 13550 • 1 ♂; same locality as for preceding; 5 Jan. 2021; A. Mailhos leg.; walking on fence (in grassland); FCE-Ar 13778 • 1 ♂; same locality as for preceding; 32.53036° S, 58.03533° W; 1 May 2021, A. Mailhos leg.; walking on fence (in grassland); FCE-Ar 13563. – **Rivera** • 3 ♂♂, 2 ♀♀; Mi Lucha; 31.43711° S, 55.27094° W; 10 Jul. 2019; G. Pompozzi leg.; in grassland; FCE-Ar 11251 • 1 ♂; Ypoá; 31.73139° S, 55.56972° W; 14 Oct. 2019; G. Pompozzi leg.; in grassland; FCE-Ar 11246 • 1 ♀; same locality as for preceding; 9 Jul. 2019; G. Pompozzi leg.; in grassland; FCE-Ar 11408 • 1 ♀; same locality as for preceding; 12 Feb. 2020; G. Pompozzi leg.; vFCE-Ar 11414.

Description

Male (holotype FCE-Ar 13551)

COLOR. Carapace black with white scales on face and on back of cephalic constriction. Abdomen as carapace, with slight dorsal constriction in middle and with transversal band of white scales, some of those scales scattered in lower density in posterior region of abdomen (Figs 1D, F, 2A–C, 9A–C). Ventrally black, sternum narrow as shown in Fig. 9D. Spinnerets dark brown. Palps dark brown. Legs I and II ventrally and dorsally yellowish, with lateral sides brown, being only the tarsus of leg I dark brown. Legs III with coxae, trochanter, femur and patella dark brown, tibiae, metatarsus and tarsus as




Fig. 1. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., photographs in vivo. **A.** Site where the species was collected. **B, D, F.** Male. **C, E, G.** Female.

leg II. Leg IV darker, with yellowish coxa and patella, rest of segments dark brown (Figs 1B, D, F, 2A–C, 3A, C). Trichobothria present in tarsus of leg I as shown in Fig. 9E. Cheliceral paturon dark brown with four promarginal teeth (distal one being bigger) and one bigger, not curved, and flat retromarginal tooth. Mastidion near base of claw (Fig. 10A). White spoon-shaped scales homogeneously scattered all over anterior surface of paturon (Figs 1B, D, 3A, 10).

MEASUREMENTS. Total length 3.80. Carapace length 2.0, width 1.10, height 1.0. AME diameter 0.35. Ocular quadrangle length 0.90. Anterior eye row width 1.0. Posterior eye row width 0.90. Abdomen length 1.80.

LEGS. Leg I: femur 1.0, patella 0.50, tibia 1.10, metatarsus 0.70, tarsus 0.40; II: fe 0.80, pa 0.40, ti 0.70, mt 0.55, ta 0.30; III: fe 0.80, pa 0.45, ti 0.70, mt 0.70, ta 0.30; IV: fe 1.40, pa 0.40, ti 1.50, mt 1.50,



Fig. 2. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., habitus. A–C. Holotype, ♂ (FCE-Ar 13551). D–F. Paratype, ♀ (FCE-Ar 5896). A, D. Dorsal view. B, E. Lateral view. C, F. Ventral view.

ta 0.35. Leg formula 4132. Leg macrosetae: femur and patella I–IV d0, p0, r0, v0; tibia I v2-2-2; II v1r-1r-1r; III 0, IV v1p; metatarsus I–II v2-2; III v2-1r, p 2ap, r 2ap; IV v2-1r2, p1ap, r2ap, d1p-1p-2.

PALP. Tibia longer than wide; RTA tooth like, curved dorsally, longer than RvTA. RvTA thumb-like, curved ventrally, embolus simple, fixed to tegulum, arising distally (11:00) ($3T+90^\circ$) (Figs 4A–D, 5A–D, 6).

ABDOMEN. Male without epiandrous fusules (Fig. 7C–D).

Female (paratype FCE-Ar 5896)

COLOR. As in male (Figs 1C, E, G, 2D–F, 3B, D). Modified scales present along constriction between cephalic and thoracic regions (Fig. 9A–C).

MEASUREMENTS. Total length 4.50. Carapace length 2.10, width 1.0, height 0.90. AME diameter 0.30. Ocular quadrangle length 0.80. Anterior eye row width 1.0. Posterior eye row width 1.0. Abdomen length 2.40.



Fig. 3. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., face and chelicera. **A, C.** Holotype, ♂ (FCE-Ar 13551). **B, D.** Paratype, ♀ (FCE-Ar 5896). **A–B.** Face. **C–D.** Chelicera.

LEGS. Leg I: femur 1.0, patella 0.38, tibia 1.0, metatarsus 0.50, tarsus 0.40. II: fe 0.80, pa 0.45, ti 0.60, mt 0.50, ta 0.38. III: fe 1.0, pa 0.50, ti 0.65, mt 0.60, ta 0.30. IV: fe 1.40, pa 0.50, ti 1.30, mt 1.0, ta 0.40. Leg formula 4132. Leg macrosetae: femur I–II d1-1-0, p0, r0; III d0-1-0 (d0-1-1 right), p0, r0; IV d1-1-1, p0, r0; patella I–IV 0; tibia I p0, r0, v2*-2a-0; II p0, r0, v1r-1r-0; III–IV 0; metatarsus I p0, r0, v2-2; II p0, r0, v1r-2; III–IV 0.

PALP. Tibiae wider than cymbium, both with saber-like setae (Fig. 8). Chelicerae paturon with five promarginal teeth and one bigger and flat retromarginal tooth (Fig. 3D).

EPIGYNE. Atrium horse-shoe like, slightly sclerotized with narrow copulatory openings. Posterior margin bilobed with constriction reaching entrance of atrium (Figs 4E, 5E, 7A). Spermatheca can be seen through cuticle, secondary (posterior ones) bigger. Internally, spermatheca (anterior) touching each other, connected with slightly spiraled duct with four turns around gland to secondary spermatheca (posterior). Internal gland thicker than inner duct and head almost touching primary spermathecae. Copulatory ducts membranous with five turns before connecting with secondary spermatheca (Figs 4F, 5F, 7B).




Fig. 4. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., photographs of genitalia. A–D. Holotype, ♂ (FCE-Ar 13551), pedipalp. A. Ventral view. B. Retrolateral view. C. Dorsal view. D. Prolateral view. E–F. Paratype, ♀ (FCE-Ar 5896), epigynum. E. Ventral view. F. Dorsal view.

SPINNERETS. Posterior lateral spinnerets with aciniform glands, posterior median spinnerets with aciniform and minor ampullate glands and anterior lateral spinnerets with piriform glands and major ampullate glands (Fig. 12).

Natural history

This species was found in bushes, herbaceous vegetation and on fences placed in grasslands (Fig. 1A), collected by hand and G-Vac vacuum. It has been observed in the same foraging places as the ant *Camponotus termitarius* Emery, 1902. In the laboratory, the specimens were fed different species of *Drosophila* Fallén, 1823. Specimens were recorded lifting the first pair of legs and moving the pedipalps up and down when walking (<https://www.youtube.com/watch?v=obBZqjP1b8I>) and eating a fly (<https://www.youtube.com/watch?v=tma8CZ6OCOc>).




Fig. 5. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., drawings of genitalia. A–D. Holotype, ♂ (FCE-Ar 13551), pedipalp. A. Ventral view. B. Retrolateral view. C. Dorsal view. D. Prolateral view. E–F. Paratype, ♀ (FCE-Ar 5896), epigynum. E. Ventral view. F. Dorsal view. Abbreviations: CD = copulatory duct; CO = copulatory opening; E = embolus; FD = fertilization duct; gl = gland; PS = primary spermatheca; RTA = retrolateral tibial apophysis; RvTA = retroventral tibial apophysis; SS = secondary spermatheca.




Fig. 6. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., paratype, ♂ (FCE-Ar 5896), scanning electronic microphotography (SEM) of pedipalp. **A.** Ventral view. **B.** Retrolateral view, arrow indicating paracymbium. **C.** Retroventral tibial apophysis, ventral view. **D–E.** RvTA, retrolateral view, with an arrow indicating the paracymbium. Abbreviations: E = embolus; RTA = retrolateral tibial apophysis; RvTA = retroventral tibial apophysis; T = tegulum.




Fig. 7. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., SEM of genitalia. **A–B.** Paratype, ♀ (FCE-Ar 5896), epigynum. **A.** Ventral view. **B.** Dorsal view. **C–D.** Paratype, ♂ (FCE-Ar 5896), epigastric male zone, showing the absence of epiandric spigots. Abbreviations: CD = copulatory duct; CO = copulatory opening; SS = secondary spermatheca.




Fig. 8. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., paratype, ♀ (FCE-Ar 5896), SEM of female palp. **A.** General view, upper arrow pointing to distal macrosetae, lower pointing to the abundance of setae in tibia and cymbium. **B.** Distal macrosetae. **C–D.** Saber-shaped macrosetae details.




Fig. 9. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., SEM. **A–C.** Paratype, ♀ (FCE-Ar 5896). **A.** Prosoma general view, arrow pointing to scales in the constriction between cephalic and thoracic regions. **B–C.** Arrangement and details of scales. **D–E.** Paratype, ♂ (FCE-Ar 5896). **D.** Sternum. **E.** Tarsus, arrows pointing to trichobothria.




Fig. 10. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., paratype, ♂ (FCE-Ar 5896), SEM of left chelicerae. **A.** Chelicerae general view, black arrows pointing to the teeth of the promargin and retromargin, white arrows pointing to scales. **B.** Scale details. **C.** Teeth of the promargin and retromargin.




Fig. 11. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., paratype, ♂ (FCE-Ar 5896), SEM of opisthosoma. **A.** General view, white arrows pointing to scales. **B–D.** Scale details.




Fig. 12. *Sarinda sombraluminosa* Hagopián, Laborda & Simó sp. nov., paratype, ♀ (FCE-Ar 5896), SEM of spinnerets. **A.** General view. **B.** Anterior lateral spinneret. **C.** Posterior median spinneret. **D.** Posterior lateral spinneret. Abbreviations: ac = aciniform gland; alS = anterior lateral spinneret; mAP = minor ampullate gland; MAP = major ampullate gland; pi = piriform spigot; pIS = posterior lateral spinneret; pmS = posterior medial spinneret.

Taxonomic remarks

This species is mentioned as “*Sarinda* sp.1” in Hagopián & Laborda (2020), who recorded it as a host of mantispid larvae. Also, this species was misidentified as *S. capibarae* in Pompozzi *et al.* (2022: table s1 in supplementary material).

Distribution

Known from Uruguay (Durazno, Flores, Montevideo, Río Negro and Rivera) (Fig. 33).

Sarinda contraluz Hagopián & Bustamante sp. nov.

[urn:lsid:zoobank.org:act:A57C501C-8E3E-4D0B-B47A-8FF1C8DBF1DA](https://zoobank.org/act:A57C501C-8E3E-4D0B-B47A-8FF1C8DBF1DA)

Figs 13–24, 33

Diagnosis

Among *Sarinda* with shorter embolus (less than 2T compared with *S. armata* Peckham & Peckham, 1892, *S. hentzi* (Banks, 1903), *S. cutleri* (Richman, 1965), *S. imitans* Galiano, 1965), males of *S. contraluz* Hagopián & Bustamante sp. nov. resemble those of *S. imitans* by having a mastidion on the male chelicera (see Fig. 22A) and by the shape of the RTA (Figs 17C, 18B, 18D), but can be distinguished from those of *S. imitans* by having a developed RvTA and a shorter pedipalp tibia, a longer embolus that arises at 5:30 (8:30 in *S. imitans*, see Galiano 1967: 30, fig. 11), by having a more rounded tegulum (Figs 16–18), and for the retromarginal teeth more developed, not so close to each other, and with the




Fig. 13. *Sarinda contraluz* Hagopián & Bustamante sp. nov., photographs in vivo **A.** Site where the species was collected. **B, D, F.** Male. **C, E, G.** Female.

distal one larger than the others (compare Fig 15C with Galiano 1967: fig. 12). Among *Sarinda* with two pairs of spermathecae, females of *S. contraluz* resemble those of *S. imitans* by having a membranous portion of the copulatory ducts with less than three turns (two to three in *S. imitans*, see Galiano 1965: 289, fig. 8; two turns in *S. contraluz*), but can be distinguished by having more developed and longer copulatory ducts and the internal duct connecting the secondary spermatheca with the primary is less spiraled (two to three in *S. imitans*, see Galiano 1965: 294, fig. 3; two in *S. contraluz* see Figs 16–17, 19).




Fig. 14. *Sarinda contraluz* Hagopían & Bustamante sp. nov., habitus. A–C. Holotype, ♂ (FCE-Ar 10509). D–F. Paratype, ♀ (FCE-Ar 13968). A, D. Dorsal view. B, E. Lateral view. C, F. Ventral view.

Etymology

The specific name is a compound noun in apposition that means ‘backlighting’ in Spanish, because of the presence of the thin transparent-white setae on the abdomen of the species, which can be observed more clearly at backlighting.

Type material

Holotype

URUGUAY • ♂; Montevideo, Melilla; 34.73250° S, 56.32083° W; 5 Nov. 2022; D. Hagopían and A. Mailhos leg.; collected with G-Vac vacuum in *Cortaderia selloana* (Schult. & Schult.f.) Asch. & Graebn. (Poaceae); FCE-Ar 10509.

Paratype

URUGUAY • 1 ♀; same collection data as for holotype; FCE-Ar 13968.




Fig. 15. *Sarinda contraluz* Hagopían & Bustamante sp. nov., face and chelicera. **A, C.** Holotype, ♂ (FCE-Ar 10509). **B, D.** Paratype, ♀ (FCE-Ar 13968). **A–B.** Face. **C–D.** Chelicera.

Other material examined

URUGUAY. – **Canelones** • 1 ♂, 2 ♀♀; Barra de Carrasco; 34.87700° S, 56.02278° W: 10 Dec. 2005; Maddison, Ruiz, Simó, Rodríguez and Laborda leg.; WPM#05-049; UBCZ. – **Lavalleja** • 1 ♀; Sierra de Minas, Parque de Vacaciones; 34.42583° S, 55.19500° W; 5–8 Dec. 2005; W. Maddison, G. Ruiz, M. Simó and M.E. Rodríguez leg.; WPM#05-046; UBCZ. – **Montevideo** • 1 ♂; Carrasco; 34.87972° S, 56.03056° W; 9 Oct. 2022; D. Hagopían and A. Mailhos leg.; FCE-Ar 13925 • 3 ♀♀, 2 ♂♂; same locality as for preceding; 19 Nov. 2022; A. Mailhos leg.; FCE-Ar 13981 • 8 ♂♂, 2 ♀♀; same collection data as for holotype; FCE-Ar 13968 • 1 ♂ same locality as for holotype; 20 Nov. 2022; D. Hagopían leg.; FCE-Ar 11414. – **Rivera** • 1 ♂; Cofusa; 30.99861° S, 55.67500° W; 23 Nov. 2021; A. Mailhos leg.; in *Paspalum* sp. (Poaceae) on a truncated hill; FCE-Ar 13791.

Description

Male (holotype FCE-Ar 10509)

COLOR. Carapace black with scales on back of cephalic constriction, abdomen as carapace, with thin transparent-white setae all over (Fig. 13B, D, F). Ventrally black, sternum shown in Fig. 21D. Spinnerets




Fig. 16. *Sarinda contraluz* Hagopían & Bustamante sp. nov., photographs of genitalia. **A–D.** Holotype, ♂ (FCE-Ar 10509), pedipalp. **A.** Ventral view. **B.** Retrolateral view. **C.** Dorsal view. **D.** Prolateral view. **E–F.** Paratype, ♀ (FCE-Ar 13968), epigynum. **E.** Ventral view. **F.** Dorsal view.

dark brown. Palps brown. Legs I ventrally and dorsally yellowish, with lateral sides brown. Legs II as legs I, being brown only in prolateral side. Legs III with coxae, trochanter, femur and patella brown, tibiae, metatarsus and tarsus as legs I. Leg IV darker, with yellowish coxa, trochanter and patella, rest of articles dark brown (Figs 13B, D, F, 14A–C, 15A). Trichobothria present in metatarsus of leg I as shown in Fig. 21E.

MEASUREMENTS. Total length 4.460. Carapace length 2.156, width 1.168, height 0.869. AME diameter 0.352. Ocular quadrangle length 1.00. Anterior eye row width 1.062. Posterior eye row width 1.106. Abdomen length 2.079. Chelicera paturon light brown with four promarginal teeth (distal one being bigger) and three retromarginal teeth (distal one being bigger) (Figs 15A, C, 22). Dorsal mastidion at middle of base of claw (Fig. 22A). Setae white and scarce, being mostly at base of paturon (Figs 15A, 22).




Fig. 17. *Sarinda contraluz* Hagopián & Bustamante sp. nov., drawings of genitalia. **A–D.** Holotype, ♂ (FCE-Ar 10509), pedipalp. **A.** Ventral view. **B.** Retrolateral view. **C.** Dorsal view. **D.** Prolateral view. **E–F.** Paratype, ♀ (FCE-Ar 13968), epigynum. **E.** Ventral view. **F.** Dorsal view. Abbreviations: CD = copulatory duct; CO = copulatory opening; E = embolus; FD = fertilization duct; gl = gland; PS = primary spermatheca; RTA = retrolateral tibial apophysis; RvTA = retroventral tibial apophysis; S = spermatheca; SS = secondary spermatheca.




Fig. 18. *Sarinda contraluz* Hagopíán & Bustamante sp. nov., non-type specimen, ♂ (FCE-Ar 13968), SEM of male pedipalp. **A.** Ventral view. **B.** Retrolateral view. **C.** RvTA, ventral view. **D.** Retrolateral view, white arrow showing the paracymbium. Abbreviations: E = embolus; RTA = retrolateral tibial apophysis; RvTA = retroventral tibial apophysis; T = tegulum.




Fig. 19. *Sarinda contraluz* Hagopíán & Bustamante sp. nov., paratype, ♀ (FCE-Ar 13968), SEM of genitalia. **A–C.** Epigynum. **A.** Ventral view. **B.** Dorsal view. **C.** Dorsal view, white arrow pointing to the pore of the gland. **D.** Spermatheca, white arrow pointing to the pore of the bennet gland. Abbreviations: CD = copulatory duct; CO = copulatory opening; FD = fertilization duct; PS = primary spermatheca; S = spermatheca; SS = secondary spermatheca.




Fig. 20. *Sarinda contraluz* Hagopián & Bustamante sp. nov., paratype, ♀ (FCE-Ar 13968), SEM of palp. **A.** General view, white arrow pointing to the abundance of setae in the tibia. **B–C.** Saber-shaped macrosetae details.




Fig. 21. *Sarinda contraluz* Hagopián & Bustamante sp. nov., SEM. **A–C.** Paratype, ♀ (FCE-Ar 13968). **A.** Prosoma general view, arrow pointing to scales in the constriction between cephalic and thoracic regions. **B–C.** Arrangement and details of macrosetae. **D–E.** Non-type specimen, ♂ (FCE-Ar 13968). **D.** Sternum. **E.** Metatarsus, arrows pointing to trichobothria.




Fig. 22. *Sarinda contraluz* Hagopián & Bustamante sp. nov., non-type specimen, ♂ (FCE-Ar 13968), SEM of chelicerae. **A.** Chelicerae dorsal view, black arrows pointing to the teeth of the promargin and retromargin, white arrows pointing to macrosetae. **B.** Chelicerae ventral view, black arrow pointing to the retromargin teeth. **C.** Macrosetae details. **D.** Teeth of the promargin and retromargin.




Fig. 23. *Sarinda contraluz* Hagopián & Bustamante sp. nov., non-type specimen, ♂ (FCE-Ar 13968); SEM of opisthosoma. **A.** General view, white arrows pointing to macrosetae. **B–D.** Setae details.

LEGS. Leg I: femur 1.024, patella 0.440, tibia 1.142, metatarsus 0.643, tarsus 0.369; II: fe 0.847, pa 0.317, ti 0.661, mt 0.651, ta 0.229; III: fe 0.869, pa 0.373, ti 0.662, mt 0.606, ta 0.400; IV: fe 1.303, pa 0.502, ti 1.162, mt 1.036, ta 0.328. Leg formula 4132. Leg macrosetae: femur and patella I–IV d0, p0, r0, v0; tibia I v2-2-2-2, II v1r-1r-1r-1r, III v2ap, p0-1-0, IV v2ap; metatarsus I–II v2-2, III v2-2-0, p 2ap, r 1v-2ap; IV v2-2, p 1ap, r 1-0-1.

PALP. Tibia longer than wide; RTA bifurcated, curved ventrally, longer than RvTA. RvTA like a lump, embolus simple, fixed to tegulum, arising ventrally (6:00) (280°) (Figs 17A–D, 18).

Female (paratype FCE-Ar 13968)

COLOR. As in male (Figs 13C, E, G, 14D–F, 15B). Modified scales present on constriction between cephalic and thoracic regions (Fig. 21A–C).

MEASUREMENTS. Total length 4.68. Carapace length 2.13, width 1.00, height 0.78. anterior median eye diameter 0.36. Ocular quadrangle length 0.88. Anterior eye row width 1.0. Posterior eye row width 1.07. Abdomen length 2.32.

LEGS. Leg I: femur 1.10, patella 0.49, tibia 1.07, metatarsus 0.65, tarsus 0.38. II: fe 0.68, pa 0.41, ti 0.65, mt 0.46, ta 0.31. III: fe 0.94, pa 0.45, ti 0.69, mt 0.60, ta 0.34. IV: fe 1.19, pa 0.54, ti 1.21, mt 1.05, ta 0.43. Leg formula 4132. Leg macrosetae: femur and patella I–IV d0, p0, r0, v0; tibia I v2-2-2-2-2, II v1r-1r-1r, III d0, p0, v0, IV v2ap; metatarsus I–II v2-2, III v2ap-1p, p 1ap, r 1ap; IV v0-0-2, p 1ap, r 2ap.




Fig. 24. *Sarinda contraluz* Hagopián & Bustamante sp. nov., paratype, ♀ (FCE-Ar 13968), SEM of spinnerets. **A.** General view. **B.** Anterior lateral spinneret. **C.** Posterior median spinneret. **D.** Posterior lateral spinneret. Abbreviations: ac = aciniform gland; aLS = anterior lateral spinneret; mAP = minor ampullate spigot; MAP = major ampullate spigot; pi = piriform gland; pLS = posterior lateral spinneret; pmS = posterior median spinneret.

PALP. Tibiae wider than tarsus, with saber-like setae (Fig. 20). Chelicerae paturon with four promarginal teeth, three retromarginal teeth on right and four on left chelicerae (Fig. 15D).

EPIGYNE. Atrium slightly sclerotized with narrow copulatory openings. Posterior margin bilobated, septum with slightly blunt projection (Figs 16E, 17E, 19A). Spermatheca can be seen through cuticle, posterior ones being bigger. Internally, primary spermatheca touching each other connected with slightly spiraled duct to secondary spermatheca, being more separated and with wider wall. Pore of Bennett's gland situated apically. Internal gland thin and sinuous, reaching middle distance between spermathecae. Copulatory ducts membranous with two turns before connecting with secondary spermatheca (Figs 17F, 19B–D).

SPINNERETS. Anterior lateral spinnerets with piriform glands and major ampullate glands, posterior median spinnerets with aciniform and minor ampullate glands, and posterior lateral spinnerets with aciniform glands (Fig. 24).

Variation

Males (n = 10)

Total length (3.66–4.91), prosoma (1.87–2.46), abdomen (1.79–2.45), leg I (3.14–4.11), leg II (1.99–2.82), leg III (2.23–2.74), leg IV (3.47–4.67). Cheliceral teeth variation: four teeth in promargin, three in retromargin (n = 8, in one case the apical tooth of promargin was bifurcated, and in another two cases the apical tooth of retromargin was bifurcated); four teeth in promargin, three in the retromargin of right chelicera and two in the left (n = 1); four teeth in promargin (apical tooth bifurcated), three in the retromargin of right chelicera and four in left (n = 1).

Females (n = 4)

Total length (4.21–4.43), prosoma (2.08–2.20), abdomen (2.09–2.23), leg I (2.71–3.40), leg II (2.12–2.65), leg III (2.35–2.73), leg IV (4.04–4.47). Cheliceral teeth variation on margins: four teeth in promargin and three in retromargin (n = 2); four teeth in promargin, four in retromargin of right chelicera and three in left (n = 1); four teeth in promargin and four in retromargin (n = 1).

Natural history

This species was found primarily in *Cortaderia selloana* (Schult. & Schult.f.) Asch. & Graebn. in grasslands, collected by hand and G-Vac vacuum. It has been observed on sunny days in spring in the same foraging places as the ant *Camponotus termitarius*. In the laboratory, specimens were fed different species of *Drosophila*. Specimens of this species were recorded lifting the first pair of the legs and moving the pedipalps and the abdomen up and down when walking (https://www.youtube.com/watch?v=B_509xPe4zo). The courtship was recorded, during which the male lifts the first pair of the legs and moves sideways while approaching the female (<https://www.youtube.com/watch?v=MduAXI0r7wo>). Also, the mounting position of one couple was recorded, in which the receptive female lowered its body and the male inserted the left pedipalp, showing the ejaculation pulses as the spines of the legs rose (<https://www.youtube.com/watch?v=tM-QHw44ajE>).

Distribution

Known from Uruguay (Canelones, Lavalleja, Montevideo and Rivera) (Fig. 33).

Sarinda marcosi Piza, 1937

Figs 25–28, 33

Sarinda marcosi Piza, 1937: 309, pl. 1 figs 1–2 (female and male syntypes from São Paulo, Brazil, deposited in the Museu de Zoologia da Universidade de São Paulo 'Luiz de Queiroz', not examined). *Sarinda australis* Mello-Leitão, 1944: 387, fig. 85 (synonymized by Galiano 1965: 292).




Fig. 25. *Sarinda marcosi* Piza, 1937, photographs in vivo. **A.** Site where the species was collected. **B, D, F.** Male. **C, E, G.** Female.

Sarinda marcosi – Galiano 1965: 292, pl. II figs 1–7, pl. III fig. 4, pl. IV fig. 3, pl. V fig. 6, pl. VI fig. 3, pl. VIII figs 1, 4. — Jackowska & Prószyński 1975: 42, fig. 4h. — Hagopíán *et al.* 2021: 264, figs 1a–j, 3b–e, 4a–d, 5a–h. — Pett *et al.* 2021: 927, fig. 7a, c–d. — World Spider Catalog 2023.

Diagnosis and description

See Galiano (1965) and Hagopíán *et al.* (2021). Here, we include new color photographs of preserved and living specimens from Uruguay (Figs 25–28).

New records

URUGUAY – **Cerro Largo** • 4 ♂♂; Arévalo; 32.48422° S, 55.13208° W; 31 Oct. 2019; D. Hagopíán leg.; FCE-Ar 10511 • 1 ♀; Paso Arriera, Paraje Palleros; 32.00528° S, 54.49000° W; 5–6 Nov. 2020; M. Simó leg.; FCE-Ar 12532. – **Durazno** • 2 ♂♂; La Paloma; 32.72722° S, 55.57806° W; 25 Oct. 2021;



Fig. 26. *Sarinda marcosi* Piza, 1937, habitus A–C. Male (FCE-Ar 14321). D–F. Female (FCE-Ar 14321). A, D. Dorsal view. B, E. Lateral view. C, F. Ventral view.

D. Hagopíán and Á. Laborda leg.; FCE-Ar 13261 • 1 ♂; same locality as for preceding; 20 Oct. 2022; D. Hagopíán leg.; FCE-Ar 14323 • 1 ♂ 1 ♀; near La Paloma; 32.67472° S, 55.67272° W; 20 Oct. 2022; D. Hagopíán leg.; FCE-Ar 14321 • 2 ♂♂; La Paloma, Gruta “La Llorona”; 32.72944° S, 55.57583° W; 20 Oct. 2022; D. Hagopíán and A. Laborda leg.; FCE-Ar 13922. – **Lavalleja** • 1 ♂; Sierra de Minas, Parque de Vacaciones; 34.42600° S, 55.19500° W; 5–8 Dec. 2005; W. Maddison, G. Ruiz, M. Simó and M.E. Rodríguez leg.; WPM#05-046; UBCZ. – **Montevideo** • 1 ♂; Malvín, Instituto de Investigaciones Biológicas Clemente Estable (IIBCE); 34.88750° S, 56.14250° W; 13 Mar. 2020; D. Hagopíán leg.; FCE-Ar 12198 • 2 ♂♂; Parque Lecocq; 34.79278° S, 56.33167° W; 7 Dec. 2020; D. Hagopíán and A. Mailhos leg.; beating-sheet on a palm; FCE-Ar 12216. – **Río Negro** • 2 ♀♀; Ruta 24 Km 85 Estancia “Las Cadenas”; 32.52007° S, 58.03604° W; 8–14 Jan. 2021; D. Hagopíán and A. Mailhos leg.; beating-sheet in riparian forest; FCE-Ar 13790. – **Rocha** • 1 ♀; La Ribiera; 34.54500° S, 54.32306° W; 25 Mar. 2023; D. Hagopíán leg.; on *Eucalyptus* bark in the grass; FCE-Ar 14060.




Fig. 27. *Sarinda marcosi* Piza, 1937, face and chelicerae. **A, C.** Male (FCE-Ar 14321). **B, D.** Female (FCE-Ar 14321). **A–B.** Face. **C–D.** Chelicera.

Variation

Males (n = 10)

Total length (3.55–4.73), prosoma (1.63–2.47), abdomen (1.80–2.26), leg I (3.12–3.92), leg II (2.15–3.31), leg III (2.27–3.23), leg IV (3.88–5.42). Cheliceral teeth variation: four teeth in the promargin, one in retromargin (n = 5); four teeth in the promargin, one in the retromargin of right chelicera and two in the left (n = 1); five teeth in the promargin (apical tooth bifurcated), one in retromargin (n = 3); five teeth in the promargin of the right chelicera and four in the left, one in retromargin (n = 1).

Females (n = 5)

Total length (4.30–4.85), prosoma (2.00–2.31), abdomen (2.18–2.58), leg I (3.23–3.60), leg II (2.44–2.78), leg III (2.53–2.93), leg IV (3.78–4.65). Cheliceral teeth variation: four teeth in the promargin of the right chelicera and five in the left, one in retromargin (n = 4); four teeth in promargin, one in retromargin (n = 1).

Distribution

Known from Brazil (São Paulo), Argentina (Santa Fe, Chaco, Salta, Tucumán and Buenos Aires), Paraguay (Asunción). In Uruguay, it was found in Canelones, Cerro Largo, Durazno, Lavalleja, Maldonado, Montevideo, Río Negro, Rocha and Tacuarembó (Piza 1937; Galiano 1965; Hagopían *et al.* 2021; Pett *et al.* 2021; World Spider Catalog 2023) (Fig. 33).




Fig. 28. *Sarinda marcosi* Piza, 1937, photographs of genitalia. **A–D.** Male pedipalp (FCE-Ar 14321). **A.** Ventral view. **B.** Retrolateral view. **C.** Dorsal view. **D.** Prolateral view. **E–F.** Epigynum (FCE-Ar 14321). **E.** Ventral view. **F.** Dorsal view.




Fig. 29. *Parafuda banksi* Chickering, 1946, photographs in vivo. A. Site where the species was collected. B, D, F. Male. C, E, G. Female.

Genus *Parafluda* Chickering, 1946

Type species

Parafluda banksi Chickering, 1946.

Parafluda banksi Chickering, 1946
Figs 29–33

Parafluda banksi Chickering, 1946: 456 (male holotype from Panama, El Valle [Valle de Antón], [8°37'12" N, 80°07'48" W], deposited in the Museum of Comparative Zoology (MCZ), USA, not examined).



Fig. 30. *Parafluda banksi* Chickering, 1946, habitus. A–C. Male (FCE-Ar 11094). D–F. Female (FCE-Ar 9542). A, D. Dorsal view. B, E. Lateral view. C, F. Ventral view.

Sarinda albianus Mello-Leitão, 1947: 29 (male holotype from Carmo do Rio Claro, Minas Gerais, Brazil deposited in MNRJ 2185, destroyed by the fire, not examined. Synonymized by Galiano 1965: 292).

Parafluda banksi – Prószyński 2017: 61, fig. 27v. — Pett *et al.* 2021: 927, fig. 7h–i. — World Spider Catalog 2023.




Fig. 31. *Parafluda banksi* Chickering, 1946, face and chelicerae. A, C. Male (FCE-Ar 11094). B, D. Female (FCE-Ar 9542). A–B. Face. C–D. Chelicera.

Diagnosis and description

See Chickering (1946) and Galiano (1971). Here, we include color photographs of preserved and living specimens to complement the descriptions (Figs 29–32).

New records

URUGUAY – **Canelones** • 1 ♂; INIA, Las Brujas; 34.66236° S, 56.33886° W; 2 Feb. 2005; M. Simó leg.; FCE-Ar 3080 • 1 ♂; same locality as for preceding; 27 Dec. 2004; M. Simó leg.; FCE-Ar 3147 • 1 ♂; San Ramón; 34.29007° S, 55.95536° W; 21 Oct. 2018; R. Lauría leg.; in dry grass; FCE-Ar 9563 • 1 ♂; Las Toscas; 34.76639° S, 55.73972° W; 3 Nov. 2019; P. Martínez leg.; inside house; FCE-Ar 10513. – **Cerro Largo** • 1 ♀; Arévalo, Oficina UPM; 32.63611° S, 55.07194° W; 1 Nov. 2019; D. Hagopíán and Á. Laborda leg.; in fallen bark in grassland; FCE-Ar 12125. – **Durazno** • 1 ♀; Santa Bernardina; 33.35164° S, 56.51121° W; D. Hagopíán leg.; under cardboard in the grass; FCE-Ar 9509 • 1 ♂; La Paloma; 32.72745° S, 55.57797° W; 24 Oct. 2021; Á. Laborda leg.; under piece of wood in




Fig. 32. Photographs of *Parafuda banksi* Chickering, 1946, genitalia. **A–D** Male pedipalp (FCE-Ar 11094). **A.** Ventral view. **B.** Retrolateral view. **C.** Dorsal view. **D.** Prolateral view. **E–F.** Epigynum (FCE-Ar 9542). **E.** Ventral view. **F.** Dorsal view.

garden; FCE-Ar 11094. – **Lavalleja** • 1 ♂; Cerro de los Cuervos; 34.27125° S, 55.24791° W; 22 Dec. 2003; Simó and Pérez-Miles leg.; FCE-Ar 9713. – **Maldonado** • 1 ♂; Punta Negra; 34.88267° S, 55.22058° W; 5 Nov. 2018; R. Roibal leg.; FCE-Ar 9614. – **Montevideo** • 1 ♂; Barrio Malvín Norte, Facultad de Ciencias; 34.88239° S, 56.11824° W; 4 Oct. 2008, M. Simó leg; pitfall; FCE-Ar 4640 • 1 ♂; Cerro de Montevideo; 34.88869° S, 56.26094° W; 23 Nov. 1998; Sección Entomología leg.; FCE-Ar 7897 • 1 ♀; same locality as for preceding; 20 Dec. 1997; Sección Entomología leg.; FCE-Ar 5945 • 1 ♂; Melilla; 34.73261° S, 56.32110° W; 17 Nov. 2017; D. Hagopían leg.; FCE-Ar 9575; • 1 ♂; same locality as for preceding; 11 Aug. 2018; D. Hagopían leg.; under metal tank in grass; FCE-Ar 9377 • 1 ♀; same locality as for preceding; 30 Sep. 2018; D. Hagopían leg.; under metal tank in grass; FCE-Ar 9558. – **Paysandú** • 1 ♀; Rincón de Pérez; 32.17530° S, 57.51699° W; 18 Mar. 2004; Á. Laborda and M. Castro leg.; entomological net in grassland; FCE-Ar 4833. – **Rocha** • 1 ♀; 12 km from Castillos; 34.09417° S, 53.87166° W; 6 Apr. 2017; Á. Laborda leg.; in *Eryngium* sp.; FCE-Ar 9542.

Distribution

Known from Argentina (Buenos Aires), Brazil (Minas Gerais), Panamá (El Valle) and Paraguay (Ñeembucú) (Chickering 1946; Galiano 1965, 1971; Pett *et al.* 2021; World Spider Catalog 2023). Here, we report the first records of the species from Uruguay (Canelones, Cerro Largo, Durazno, Maldonado, Montevideo, Paysandú and Rocha) (Fig. 33).




Fig. 33. Distribution map with records in Uruguay of *Parafudra banksi* Chickering, 1946 (blue circles), *Sarinda marcosi* Piza, 1937 (yellow squares), *Sarinda contraluz* Hagopían & Bustamante sp. nov. (red inverted triangles) and *Sarinda sombraluminosa* Hagopían, Laborda & Simó sp. nov. (green triangles).

Discussion

The two new species are currently known only from Uruguay and both were observed in grasslands. *Sarinda sombraluminosa* sp. nov. was observed in livestock farms in natural grasslands (Pompozzi *et al.* 2022). *Sarinda contraluz* sp. nov. was mainly observed on *Cortaderia selloana*, known as “Pampa grass”. Although this grass was introduced in several countries in the world, it is native to South America (ranging from southern Brazil to Argentina). These results suggest that both species are associated with the Pampa biome so it is expected that their distributions also should extend to other countries such as southern Brazil and part of Argentina.

Several studies mention that representatives of *Sarinda* mimic ant species of the genus *Camponotus* (Galiano 1965, 1967; Maddison 2015; Hagopián *et al.* 2021). The new species here described share their habitat with the ant *Camponotus termitarius*, but behavioral studies are needed to determine the type of mimicry for these species. Also, the walking behavior is similar to that recorded for *S. marcosi* (Hagopián *et al.* 2021), in which the spider moves up and down the first pair of legs, the pedipalps and the abdomen while walking. Galiano (1971) mentions that *P. banksi* lifts the first pair of the legs while walking, as the antennae of ants, but the ant model and the mimicry type of this species remain unknown.

Sexual behavior was already described for the genus *Martella* (Galiano 1996), but here, we record the courtship and mating for the genus *Sarinda* for the first time. Galiano (1996) mentions that copulation took place on the walls of the recipient, outside the nest, as seen in *S. contraluz* sp. nov., but she did not describe any courtship behavior. Additional studies are needed to characterize the courtship and mating behaviors of the species of this genus.

Parafluda banksi shows a Neotropical distribution from Panama to Argentina (World Spider Catalog 2023). The species was found in synanthropic sites such as buildings or gardens but also in grasslands, as mentioned by Galiano (1971). *Sarinda marcosi* is known from southern South America (World Spider Catalog 2023) and, as *P. banksi*, the species inhabits a variety of habitats. The new records here reported are from native riparian forests, native ravine forests but also in synanthropic sites such as urban parks, gardens and *Eucalyptus* L'Hér. plantations.

This study presents different kinds of information, such as in-alcohol and live photography, SEM, natural history, distribution records, and behavioral videos. All this information complements the descriptions of the new taxa, and in turn, opens the way for future taxonomic and biogeographic studies, as well as mimetic and sexual behaviors.

Acknowledgments

We thank Ary Mailhos for his help in the field in collecting the specimens. To Stephany Arizala (Smithsonian Tropical Research Institute, Panama) for her helping on the location of the type locality of *Parafluda banksi*. To Ana Laura Reyes (Unidad de Microscopía Electrónica de Barrido, Facultad de Ciencias, UdelaR) for her assistance with the SEM photographs. This work was supported (in part) by a 2022 grant of the Hebert-Levi-Memorial-Fund-for-Arachnological-Research from the American Arachnological Society to AAB. MS acknowledges financial support by the Programa Desarrollo de Ciencias Básicas (PEDECIBA) and the Sistema Nacional de Investigadores (SNI), Uruguay. We also thank the editor and two anonymous reviewers for their helpful comments and suggestions, which improved the manuscript.

References

Bustamante A.A. & Ruiz G.R.S. 2017. Systematics of Thiodinini (Araneae: Salticidae: Salticinae), with description of a new genus and twelve new species. *Zootaxa* 4362 (3): 301–347.
<https://doi.org/10.11646/zootaxa.4362.3.1>

- Bustamante A.A. & Ruiz G.R.S. 2020. New species and records of thiodinines from North and South America (Araneae: Salticidae: Salticinae: Thiodinini). *Zootaxa* 4899 (1): 115–140. <https://doi.org/10.11646/zootaxa.4899.1.6>
- Cala-Riquelme F. 2021. Autodesk Sketchbook: an application that minimizes time and maximizes results of taxonomic drawing. *Zootaxa* 4963 (3): 577–586. <https://doi.org/10.11646/zootaxa.4963.3.10>
- Chickering A.M. 1946. The Salticidae (spiders) of Panama. *Bulletin of the Museum of Comparative Zoology at Harvard College* 97: 1–474. Available from <https://www.biodiversitylibrary.org/page/2808637> [accessed 5 Feb. 2024].
- Edwards G.B. 2004. Revision of the jumping spiders of the genus *Phidippus* (Araneae: Salticidae). *Occasional Papers of the Florida State Collection of Arthropods* 11: 1–156.
- Galiano M.E. 1964. Salticidae (Araneae) formiciformes. I. Revisión del género *Martella* Peckham, 1892. *Physis, Revista de la Sociedad Argentina de Ciencias Naturales (C)* 24: 353–363.
- Galiano M.E. 1965. Salticidae (Araneae) formiciformes IV. Revisión del género *Sarinda* Peckham, 1892. *Revista del Museo Argentino de Ciencias Naturales Bernardino Rivadavia (Entomología)* 1: 267–312.
- Galiano M.E. 1967. Salticidae (Araneae) formiciformes. VIII. Nuevas descripciones. *Physis, Revista de la Sociedad Argentina de Ciencias Naturales (C)* 27: 27–39.
- Galiano M.E. 1971. Salticidae (Araneae) formiciformes. XI. El género *Parafluda* Chickering, 1946. *Revista de la Sociedad Entomológica Argentina* 33: 63–68.
- Galiano M.E. 1996. Formiciform Salticidae (Araneae). Two new combinations and four new species of the genera *Martella* and *Sarinda*. *Miscel-lània Zoològica* 19: 105–115.
- Hagopíán D. & Laborda Á. 2020. Attack on Salticidae: new records of mantispids (Neuroptera: Mantispidae) in neotropical jumping spiders (Araneae: Salticidae). *Boletín de la Sociedad Zoológica del Uruguay* 29 (2): 167–170.
- Hagopíán D., Aisenberg A., Laborda Á. & Simó M. 2021. Morphological and behavioral traits associated with myrmecomorphy in *Sarinda marcosi* Piza, 1937 (Araneae: Salticidae: Sarindini). *Journal of Arachnology* 48 (3): 262–271. <https://doi.org/10.1636/JoA-S-19-069>
- Jackowska B. & Prószyński J. 1975. In search of the natural system of ant-like Salticidae. *Proceedings of the 6th International Arachnological Congress*: 39–43.
- Levi H.W. 1965. Techniques for the study of spider genitalia. *Psyche* 72: 152–158. <https://doi.org/10.1155/1965/94978>
- Maddison W.P. 2015. A phylogenetic classification of jumping spiders (Araneae: Salticidae). *Journal of Arachnology* 43 (3): 231–292. <https://doi.org/10.1636/arac-43-03-231-292>
- Mello-Leitão C.F. de 1944. Arañas de la provincia de Buenos Aires. *Revista del Museo de La Plata (N.S., Sección Zoología)* 3: 311–393.
- Mello-Leitão C.F. de 1947. Aranhas de Carmo do Rio Claro (Minas Gerais) coligidas pelo naturalista José C.M. Carvalho. *Boletim do Museu Nacional do Rio de Janeiro (N.S., Zoologia)* 80: 1–34.
- Müller H.G. & Cutler B. 1989. The genus *Sarinda* Peckham 1892 in N-Colombia (Arachnida: Araneae: Salticidae). *Senckenbergiana Biologica* 69: 73–76.
- Petrunkevitch A. 1925. Arachnida from Panama. *Transactions of the Connecticut Academy of Arts and Sciences* 27: 51–248.

Pett B.L., Rubio G.D. & Stolar C.E. 2021. A first baseline for the salticid (Araneae: Salticidae) fauna of Paraguay, with thirty-two new records and description of a new species from Paraguay and Argentina. *Arachnology* 18 (8): 922–935. <https://doi.org/10.13156/ arac.2021.18.8.922>

Piza Jr S. de T. 1937. Novas especies de aranhas myrmecomorphas do Brasil e considerações sobre o seu mimetismo. *Revista do Museu Paulista* 23: 307–319.

Pompozzi G., de Santiago F., Blumetto O. & Simó M. 2022. Livestock systems preserving natural grasslands are biodiversity reservoirs that promote spiders' conservation. *Journal of Insect Conservation* 26: 453–462. <https://doi.org/10.1007/s10841-022-00399-y>

Prószyński J. 2017. Pragmatic classification of the world's Salticidae (Araneae). *Ecologica Montenegrina* 12: 1–133. <https://doi.org/10.37828/em.2017.12.1>

Shorthouse David P. 2010. *SimpleMappr, an Online Tool to Produce Publication-quality Point Maps*. Available from <https://www.simplemappr.net> [accessed 21 Apr. 2023].

World Spider Catalog 2023. *World Spider Catalog, Version 24*. Natural History Museum Bern. Available from <http://wsc.nmbe.ch> [accessed 21 Apr. 2022]. <https://doi.org/10.24436/2>

Manuscript received: 30 May 2023

Manuscript accepted: 31 October 2023

Published on: 7 March 2024

Topic editor: Tony Robillard

Section editor: Rudy Jocqué

Desk editor: Pepe Fernández

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic.