

This work is licensed under a Creative Commons Attribution License (CC BY 4.0).

Research article

<urn:lsid:zoobank.org:pub:B935D2BA-84E7-44C5-A985-61CCD87134C7>

Taxonomic reassessment of the *Herpetoreas xenura* (Wall, 1907) (Squamata: Serpentes: Natricidae) from Myanmar with description of a new species

Tan Van NGUYEN^{1,*}, Hmar Tlawmte LALREMSANGA²,
Lal BIAKZUALA³ & Gernot VOGEL^{4,*}

¹Institute for Research and Training in Medicine, Biology and Pharmacy, Duy Tan University,
Da Nang, 550000, Vietnam.

¹College of Medicine and Pharmacy, Duy Tan University, 120 Hoang Minh Thao,
Lien Chieu, Da Nang, 550000, Vietnam.

^{2,3}Developmental Biology and Herpetology Laboratory, Department of Zoology, Mizoram University,
796004 Tuanhil, Mizoram, India.

⁴Society for South East Asian Herpetology, Im Sand-3, Heidelberg, Germany.

*Corresponding authors: tan.sifasv@gmail.com; Gernot.Vogel@t-online.de

²Email: htlrsa@yahoo.co.in

³Email: bzachawngthu123@gmail.com

¹<urn:lsid:zoobank.org:author:94EABB6E-180C-424E-8E9C-60DB36125550>

²<urn:lsid:zoobank.org:author:F7D2FBD8-87EB-4962-BC8E-87AD754F7E8C>

³<urn:lsid:zoobank.org:author:D4198FBC-CB58-453B-B0E1-568F47ECE876>

⁴<urn:lsid:zoobank.org:author:26C84E11-E43F-4076-B268-28C32E9FD2D0>

Abstract. The review of the taxonomic status of *Herpetoreas xenura* species complex, based on morphological and molecular data, revealed a new species from Myanmar, which we describe as *Herpetoreas davidi* sp. nov. from the Rakhine Yoma Elephant Wildlife Sanctuary in Rakhine State. It is suggested to be a sister species to *Herpetoreas peali* and *Herpetoreas xenura* sensu stricto, and can be separated from the latter species by a combination of morphological and scalation characters, and by its pattern. Another population from the Htamanthi Wildlife Sanctuary, Sagaing Region, was previously confused with *H. xenura*. However, as a closer examination of its morphology, shows it to be closer to *Hebius khasiensis*, we herein tentatively refer to this population as *Hebius khasiensis*, pending further molecular data confirming its taxonomic status. Consequently, we propose to temporarily remove *Herpetoreas xenura* from the fauna of Myanmar. Further studies, especially in the northwestern region of Myanmar, such as near border with India in Chin Hills and Naga Hills, are required. An updated key for the species of *Herpetoreas* is also provided.

Keywords. Distribution, *Herpetoreas davidi* sp. nov., Indo-Burmese region, Naga Hills, Rakhine Hills, taxonomy.

Nguyen T.V., Lalremsanga H.T., Biakzuala L. & Vogel G. 2024. Taxonomic reassessment of the *Herpetoreas xenura* (Wall, 1907) (Squamata: Serpentes: Natricidae) from Myanmar with description of a new species. *European Journal of Taxonomy* 931: 158–203. <https://doi.org/10.5852/ejt.2024.932.2519>

Introduction

The Himalaya mountain keelbacks of the genus *Herpetoreas* Günther, 1860 are widely distributed across the southern foothills of the Himalaya Mountains as well as Eastern Himalayas, including northeastern Pakistan, northern and northeastern India, southwestern China (Region of Tibet), Nepal, Bhutan, Bangladesh, and Myanmar (Lalremsanga *et al.* 2022; Ren *et al.* 2022). Several recent studies have demonstrated that the diversity of *Herpetoreas* is still underestimated and the species received much controversy and confusion with species of the genera *Amphiesma* Duméril, Bibron & Duméril, 1854 and, especially, *Hebius* Thompson, 1913 (Smith 1943; Malnate 1966; Guo *et al.* 2014; Peng *et al.* 2021; Deepak *et al.* 2022; Lalremsanga *et al.* 2022; Ren *et al.* 2022). The genus *Herpetoreas* is morphologically characterised by: head moderately distinct from neck; body cylindrical, maximum TL 943 mm; tail relatively long, TaL/TL 0.227–0.317; nostrils and eye directed laterally; supralabials 8–9, usually 3rd–5th or 4th and 5th entering orbit; dorsal scale rows 19–19–17; scales weakly to distinctly keeled, notched at their apical part; ventrals 136–234; cloacal plate and subcaudals divided or not; maxillary teeth 13–23, slightly increasing in size, last two to three teeth distinctly enlarged, separated from anterior teeth by a small diastema; hemipenis short and thin, shallowly bilobed, spinous throughout with single basal hook; sulcus spermaticus single, centripetal, extends to the inner right lobe or to the crotch only; apical naked area on the crotch weakly developed, not protruding, not visible from asulcate surface; venter yellowish-beige, each ventral scale decorated with dark spots at lateral edge or not (Günther 1860; Guo *et al.* 2014; Ren *et al.* 2022). Currently, the genus *Herpetoreas* comprises seven species, namely: *Herpetoreas burbrinki* Guo, Zhu, Liu, Zhang, Li, Huang & Pyron, 2014; *H. murlen* Lalremsanga, Bal, Vogel & Biakzuala, 2022; *H. pealii* (Sclater, 1891); *H. platyceps* (Blyth, 1854); *H. sieboldii* Günther, 1860; *H. xenura* (Wall, 1907); and *H. tpsen* Ren, Jiang, Huang, David & Li, 2022 (Lalremsanga *et al.* 2022; Ren *et al.* 2022).

The Strange-tailed keelback or Wall's Keelback, *Herpetoreas xenura*, was described by Wall (1907) based on a specimen in bad condition from an unknown type locality (see Fig. 1A), and, furthermore, the holotype of this species has been lost (Smith 1943). Subsequently, Wall (1909) reported about four additional specimens of this species from Cherrapunji in the Khasi Hills, Assam (now Meghalaya State) which were discovered in the Indian Museum (now the Zoological Survey of India [ZSI], Kolkata [Calcutta], West Bengal, India). These specimens were labelled *Hebius modestus* (Günther, 1875), to which the species bears resemblance in general coloration but is differentiated by its undivided subcaudals, number of preoculars, number of temporals, and number of keeled scales on body (Wall 1907). Therefore, "Khasi Hill, Meghalaya State, India" is assumed to be the type locality. The semi-aquatic species *Herpetoreas xenura* has been reported to be distributed from Northeast India to Bangladesh and Myanmar (Wall 1907; Wogan *et al.* 2008; Ren *et al.* 2022; Uetz *et al.* 2023). But the records of this species from Rakhine State and Sagaing Region of Myanmar, however, were not accompanied by information on morphology or molecular data. During examination of the herpetological collections of the California Academy of Sciences (CAS, USA), one of us (G. Vogel) encountered six specimens of *Herpetoreas* from Myanmar (five from Rakhine State and one from Sagaing Region), which were originally identified as *Herpetoreas xenura* by Wogan *et al.* (2008). However, a more detailed morphological re-examination of these six specimens revealed morphological differences between these two populations. In the present study, we provide a detailed morphological description and comparison of the populations of *Herpetoreas* in Myanmar and demonstrate that they belong to two taxa that are morphologically distinct from true *H. xenura* and other congeners. Molecular analyses based on mitochondrial cytochrome b (Cytb) confirm the distinctiveness of the Rakhine population,




Fig. 1. *Herpetoreas xenura* (Wall, 1907), in preservative. **A.** Holotype (lost), sex unknown. **B.** ♂, general dorsal, ventral, and head view (MZMU 1211). **C.** ♀, general dorsal and ventral view (MZMU 3271). Drawing (A) by J. Green Delet Lith in Wall (1907). Photos by G. Vogel (A–C).

which is formally described below as a new species. Meanwhile the population at Sagaing should be assigned to *Hebius khasiensis* (Boulenger, 1890).

Material and methods

Material examined

For this study, a total of 19 preserved specimens of the *Herpetoreas xenura* group were examined for their external morphological characters (five specimens from Rakhine State and one specimen from Sagaing Region, Myanmar; and 13 specimens of *Herpetoreas xenura* from India, see Table 1). The other 58 examined specimens of Natricidae Bonaparte, 1838, including *Hebius khasiensis* (28 specimens), *Herpetoreas sieboldii* (19 specimens), and *Herpetoreas platyceps* (six specimens), are listed in Appendix 1. Measurements were taken with a slide calliper to the nearest 0.1 mm, except for body and tail lengths, which were measured to the nearest mm with a measuring tape.

The following morphometric and meristic characters were recorded:

ATEM	=	anterior temporals
CS	=	condition of the cloacal plate
DSR	=	dorsal scale rows
E-Ns	=	eye to nostril distance
E-Sn	=	eye to snout distance
ED	=	eye diameter
FL	=	length of frontal
HD	=	head depth
HL	=	head length
HW	=	head width
IL	=	infralabials
INL	=	length of internasal scales
Lor	=	loreal
PFL	=	length of prefrontals
PL	=	length of parietals
PoO	=	postoculars
PrO	=	preoculars
PTEM	=	posterior temporals
PVEN	=	pre-ventrals
SC	=	subcaudals
SL	=	supralabials
SL-E	=	supralabials contacting the eye
SVL	=	snout-vent length
SW	=	snout width
TaL	=	tail length
TL	=	total length
VEN	=	ventrals

The number of ventral scales was counted according to Dowling (1951). Dorsal scales were counted at one head length posterior to the neck, at the midpoint of snout-vent length, and at one head length anterior to the cloacal plate. Subcaudals were counted from immediately posterior to the anal scute to the tail tip, but excluding the terminal scute. Values for paired head characters were recorded on both sides of the head and were reported in a left/right order. The sex was determined by dissection of the ventral tail base.

Table 1 (continued on next page). Main measurements and meristic characters of the *Herpetoreas davidi* sp. nov., *H. xenura* (Wall, 1907), and *Hebius khasiensis* (Boulenger, 1890). Abbreviations: F = adult female; *Heb.* = *Hebius*; *Her.* = *Herpetoreas*; M= adult male; N/a = not available; SF = subadult female; SM = subadult male.

No.	Previously taxa	Propose taxa	Locality	Number	Sex	Status	SVL (mm)	TaL (mm)
1	<i>Hep. xenura</i>	<i>Hep. davidi</i> sp. nov.	Gawa, Rakhine, Myanmar	CAS 222969	M	Holotype	384	168
2	<i>Hep. xenura</i>	<i>Hep. davidi</i> sp. nov.	Gawa, Rakhine, Myanmar	CAS 220378	SM	Paratype	292	126
3	<i>Hep. xenura</i>	<i>Hep. davidi</i> sp. nov.	Gawa, Rakhine, Myanmar	CAS 220550	F	Paratype	397	178
4	<i>Hep. xenura</i>	<i>Hep. davidi</i> sp. nov.	Gawa, Rakhine, Myanmar	CAS 220256	F	Paratype	482	N/a
5	<i>Hep. xenura</i>	<i>Hep. davidi</i> sp. nov.	Gawa, Rakhine, Myanmar	CAS 220332	SF	Paratype	259	115
6	<i>Hep. xenura</i>	<i>Heb. khasiensis</i>	Khandi, Sagaim, Myanmar	CAS 232203	F		153	72
7	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 810	M		650	N/a
8	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 811	M		635	195
9	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 812	M		675	195
10	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 813	M		620	175
11	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 814	M		670	200
12	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 816	M		635	185
13	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 817	M		560	170
14	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 818	M		480	143
15	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 1211	M		501	198
16	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 819	F		565	180
17	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 820	F		424	124
18	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 3271	F		417	161
19	<i>Hep. xenura</i>	<i>Hep. xenura</i>	Mizoram, India	MZMU 815	SF		110	N/a

NGUYEN T.V. *et al.*, Taxonomic reassessment of the *Herpetoreas davidi* sp. nov., *H. xenura* (Wall, 1907), and *Hebius khasiensis* (Boulenger, 1890). Abbreviations: F = adult female; *Heb.* = *Hebius*; *Her.* = *Herpetoreas*; M = adult male; N/a = not available; SF = subadult female; SM = subadult male.

No.	Propose taxa	Number	Sex	VEN	SC	SC	CP	DSR	SL	SL-E	IL
1	<i>Hep. davidi</i> sp. nov.	CAS 222969	M	154	99	single	single	19-19-17	9/9	4-6	9/10
2	<i>Hep. davidi</i> sp. nov.	CAS 220378	SM	156	97	single	single	19-19-17	9/9	4-6	10/10
3	<i>Hep. davidi</i> sp. nov.	CAS 220550	F	151	97	single	single	19-19-17	9/9	4-6	10/10
4	<i>Hep. davidi</i> sp. nov.	CAS 220256	F	155	N/a	single	single	19-19-17	9/9	4-6	10/10
5	<i>Hep. davidi</i> sp. nov.	CAS 220332	SF	152	100	single	single	19-19-17	9/9	4-6	10/9
6	<i>Heb. khasiensis</i>	CAS 232203	F	134	92	divided	divided	19-19-17	9/9	4-6	9/10
7	<i>Hep. xenura</i>	MZMU 810	M	160	N/a	single	single	19-19-17	9/9	4-6	9/9
8	<i>Hep. xenura</i>	MZMU 811	M	159	102	single	single	19-19-17	9/9	4-6	9/9
9	<i>Hep. xenura</i>	MZMU 812	M	158	86	single	single	19-19-17	7/8	4-6	8/8
10	<i>Hep. xenura</i>	MZMU 813	M	158	95	single	single	19-19-17	7/8	4-6	8/8
11	<i>Hep. xenura</i>	MZMU 814	M	164	103	single	single	19-19-17	7/7	4-5/4-6	7/7
12	<i>Hep. xenura</i>	MZMU 816	M	161	86	single	single	19-19-17	9/9	4-6	9/9
13	<i>Hep. xenura</i>	MZMU 817	M	160	92	single	single	19-19-17	8/8	4-6	8/8
14	<i>Hep. xenura</i>	MZMU 818	M	160	101	single	single	19-19-17	8/8	4-6	8/8
15	<i>Hep. xenura</i>	MZMU 1211	M	165	105	single	single	19-19-17	9/9	4-6	9/10
16	<i>Hep. xenura</i>	MZMU 819	F	165	110	single	single	19-19-17	9/9	4-6	9/9
17	<i>Hep. xenura</i>	MZMU 820	F	158	97	single	single	19-19-17	8/8	4-6	8/8
18	<i>Hep. xenura</i>	MZMU 3271	F	155	87	single	single	19-19-17	9/9	4-6	10/10
19	<i>Hep. xenura</i>	MZMU 815	SF	164	87	single	single	19-19-17	8/8	4-6	7/7

For comparison with other taxa, we relied on previously published data (e.g., Wall 1907, 1909; Smith 1943; Malnate 1966; Mathew & Meetei 2004; Reza 2010; Guo *et al.* 2014; Das *et al.* 2020; Malsawmdawngliana *et al.* 2021; Peng *et al.* 2021; Lalremsanga *et al.* 2022; Ren *et al.* 2022); and our examined specimens. The distribution map was prepared in QGIS ver. 3.16.2.

Museum abbreviations

AUP	= School of Agriculture and Natural Resources, University of Phayao, Phayao, Thailand
BNHS	= Bombay Natural History Society, Mumbai, India
CAS	= California Academy of Science, San Francisco, USA
CHS	= Private catalogue of Huang Song (China)
CIB	= Chengdu Institute of Biology, Chinese Academy of Sciences, Chengdu, Sichuan, China
FMNH	= The Field Museum, Chicago, USA
GP	= Private catalogue of Guo Peng (China)
JU	= Jahangirnagar University, Dhaka, Bangladesh
KIZ	= Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, China
KUZ	= Kyoto University, Department of Zoology Museum, Japan
MNHN	= Muséum national d'histoire naturelle, Paris, France
MZMU	= Departmental Museum of Zoology, Mizoram University, India
NHMUK	= The Natural History Museum (formerly BMNH), London, United Kingdom
NHMW	= Naturhistorisches Museum (formerly NMW), Wien, Austria
PUCZM	= Pachhunga University College, Mizoram University, Mizoram, India
QSMI	= Queen Saovabha Memorial Institute, Bangkok, Thailand
RMNH	= Naturalis-Nationaal Natuurhistorisch Museum (Naturalis), Leiden, Netherlands
SYNU	= Shenyang Normal University, Shenyang, China
USNM	= National Museum of Natural History, Smithsonian Institution, Washington, USA
VNMN	= Vietnam National Museum of Nature, Hanoi, Vietnam
VNUF	= Vietnam National University of Forestry, Hanoi, Vietnam
WII	= Wildlife Institute of India, Uttarakhand, India
YBU	= Yibin University, Sichuan, China
ZMMU	= Zoological Museum of Moscow, Moscow, Russia
ZSI	= Zoological Survey of India (formerly ZSIC), Kolkata, West Bengal, India

Molecular analyses

We extracted the Genomic DNA from the liver tissues using QIAamp DNA Mini Kit following the manufacturer's procedure. The fragment of Cytb was amplified using the forward (L14910) and reverse (H16064) primers (Burbrink *et al.* 2000) at the thermal conditions of 94°C for 3 min, followed by 35 cycles of 94°C for 30 sec, 49°C for 40 sec, 72°C for 30 sec, and with a final extension at 72°C for 5 min. The amplified products were purified and sequenced using Sanger's dideoxy method at Barcode BioSciences, Bangalore, India.

The datasets of natricid snakes comprise the sequence of Cytb gene of *Herpetoreas* sp. (CAS 220256) from Rakhine State, Myanmar, and *Hebius venningi* (Wall, 1910) (MZMU 1313 [isolate A, B] with sequence accession numbers: OR389974-75, respectively) from Mizoram State, India along with 52 congeneric sequences and outgroups obtained from the NCBI database (see Appendix 2). The dataset was aligned with the MUSCLE algorithm (Edgar 2004) using default parameters in MEGA 11 (Tamura *et al.* 2021). The dataset was partitioned by codon positions. PartitionFinder ver. 2 (Lanfear *et al.* 2016) was employed for selecting the best partitioning schemes and for evolutionary model searching under the Bayesian Information Criterion (Appendix 2). Bayesian Inference (BI) and Maximum Likelihood (ML) phylogenies were performed in Mr.Bayes ver. 2.3.5 (Ronquist *et al.* 2012) and in IQTREE (Nguyen *et al.* 2015), respectively. In the BI analyses, four independent runs with one cold and three hot

chains were conducted for 20 million generations and sampled every 5000 generations. The trace plots generated by the MCMC runs were viewed in Tracer ver. 1.7 (Rambaut *et al.* 2018) and the first 25% of samples were discarded as burnin. The ML analysis was performed using the partitioning schemes selected by PartitionFinder ver. 2 (Lanfear *et al.* 2016), and the models selected under BIC scores by ModelFinder (Kalyaanamoorthy *et al.* 2017) implemented in the IQTREE (Nguyen *et al.* 2015) at 10 000 Ultrafast Bootstrap (UFB) (Minh *et al.* 2013). The uncorrected p-distances were estimated in MEGA 11 (Tamura *et al.* 2021). The standardized p-distances were further utilized for multivariate Principal Coordinate Analysis (PCoA) (Gower 1966) to visualize the genetic differentiation among the study taxa in PAST ver. 4.13 (Hammer *et al.* 2001). We applied Assemble Species by Automatic Partitioning (ASAP) (Puillandre *et al.* 2021), a single locus-based species delimitation method. This method produced species partitions based on the probabilities and barcode gap width; these two metrics are combined into a single ASAP-score for ranking the partitions, the lower the ASAP-score, the better the partition (Puillandre *et al.* 2021).

Results

Phylogenetic relationships

The Cytb dataset of the natricid species contained 1124 bp aligned characters. Two types of partitioned phylogenetic analyses (BI and ML) were performed, which largely inferred consistent tree topologies (Fig. 2A; [Supp. file 1](#)). In our phylogenetic analyses of the natricid group, the specimen of *Herpetoreas* cf. *xenura* from Rakhine, Myanmar, is nested with the putative *Herpetoreas* clade and forms a sister lineage to the sub-clade consisting of *H. pealii* and *H. xenura* sensu stricto with well-supported Bayesian posterior probabilities (PP) and UFB values (PP=0.96; UFB=98). It is also distinct from the other species of *Herpetoreas* by considerable genetic divergences of 19.2% (to *H. murlen*) to 30.8% (to *H. tpser*), and 26.9–30.8% with *H. xenura* sensu stricto; while 26.9% genetic distances are recovered with the two species of *Amphiesma*, and 38.5% genetic divergence from *Sahyadriophis beddomei* (Günther, 1864); the genetic divergence with species of *Hebius* ranged from 15.4–38.5% (to *Hebius andreae* (Ziegler & Le, 2006) and *H. pryeri* (Boulenger, 1887), respectively) and 46.2% to the outgroup (*Amphiesmoides ornaticeps* (Werner, 1924)) (Appendix 3). The ordination of the genetic divergence along the first two Principal Coordinate (PCo) axes also reveal the discrete clustering of the genera: *Amphiesma*, *Hebius*, *Herpetoreas* and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023 (Fig. 2B). The Rakhine specimen of *H. cf. xenura* is also clearly discriminated from *H. xenura* sensu stricto as well as from the other species of *Herpetoreas* (Fig. 2C). The molecular species delimitation produced 10 partitions, out of which 90% of the partitions including the optimal partition selected (ASAP-score = 2.0) inferred *H. cf. xenura* (Rakhine) as distinct species (Fig. 3).

Taxonomy

The morphological features in five specimens of *Herpetoreas* cf. *xenura* from Rakhine Yoma Elephant Sanctuary, Rakhine State, agree with the generic characters of *Herpetoreas* provided by Guo *et al.* (2014) and Ren *et al.* (2022), but exhibit a distinct lineage with strong genetic divergence from the seven nominal species of *Herpetoreas*. Overall, both molecular and morphological differentiation leave no doubt that the species of *Herpetoreas* from Rakhine State represents a distinct undescribed species. Another juvenile specimen (CAS 232203) originally classified as *Herpetoreas* cf. *xenura* from Htamanthi Wildlife Sanctuary, Sagaing Region, Myanmar (Wogan *et al.* 2008), was examined. The morphological characteristics (pholidosis and colour pattern) of CAS 232203 are not consistent with the generic characters of *Herpetoreas*. The important morphological characters and photos of the specimen CAS 232203 are presented in Table 1 and Fig. 4A–F. However, the specimen CAS 232203 is very similar to the diagnostic characters of *Hebius khasiensis* see Table 2 and Figs 4G–H, 6G–H. Due to the lack of genetic information for the Sagaing specimen, we herein tentatively refer to this specimen as *Hebius*





Fig. 2. **A.** Bayesian Inference (BI) phylogenetic tree based on the fragment of mitochondrial cytochrome b gene among the genera: *Amphiesma* Duméril, Bibron & Duméril, 1854, *Amphiesmoides* Malnate, 1961, *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023. The posterior probability (PP) support values are given at each branch. **B.** Ordination of standardized p-distance of the genera: *Amphiesma*, *Amphiesmoides*, *Hebius*, *Herpetoreas*, and *Sahyadriophis* (excluding outgroup) along the first and second principal coordinate (PCo) axes where a total of 56% and 13% of variance are captured by PCo1 and PCo2, respectively. **C.** Ordination of standardized p-distance of *Herpetoreas* along the first and second PCo axes where a total of 52% and 16% of variance are captured by PCo1 and PCo2, respectively.






Fig. 4. *Hebius khasiensis* (Boulenger, 1890). A–F. Specimen juvenile, sex unknown, in preservative (CAS 232203). A. Lateral view of the head, right side. B. Lateral view of the head, left side. C. Dorsal view of the head. D. Ventral view of the head. E. General dorsal view. F. General ventral view. G. *Tropidonotus khasiensis*, syntype, ♂ (NHMUK 1946.1.13.62), lateral view of the head, right side. H. *Natrix gilhodesi* Wall, 1925, holotype, ♂ (NHMUK 1946.1.12.81), lateral view of the head, right side. Photos by G. Vogel.

Table 2 (continued on next two pages). Comparison of morphological characteristics of *Herpetoreas davidi* sp. nov. with those of the genus *Herpetoreas*

Günther, 1860. Abbreviations: *Heb.* = *Hebius*; *Her.* = *Herpetoreas*; N/a = not available.

No.	Species	SVL (mm)				Tal (mm)				VEN	
		Males	Females	Males	Females	Males	Females	Males	Females	Males	Females
1	<i>Hep. davidi</i> sp. nov.	292–384	259–482	126–168	115–178	0.301–0.304	0.207–0.310	154–156	151–155		
2	<i>Hep. xenura</i>	[\bar{x} = 338.0, n = 2]	[\bar{x} = 379.3, n = 3]	[\bar{x} = 147.0, n = 2]	[\bar{x} = 146.5, n = 2]	[\bar{x} = 0.303, n = 2]	[\bar{x} = 0.309, n = 2]	[\bar{x} = 155, n = 2]	[\bar{x} = 152.7, n = 3]		
3	<i>Hep. burbrinki</i>	[\bar{x} = 590.6, n = 10]	[\bar{x} = 468.7, n = 3]	[\bar{x} = 182.9, n = 10]	[\bar{x} = 155.0, n = 3]	[\bar{x} = 0.240, n = 9]	[\bar{x} = 0.249, n = 3]	[\bar{x} = 160.7, n = 9]	[\bar{x} = 160.5, n = 4]		
4	<i>Hep. murien</i>	527	462	230	184	0.304	0.304	172	169		
5	<i>Hep. pealii</i>	[n = 1]	[n = 1]								
6	<i>Hep. platiceps</i>	355	N/a	106	N/a	0.230	N/a	179	N/a		
7	<i>Hep. sieboldii</i>	334	511	117	150	0.259	0.227	142–144	136		
8	<i>Hep. ipseser</i>	[n = 1]	[n = 2]	[n = 1]	[n = 1]						
9	<i>Heb. khasiensis</i>	328–584	318–552	115–203	102–224	0.243–0.267	0.232–0.273	205–234	191–216		
10	<i>Heb. khasiensis</i>	[\bar{x} = 476, n = 13]	[\bar{x} = 485.7, n = 6]	[\bar{x} = 170.6, n = 9]	[\bar{x} = 174.0, n = 3]	[\bar{x} = 0.252, n = 5]	[\bar{x} = 0.248, n = 10]	[\bar{x} = 220.8, n = 8]	[\bar{x} = 202.1, n = 13]		
11	<i>Hep. cf. sieboldii</i>	355–479	279–491	165	108–207	0.246–0.294	0.243–0.296	193–207	174–185		
		[\bar{x} = 417.0, n = 2]	[\bar{x} = 410.8, n = 4]	[n = 1]	[\bar{x} = 169.8, n = 4]	[n = 1]	[\bar{x} = 0.293, n = 4]	[\bar{x} = 161.5, n = 2]	[\bar{x} = 159.5, n = 4]		
		271–412	113–43.5	138–183	53–192	0.320–0.346	0.296–0.329	142–154	141–152		
		[\bar{x} = 333.4, n = 13]	[\bar{x} = 281.4, n = 20]	[\bar{x} = 164.7, n = 10]	[\bar{x} = 122.5, n = 15]	[\bar{x} = 0.338, n = 10]	[\bar{x} = 0.312, n = 15]	[\bar{x} = 148.2, n = 13]	[\bar{x} = 147.7, n = 20]		
		N/a	153	N/a	72	N/a	0.320	N/a	134		
		[n = 1]	[n = 1]								
		407	N/a	153	N/a	0.273	N/a	200		[n = 1]	
		[n = 1]									

Table 2 (continued). Comparison of morphological characteristics of *Herpetoreas davidi* sp. nov. with those of the genus *Herpetoreas* Günther, 1860.
 Abbreviations: *Heb.* = *Hebius*; *Her.* = *Herpetoreas*; N/a = not available.

No.	Species	SC			VEN+CS			DSR			Lor		AS		SC
		Males	Females	Males	Females	Males	Females	19–19–17	1	Single	Single	Single	Single		
1	<i>Hep. davidi</i> sp. nov.	97–99	97–100	253	248–252	19–19–17	1	Single	Single	Single	Single	Single	Single	Single	Single
2	<i>Hep. xenura</i>	[\bar{x} = 98.0, n = 2] 86–105	[\bar{x} = 98.5, n = 2] 87–110	[n = 2] 244–270	[\bar{x} = 250.0, n = 2] 242–275	19–19(17)–17	1	Single	Single	Single	Single	Single	Single	Single	Single
3	<i>Hep. burbrinki</i>	[\bar{x} = 96.6, n = 9] 96	[\bar{x} = 95.3, n = 4] 94	[\bar{x} = 257.3, n = 9] 268	[\bar{x} = 255.8, n = 4] 263	19–19–17	1 or 2	Divided	Divided	Divided	Divided	Divided	Divided	Divided	Divided
4	<i>Hep. murlen</i>	[n = 1] 78	[n = 1] N/a	[n = 1] 257	[n = 1] N/a	19–19–17	1	Divided	Divided	Divided	Divided	Divided	Divided	Divided	Divided
5	<i>Hep. pealii</i>	[n = 1] 75–77	[n = 1] 69	[n = 1] 219	[n = 1] 205	19–19–17	1	Single	Single	Single	Single	Single	Single	Single	Single
6	<i>Hep. platiceps</i>	[\bar{x} = 92.2, n = 6] 88–98	[\bar{x} = 86.1, n = 11] 78–96	[\bar{x} = 316.7, n = 6] 308–326	[\bar{x} = 287.3, n = 11] 271–302	19–19–17	1	Divided	Divided	Divided	Divided	Divided	Divided	Divided	Divided
7	<i>Hep. sieboldii</i>	[\bar{x} = 102.3, n = 9] 91–109	[\bar{x} = 100.7, n = 3] 97–106	[\bar{x} = 303.0, n = 9] 296–314	[\bar{x} = 283.3, n = 3] 279–287	19–19–17	1	Divided	Divided	Divided	Divided	Divided	Divided	Divided	Divided
8	<i>Hep. tpsper</i>	[n = 1] 96	[n = 1] 81–97	[n = 1] 255	[n = 1] 239–253	19–19–17	1	Divided	Divided	Divided	Divided	Divided	Divided	Divided	Divided
9	<i>Heb. khasiensis</i>	[n = 1] 96–111	[\bar{x} = 88.3, n = 4] 87–99	[n = 1] 238–264	[\bar{x} = 247.8, n = 4] 230–251	19–19–17	1	Divided	Divided	Divided	Divided	Divided	Divided	Divided	Divided
10	<i>Heb. khasiensis</i>	[\bar{x} = 103.3, n = 10] N/a	[\bar{x} = 93.8, n = 15] 92	[\bar{x} = 250.7, n = 10] N/a	[\bar{x} = 241.9, n = 15] 226	19–19–17	1	Divided	Divided	Divided	Divided	Divided	Divided	Divided	Divided
11	<i>Hep. cf. sieboldii</i>	N/a	[n = 1] 107	N/a	[n = 1] 310	19–19–17	1	Divided	Divided	Divided	Divided	Divided	Divided	Divided	Divided

Table 2 (continued). Comparison of morphological characteristics of *Herpetoreas davidi* sp. nov. with those of the genus *Herpetoreas* Günther, 1860.
 Abbreviations: *Heb.* = *Hebius*; *Her.* = *Herpetoreas*; N/a = not available.

No.	Species	SL	SL-E	IL	PrO	PoO	ATEM	PTEM	MT	VEN colours	VEN lateral blotch	Distributions	Sources
1	<i>Hep. davidi</i> sp. nov.	9	4–6	9 or 10	1	3	1–2	2	N/a	whitish or yellowish	barely present, discontinuous	Myanmar	our data
2	<i>Hep. xenura</i>	9 (8 or 7)	4–6 (4–5)	9–9	1	3	1–2	1–2	22–23	whitish or yellowish	present, discontinuous	India, Bangladesh	Wall (1907, 1909); Mathew & Meetei (2004); Reza (2010); Malsawndawnglana et al. (2021); our data
3	<i>Hep. burbrinki</i>	8 (9)	3–5	10	1–2	3	2–3	2	21	creamish-beige	present, discontinuous	China	Guo et al. (2014); Peng et al. (2021)
4	<i>Hep. murlen</i>	8	3–5	9–9	1	3	1	1	13	light sandbrown	present, discontinuous	India	Lalremanga et al. (2022)
5	<i>Hep. pealii</i>	9	4–5	10	1	3 (2)	2	2	13–21	pale with brown patches	continuous	India	Smith (1943); Das et al. (2020)
6	<i>Hep. playceps</i>	8	3–5	8 or 9	1	3	1–2	1–2 (3)	18–22	usually immaculate	present or absent	India, Nepal, Bhutan, China, Bangladesh	Malnate (1966); our data
7	<i>Hep. sieboldii</i>	8	3–5	10	1	3	1–2	1–2	19–23	darkened posteriorly	usually absent	Pakistan, India, Nepal, Bhutan, Bangladesh	Malnate (1966); our data
8	<i>Hep. tpsier</i>	8 or 9	3–5 or 4–6	9–10 (8)	1	3 (2)	1–2	2 (1)	20–21	reddish orange	present, discontinuous	China	Ren et al. (2022)
9	<i>Heb. khasiensis</i>	9	4–6	10 (9)	1	2	1 (2)	2 (1)	21–26	whitish or yellowish	usually absent	India, Myanmar, China, Thailand	our data
10	<i>Heb. khasiensis</i>	9	4–6	9 or 10	1	3	1	1	N/a	whitish	barely present, discontinuous	Myanmar	our data
11	<i>Hep. cf. sieboldii</i>	9	3–5 or 4–5	9–9	1	2	1	2	N/a	whitish	absent	Myanmar	our data

khasiensis with pending further molecular data for confirming its taxonomic status. The taxonomic status of the *Hebius khasiensis-boulengeri* complex of species is being re-evaluated in another project (T.V. Nguyen *et al.* unpubl. data).

In contrast, based on the morphological and genetic distinctiveness of specimens of the Rakhine population from other congeners, we have no hesitation in considering them representing a new species of the genus *Herpetoreas* described below:

Class Reptilia Laurenti, 1768
Order Squamata Oppel, 1811
Suborder Serpentes Linnaeus, 1758
Family Natricidae Bonaparte, 1838
Genus *Herpetoreas* Günther, 1860

***Herpetoreas davidi* sp. nov.**

[urn:lsid:zoobank.org:act:4A77C927-AC6B-4367-8813-B0D172F99B42](https://lsid.zoobank.org/act:4A77C927-AC6B-4367-8813-B0D172F99B42)

Fig. 5; Tables 1–2

Amphiesma xenura (non *Herpetoreas xenura* Wall, 1907) — Wogan *et al.* 2008: 87, in part. — Das 2010: 335, in part. — Reza 2010: 64, in part. — Wallach *et al.* 2014: 35, in part.

Herpetoreas xenura — Ren *et al.* 2022: 79, in part. — Uetz *et al.* 2023: page “*Herpetoreas xenura*”, in part.

Herpetoreas sp. — Deepak *et al.* 2022: 285, in part. — Patel *et al.* 2023: 431 in supplementary materials.

Diagnosis

Herpetoreas davidi sp. nov. differs from all other members of the genus *Herpetoreas* by the combination of the following morphological characters: (1) body cylindrical, small size, max. TL 552 mm; (2) tail relatively long in both sexes, TaL/TL 0.301–0.310; (3) dorsal scale rows 19–19–17; all strongly keeled; (4) ventrals 151–156; cloacal plate single, subcaudals single; subcaudals 97–100; (5) supralabials 9, 4th–6th entering orbit; (6) head with a faint off-white sagittal line just behind the parietal; (7) dorsum dark olive-brown or dark brown with indistinct series of pale spots; (8) venter off-whitish, each ventral decorated with a pair of indistinct and faint dark-brown spots on their lateral edges.

Etymology

The species epithet is an eponym in honour of Patrick David (MNHN, Paris, France), a French reptile taxonomist, for his contribution to the systematics of snakes of the family Natricidae and to the Asian snake fauna in general. We suggest “Rakhine Keelback” as its English common name.

Type material

Holotype

MYANMAR • ♂ adult; Daung Stream, within Rakhine Yoma Elephant Sanctuary, Gwa Township, Rakhine State; 17.5844166667° N, 94.6812777778° E, WGS 84; ca 120 m a.s.l.; 23 Jan. 2002; G.O.U. Wogan and S.W. Kyi leg.; CAS 222969.

Paratypes

MYANMAR • 1 ♀ adult, 1 ♀ subadult, 1 ♂ subadult; Kyat Stream, Rakhine Yoma Elephant Sanctuary, Gwa Township, Rakhine State; 17.7038888889° N, 94.6484166667° E, WGS 84; ca 175 m a.s.l.; 26–27 and 29 Apr. 2021, respectively; J.B. Slowinski, G.O.U. Wogan, H. Win, T. Thin, and K.S. Lwin leg.; CAS 220256, CAS 220332, CAS 220378, respectively • 1 ♂ adult; Daung Stream, Rakhine Yoma

Elephant Sanctuary, Gwa Township, Rakhine State; 17.5841833333° N, 94.67815° E, WGS 84; ca 125 m a.s.l.; 1 May 2021; J.B. Slowinski, G.O.U. Wogan, H. Win, T. Thin, and K.S. Lwin leg.; CAS 220550.

Description of the holotype (Fig. 5)

MORPHOLOGY. Body slender, cylindrical, and elongate (SVL 384 mm, TaL 168 mm, TL 552 mm, ratio TaL/TL 0.304). Head moderately large, moderately distinct from neck (ratio HL/SVL 0.048, ratio HW/HL 0.512), moderately distinct from neck (Fig. 5E–F); snout broad, obtuse as seen from above; eye large (ED 2.86 mm, ratio ED/E–Sn 1.036, ratio ED/E–Ns 2.000); pupils round; nostrils crescentic, piercing in the middle of the nasal, oriented laterally (Fig. 5A–B).

BODY SCALATION. Dorsal scales in 19–19–17 rows; all scales strongly keeled, notched at apex; 154 ventrals (plus 2 preventrals); cloacal plate single; 99 subcaudals, all single.

HEAD SCALATION. Rostral wider than high, width approximately twice as long as high, visible from above (Fig. 5D); nasals pentagonal, slightly elongate, completely divided, anterior part slightly smaller than the posterior one; nostrils located on the middle of nasals (Fig. 5A–B); internasals subtriangular, in broad contact with each other, not in contact with loreal, about equal length to width, slightly, but distinctly narrowing anteriorly, the width of the posterior margin approximately twice as long as the anterior margin; 2 prefrontals, hexagonal, slightly broader than long, about 1.3 times as long as internasals; prefrontal sutures slightly longer than length internasal sutures; frontal pentagonal, elongate, length 1.4 times as long as width, about twice as long as prefrontal; supraocular 1/1, rectangular, elongate, narrowed anteriorly, broadly in contact with prefrontal, with supraocular two-thirds as broad as the frontal; parietals in broad contact with each other, parietal suture subequal to length of frontal (Fig. 5D). Loreal 1/1, subrectangular, small, about equal in length to depth, in contact with 2nd–3rd supralabials, not entering orbit; preoculars 1/1; postoculars 3/3, size decreasing from top to bottom; supralabials 9/9, 2nd–3rd in contact with the loreal, 4th–6th entering orbit, 8th largest; temporals 2+3 on both sides; mental subtriangular, wide, 2.5 times as long as high (Fig. 5A–B); 9/10 infralabials, first pair in contact with each other behind mental, 1st–5th in contact with anterior chin shields; two pairs of chin shields; posterior chin shields 1.3 times as long as anterior ones, separated from each other by small scales; mental groove apparent (Fig. 5C).

Coloration in preservative

Dorsal surface of head dark brown, with two small and white stripes run from the posterior edge of the 8th supralabials to the nape. Ventral surface of head uniform whitish. Dorsal surface of body and tail pale dark brown with irregular brown speckling on either side. Ventral surface of body and tail whitish with scales bordered on either side by a blackish-brown stripe. According to CAS staff, no photographs of living specimens of the type series of *Herpetoreas davidi* sp. nov. were taken.

Variation (for details of information's see Table 1 and Fig. 5)

The longest known specimen is 575 mm long (SVL 397 mm, TaL 178 mm, female; CAS 220550).

BODY SCALATION. 19–19–17 DSR; 151–156 VEN, 97–100 SC, single in all examined specimens, total number of VEN+SC 248–253, without sexual dimorphism. SL: 9 in all occurrences, IL: 9–10.

MAIN CHARACTERS OF PATTERN. Head olive-brown with two small, faint off-white spots on the parietals and a faint off-white sagittal line just behind the parietal. Dorsum dark olive-brown or dark brown with indistinct narrow blackish cross-bars or series of spots; venter off-whitish or cream, each ventral decorated with a pair of dark brown square spots on their lateral edges.




Fig. 5. *Herpetoreas davidi* sp. nov. A–F. Holotype, ♂ (CAS 222969) in preservation. **A.** Lateral view of the head, right side. **B.** Lateral view of the head, left side. **C.** Ventral view of the head. **D.** Dorsal view of the head. **E.** General dorsal view. **F.** General ventral view. **G–H.** Paratype, ♀ (CAS 220256). **G.** General dorsal view. **H.** General ventral view. Photos by G. Vogel.



Fig. 6. A–F. *Herpetoreas xenura* (Wall, 1907) in life. A–C. Mizoram, India. D–F. Kaptai National Park, Chittagong, Bangladesh. G–I. *Hebius khasiensis* (Boulenger, 1890) in life. G. Tamdi, Mizoram, India. H. Bhamo, Kachin, Myanmar. I. Nakhon Thai, Phitsanulok, Thailand. Photos by G. Vogel (A–C), Reza (2010) (D–E), P. Shinde (G), K.C. Ouyang (H), and P. Pawangkhanant (I).

Comparison

We summarize the main characters separating *Herpetoreas davidi* sp. nov. from the other seven species of the genus *Herpetoreas* in Table 2. More specifically, the comparisons with these species are as follows:

Herpetoreas davidi sp. nov. is easily distinguished from *H. burbrinki*, *H. murlen*, *H. pealii*, *H. platyceps*, *H. sieboldii*, and *H. tpsen* by having single subcaudals vs divided subcaudals. Particularly, *Herpetoreas davidi* differs from *H. burbrinki*, *H. murlen*, *H. platyceps*, *H. sieboldii*, and *H. tpsen* by having a single cloacal plate (vs divided) and fewer ventrals VEN 151–156 (vs 169–172 in *H. burbrinki*, 179 in *H.*




Fig. 7. Map showing the distribution of *Herpetoreas davidi* sp. nov. in Rakhine Yoma Elephant Sanctuary, Rakhine, Myanmar (red stars). 1. Daung Stream (type locality). 2. Kyat Stream. *Herpetoreas xenura* (Wall, 1907) sensu stricto in red circles, *Hebius khasiensis* (Boulenger, 1890) in blue diamonds, and *Hebius cf. khasiensis* in blue triangle. Our estimated distribution range for *Herpetoreas xenura* sensu stricto is shaded in red. See Appendix IV for the corresponding locality numbers of the other taxa presented in the map.

murlen, 191–234 in *H. platyceps*, and 174–207 in *H. sieboldii*); differs from *H. pealii* by having more ventrals VEN 151–156 (vs 136–144); and differs from *H. tpser* by having a smaller size in both sexes (max TL 552 mm in males, 575 mm in females vs 584 mm in males, 679 mm in females).

Finally, *Herpetoreas davidi* sp. nov. can be distinguished from the superficially similar species *H. xenura* by having a smaller size in both sexes (max TL 552 mm in males, 575 mm in females vs 870 mm in males, 754 mm in females), higher ratio TaL/TL in both sexes (0.301–0.304 in males, 0.307–0.310 in females vs 0.220–0.283 in males, 0.226–0.279 in females), and slightly lower number of VEN in both sexes (154–156 in males, 151–155 in females vs 158–165 in males, 155–165 in females).

Distribution and natural history notes

Herpetoreas davidi sp. nov. is so far known only from mountain forests in two locations within Rakhine Yoma Elephant Sanctuary, Gwa Township, Rakhine State, Myanmar (Fig. 7; Appendix 4). Further studies in neighbouring areas, especially in the northern Rakhine Hills, Myanmar, are required to determine the actual distribution and population size and trend of this new species. According to data recorded for specimens deposited in the collections of CAS (<http://portal.vertnet.org/search?q=Amphiesma+xenura+CAS+Myanmar>), specimens of *Herpetoreas davidi* were collected both during the day and at night on the ground near streams. The air temperature varied from 28°C to 31°C, and relative humidity was 94%. This species seems to be distributed at low elevations. The stomach of the holotype CAS 222969 contained a frog of the family Ranidae Batsch, 1796.

Conservation status

Herpetoreas davidi sp. nov. is currently known only from Rakhine Yoma Elephant Sanctuary, Gwa Township, Rakhine State, Myanmar, a protected area. In absence of other biological data to assess its population status and potential threat, we suggest that the species should be considered a Data Deficient (DD) species, following the IUCN's Red List categories (IUCN Standards and Petitions Committee 2019).

Discussion

In this study, we re-evaluate the taxonomy of the species *Herpetoreas xenura* previously reported in Myanmar by Wogan *et al.* (2008). The evidence from the molecular inferences and morphology clearly signified that the *Herpetoreas xenura* population in Rakhine State sensu Wogan *et al.* (2008) is distinct from the other nominal species of *Herpetoreas* that we describe here as *Herpetoreas davidi* sp. nov. Furthermore, the new species *Herpetoreas davidi* is completely geographically isolated from the species of *Herpetoreas* (mainly distributed in the Himalayan foothills), thus also supporting the validity of the new species. It is the most southern representative of the genus *Herpetoreas*, the closest distance to the north to the genus *Herpetoreas* (represented by *Herpetoreas xenura* in Pala Lake Area, Mizoram State, India, location 12 in Fig. 7) is more than 500 km. Meanwhile, the other population previously identified as *Herpetoreas cf. xenura* in Sagaing Region should be referred to as *Hebius khasiensis*.

Despite some recent progress in the taxonomy of the genus *Herpetoreas*, the diversity within this genus is still underestimated (Lalremsanga *et al.* 2022; Ren *et al.* 2022). For instance, similar to the case of *Herpetoreas cf. xenura* in Myanmar (Wogan *et al.* 2008), we strongly presume that the record of *Herpetoreas cf. sieboldii* from Taunggyi City, Shan State, Myanmar cited by Malnate (1966) based on female specimen BNHS 1567 (formerly BNHS 80.4) is also ambiguous. Re-examination of this specimen BNHS 1567 (by O. Adhikari, BNHS) revealed a higher total number of VEN+SC (310 vs 279–287 in *H. sieboldii*, see Table 2). Furthermore, this locality in Shan State is also reasonably isolated from the range of the true *Herpetoreas sieboldii* that inhabits the foothills of the Himalayas. This locality

is at about 880 km southeast from the type locality of *H. sieboldii*. Therefore, we stress the need for additional specimens to clarify its taxonomic status (see also Ren *et al.* 2022).

The present ASAP analysis produced 10 unidentical species partitions, and a total of 50 species are inferred by the best partition selected. Notably, this method detected *H. venningi* (MZMU 1313) from Mizoram State, India as distinct to *Hebius venningi* (CAS 233206) from Kachin State, Myanmar, with a genetic divergence of 3.8% between them; *Hebius khasiensis* (MZMU 1304) from Mizoram State, India, as distinct to *Hebius khasiensis* (CAS 221504) from Kachin State, with a genetic divergence of 3.8% between them; and *Herpetoreas xenura* (AD 633) from Assam State, India, as distinct to the other three conspecific specimens (PUCZM/X/SL1, 16–17) from Mizoram State, with a genetic divergence of 3.8% each from them. Comparably, inter-specific genetic divergence ranged between 3.8% and 46.2% across the present dataset excluding the outgroup taxon (see Appendix 4). The substantially high inter-specific genetic divergence (46.2%) estimated from *Sahyadriophis beddomei* with respect to three species of *Hebius*, viz. *H. craspedogaster* (Boulenger, 1899), *H. concularus* (Malnate, 1963), and *H. leucomystax* (David, Bain, Nguyen, Orlov, Vogel, Vu & Ziegler, 2007), further corroborates the generic status of the recently erected *Sahyadriophis* (Patel *et al.* 2023). Henceforth, our genetic divergence estimation reveals the absence of a barcode gap in the dataset and further evinces cryptic diversity within *Hebius venningi*, *H. khasiensis*, and *Herpetoreas xenura* by taking into account their respective intra-specific genetic divergence (3.8%) and the species inferences in ASAP. The aforesaid evidence necessitates a systematic review particularly for these three taxa and will be addressed in a future study with more holistic data (T.V. Nguyen *et al.* unpubl. data). Concerning *Herpetoreas sieboldii* and *H. platyceps*, two widely distributed species that received controversies in the past and little interest from taxonomists, the best ASAP partition inferred the two specimens of *Herpetoreas platyceps* (KIZ YPX26317, WII AD.R183) as two distinct species despite the fact that the genetic divergence-based PCoA ordination and phylogenetic inferences suggested their close relationship with a shallow genetic differentiation between them. Nonetheless, seeing the minimum inter-specific genetic divergence estimated across the dataset (3.8%), the genetic divergence between them (7.7%) propounded to refer the specimen of *Herpetoreas platyceps* (WII AD.R183) to as *H. cf. sieboldii*. Conforming to our molecular inferences, we further propose to amend the status of the following specimens: *Amphiesma stolatum* (Linnaeus, 1758) (USNM 524073) into *Hebius clerki* (Wall, 1925), *Amphiesma sauteri* (Boulenger, 1909) (FMNH 232808) into *Hebius maximus* (Malnate, 1962), *Amphiesma craspedogaster* (YBU 071128) into *Herpetoreas burbrinki*, *Hebius venningi* (GP 2468, from Chiang Mai, Thailand) into *Hebius deschauenseei* (Taylor, 1934) [a re-identification in full agreement with David *et al.* 2021; Hauser *et al.* 2022], *Hebius venningi* (ZMMU NAP-08395) into *Hebius nigriventer* (Wall, 1925) [a re-identification in full agreement with David *et al.* 2021], *Hebius octolineatus* (Boulenger, 1904) (GP 1242) into *Hebius johannis* (Boulenger, 1908), *Hebius craspedogaster* (GP 1712) into *Hebius metusia* (Inger, Zhao, Shaffer & Wu, 1990), *Hebius johannis* (GP 1569) into *Hebius octolineatus* (Boulenger, 1904), *Hebius* sp. (GP 1766) into *Hebius cf. ruthveni* (Van Denburgh, 1923), *Hebius* sp. (GP 1790) into *Hebius sauteri* (Boulenger, 1909), *Hebius* sp. (GP 1618) into *Hebius taronensis* (Smith, 1940), *Hebius* sp. (ZMMU NAP-06638) into *Hebius terrakarenorum* Hauser, Smits & David, 2022, *Hebius* sp. in to (AUP NAP-06639) *Hebius cf. groundwateri* (Smith, 1922), *Hebius bitaeniatum* (Wall, 1925) (GP 1940) into *Hebius* sp., and *Hebius parallelum* (Boulenger, 1890) (CHS 849; KIZ 06681) into *Herpetoreas tpser* (see Appendix 2).

As the holotype of the *Herpetoreas xenura* was lost, a neotype designation from the four samples (stored at ZSI, India) reported by Wall (1909) was required to ensure taxonomy stability of this species. In the present study, we update the diagnosis of *Herpetoreas xenura* as follows (Wall 1907, 1909; Mathew & Meetei 2004; Reza 2010; Malsawmdawngiana *et al.* 2022; Ren *et al.* 2022; and our specimens): body cylindrical, maximum TL 870 mm; tail relatively long, TaL/TL 0.220–0.283; dorsal scale rows 19–19–17; all strongly keeled; ventrals 155–165; cloacal plate single, subcaudals single; subcaudals 81–110; supralabials 9(10), 4th–6th entering orbit; maxillary teeth 22–23, gradually enlarged posteriorly,

last two distinctly enlarged, separated from anterior teeth by a small diastema; hemipenis short and thin, shallowly bilobed, extends to 8th subcaudals; sulcus spermaticus single; dorsum dark olive-brown with indistinct narrow blackish cross-bars or series of spots; venter off-white each ventral decorated with a pair of dark brown square spots on their lateral edges (see Fig. 6A–F). The known distribution of *Herpetoreas xenura* is restricted to northeastern India (from West Garo Hills and Khasi Hills in Meghalaya through Atharamura Range in Tripura, Menam Mikir Punjee in Assam, and extending towards south in Lushai Hills and up to Naga Hills towards northeast), southeastern Bangladesh (Keokradong Mountain), and possibly to the northeastern parts of Bangladesh as well as the Chin Hills of Myanmar (Fig. 7; Appendix 4). According to our observations and secondary information, we argue that there are well thriving populations of this species within its distribution range in Northeast India. Consequently, we speculate the current IUCN's categorization of this species as “Near Threatened” due to the small distribution area and reduced quality of forest freshwater habitat may be overstated (see Wogan *et al.* 2021), therefore we suggest an amendment on its conservation status into “Least Concern” species concurring to the IUCN's Red List categories (IUCN Standards and Petitions Committee 2019).

Based on literature (Lalremsanga *et al.* 2022; Ren *et al.* 2022); and our material, we propose an updated key to the genus *Herpetoreas*.

- | | |
|---|--|
| 1. Subcaudals single | 2 |
| – Subcaudals divided | 3 |
| 2. Max TL up to 575 mm, ventrals < 155 | <i>H. davidi</i> sp. nov. |
| – Max TL up to 870 mm, ventrals > 155 | <i>H. xenura</i> (Wall, 1907) |
| 3. Cloacal plate single | <i>H. pealii</i> (Sclater, 1891) |
| – Cloacal plate divided | 4 |
| 4. Ventrals fewer than 168 | <i>H. tpsen</i> Ren, Jiang, Huang, David & Li, 2022 |
| – Ventrals no less than 168 | 5 |
| 5. Tail relatively long, TaL/TL > 0.300 | <i>H. burbrinki</i> Guo, Zhu, Liu, Zhang, Li, Huang & Pyron, 2014 |
| – Tail relatively short, TaL/TL ≤ 0.300 | 6 |
| 6. Ventrals < 190 | <i>H. murlen</i> Lalremsanga, Bal, Vogel & Biakzuala, 2022 |
| – Ventrals ≥ 190 | 7 |
| 7. A higher proportion of the length of the tail with 4 supracaudal scale rows than that with 6 supracaudal scale rows high, SC4/SC6 = 1.43; weekly keeled only on five to seven mid-dorsal scale rows; VEN 205–234 [$\bar{x} = 220.8$] in males, 191–216 [$\bar{x} = 202.1$] in females | <i>H. platyceps</i> (Blyth, 1854) |
| – A lower proportion of the length of the tail with 4 supracaudal scale rows than that with 6 supracaudal scale rows high, SC4/SC6 = 0.53; all keeled but some may not be keeled in outermost mid-dorsal scale rows; VEN 193–207 [$\bar{x} = 200.3$] in males, 174–185 [$\bar{x} = 181.2$] in females | <i>H. sieboldii</i> Günther, 1860 |

Acknowledgements

T.V. Nguyen thanks Bao Nguyen Le (DTU, Vietnam) for invaluable support. We are deeply grateful to Parag Shinde (India), Ahm Reza (Bangladesh), Parinya Pawangkhanant (Thailand), and Kaichen Ouyang (China) for providing information and photos of *Herpetoreas* spp. and *Hebius* spp. We are especially thankful to Patrick David (MNHN, France), Omkar Adhikari (BNHS, India), Nikolay A.

Poyarkov (ZMMU, Russia), and Parinya Pawangkhanant (AUP, Thailand) for sharing data related to *Herpetoreas* spp. and *Hebius* spp. We thank Patrick D. Campbell (NHMUK), Lauren Scheinberg (CAS), Alain Dubois, Nicolas Vidal and Annemarie Ohler (MNHN), Andreas Schmitz (MHNG), Mark-Oliver Rödel and Frank Tillack (ZMB), Silke Schweiger, and Georg Gassner (NHMW), Pim Arntzen and Esther Dondorp (RMNH), Gunther Köhler and Linda Mogk (SMF), Jakob Hallermann (ZMH), Dennis Rödder and Wolfgang Böhme (ZFMK, Germany) for the possibility to examine specimens in the collections under their care. We also warmly thank Ngoc Quynh Nguyen (SIFASV, Vietnam) for help in the preparation of the figures. We are grateful to Patrick David (MNHN) for proofreading and linguistic help. We thank Aurélien Miralles and three anonymous reviewers for kindly reviewing a previous version of the manuscript. This research is supported in part by the Rufford Foundation (Grant No. 39897-1, data collection and analysis).

References

- Ahsan M.F., Asmat G.S.M., Islam M.A., Khan M.M.H., Muzaffar S.B., Kamruzzaman M., Chakma S. 2009. Amphibians and reptiles. In: Kabir S.M.H., Ahmed M., Ahmed A.T.A., Rahman A.K.A., Ahmed Z.U., Begum Z.N.T., Hassan M.A. & Khondker M. (eds) *Encyclopedia of Flora and Fauna of Bangladesh* Vol. 25. Asiatic Society of Bangladesh, Dhaka, Bangladesh.
- Alfaro M.E. & Arnold S.J. 2001. Molecular systematics and evolution of *Regina* and the Thamnophiine snakes. *Molecular Phylogenetics and Evolution* 21 (3): 408–423.
<https://doi.org/10.1006/mpev.2001.1024>
- Burbrink F.T., Lawson R. & Slowinski J.B. 2000 Mitochondrial DNA phylogeography of the polytypic North American rat snake (*Elaphe obsoleta*): a critique of the subspecies concept. *Evolution* 54: 2107–2118. <https://doi.org/10.1111/j.0014-3820.2000.tb01253.x>
- Chan-ard T., Nabhitabhata J. & Parr J.W. 2015. *A Field Guide to the Reptiles of Thailand*. Oxford University Press, New York.
- Che J., Jiang K., Yan F. & Zhang Y.-P. 2020. *Amphibians and Reptiles in Tibet, Diversity and Evolution*. Science Press, Beijing, China.
- Das A., Gower D.J. & Deepak V. 2020. Lost and found: rediscovery and systematics of the Northeast Indian snake *Hebius pealii* (Sclater, 1891). *Vertebrate Zoology* 70: 305–318.
<https://doi.org/10.26049/VZ70-3-2020-04>
- Das I. 2010. *A Field Guide to the Reptiles of South-east Asia*. New Holland Publishers Ltd, London.
- David P., Vogel G. & van Rooijen J. 2013 On some taxonomically confused species of the genus *Amphiesma* Duméril Bibron & Duméril 1854 related to *Amphiesma khasiense* (Boulenger 1890) (Squamata Natricidae). *Zootaxa* 3694 (4): 301–335. <https://doi.org/10.1111/zootaxa.3694.4.1>
- David P., Vogel G., Nguyen T.Q., Orlov N.L., Pauwels O.S.G., Teynié A. & Ziegler T. 2021. A revision of the dark-bellied, stream-dwelling snakes of the genus *Hebius* (Reptilia: Squamata: Natricidae) with the description of a new species from China, Vietnam and Thailand. *Zootaxa* 4911 (1): 1–61.
<https://doi.org/10.1111/zootaxa.4911.1.1>
- Deepak V., Cooper N., Poyarkov N.A., Kraus F., Burin G., Das A., Narayanan S., Streicher J.W., Smith S.-J. & Gower D.J. 2022. Multilocus phylogeny, natural history traits and classification of natricine snakes (Serpentes: Natricinae). *Zoological Journal of the Linnean Society* 1: 279–298.
<https://doi.org/10.1093/zoolinnean/zlab099>
- Dowling H.G. 1951. A proposed standard system of counting ventrals in snakes. *British Journal of Herpetology* 1: 97–99.

- Edgar R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Gower J.C. 1966. Some distance properties of latent root and vector methods used in multivariate analysis. *Biometrika* 53: 325–338.
- Günther A.C.L.G. 1864. *The Reptiles of British India*. Taylor & Francis, London.
<https://doi.org/10.5962/bhl.title.5012>
- Guo P., Liu Q., Xu Y., Jiang K., Hou M., Ding L., Pyron R.A. & Burbrink F.T. 2012. Out of Asia: natricine snakes support the Cenozoic Beringian Dispersal Hypothesis. *Molecular Phylogenetics and Evolution* 63 (3): 825–33. <https://doi.org/10.1016/j.ympev.2012.02.021>
- Guo P., Zhu F., Liu Q., Zhang L., Li J.X., Huang Y.Y. & Pyron R.A. 2014. A taxonomic revision of the Asian keelback snakes, genus *Amphiesma* (Serpentes: Colubridae: Natricinae), with description of a new species. *Zootaxa* 3873: 425–440. <https://doi.org/10.11646/zootaxa.3873.4.5>
- Hammer Ø., Harper D.A.T. & Ryan P.D. 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4 (1): 9.
- Hauser S., Smits T. & David P. 2022. On the distribution of the species of the genus *Hebius* Thompson, 1913 (Squamata: Natricidae) in North Thailand, including the description of a new species and a discussion on snake diversity of this region. *Zootaxa* 5116 (1): 1–39. <https://doi.org/10.11646/zootaxa.5116.1.1>
- Hmar G.Z., Lalmuangsanga, Lalbiakzuala, Lalruatthara, Lalrinsanga & Lalremsanga H.T. 2020. Inventory survey on the ophidian fauna of Reiek Community Reserved Forest, Mamit District, Mizoram, India. *Journal of Environmental Biology* 41: 821–826.
- Hou S.B., Yuan Z.Y., Wei P.F., Zhao G.G., Liu G.H., Wu Y.H., Shen W.J., Chen J.M., Guo P. & Che J. 2021. Molecular phylogeny and morphological comparisons of the genus *Hebius* Thompson, 1913 (Reptilia: Squamata: Colubridae) uncover a new taxon from Yunnan Province, China, and support revalidation of *Hebius septemlineatus* (Schmidt, 1925). *Zoological Research* 42 (5): 620–625.
<https://doi.org/10.24272/j.issn.2095-8137.2021.093>
- IUCN Standards and Petitions Committee 2019. *Guidelines for Using the IUCN Red List Categories and Criteria. Version 14*. Prepared by the Standards and Petitions Committee. Available from <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [accessed 4 Apr. 2024].
- Kaito T. & Toda M. 2016. The biogeographical history of Asian keelback snakes of the genus *Hebius* (Squamata: Colubridae: Natricinae) in the Ryukyu Archipelago, Japan. *Biological Journal of the Linnean Society* 118 (2): 187–199. <https://doi.org/10.1111/bij.12726>
- Kalyaanamoorthy S., Minh B.Q., Wong T.K., Von Haeseler A. & Jermiin L.S. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* 14: 587–589.
<https://doi.org/10.1038/nmeth.4285>
- Lalremsanga H.T., Bal A.K., Vogel G. & Biakzuala L. 2022. Molecular phylogenetic analyses of lesser known colubrid snakes reveal a new species of *Herpetoreas* (Squamata: Colubridae: Natricinae), and new insights into the systematics of *Gongylosoma scriptum* and its allies from northeastern India. *Salamandra* 58 (2): 101–115.
- Lalironunga S., Lalrinchhana C., Vanramliana V., Das A., Gower D.J. & Deepak V.A. 2020. A multilocus molecular perspective on the systematics of the poorly known Northeast Indian colubrid snakes *Blythia reticulata* (Blyth, 1854), *B. hmuifang* Vogel, Lalremsanga & Vanlalhrima, 2017, and *Hebius xenura* (Wall, 1907). *Zootaxa* 4768: 193–200. <https://doi.org/10.11646/zootaxa.4768.2.2>

Lanfear R., Frandsen P.B., Wright A.M., Senfeld T. & Calcott B. 2016. PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* 34: 772–773. <https://doi.org/10.1093/molbev/msw260>

Li J.N., Liang D., Wang Y.Y., Guo P., Huang S. & Zhang P. 2020. A large-scale systematic framework of Chinese snakes based on a unified multilocus marker system. *Molecular Phylogenetics and Evolution* 148: 106807. <https://doi.org/10.1016/j.ympev.2020.106807>

Li M.L., Ren L.L., Huang J.J., Lyu Z.T., Qi S., Jiang K., Wang Y.Y. & Li J.T. 2022. On the validity of *Hebius sauteri maximus* (Malnate, 1962) (Squamata, Natricidae), with the redescription of *H. maximus* comb. nov. and *H. sauteri* (Boulenger, 1909). *Herpetozoa* 35: 265–282.

<https://doi.org/10.3897/herpetozoa.35.e94920>

Liu Q., Zhong G.H., Wang P., Liu Y. & Guo P. 2018. A new species of the genus *Hebius* (Squamata: Colubridae) from Sichuan, China. *Zootaxa* 4483 (2): 385–394.

<https://doi.org/10.11646/zootaxa.4483.2.10>

Malnate E.V. 1966. *Amphiesma platyceps* (Blyth), and *Amphiesma sieboldii* (Guenther): sibling species (Reptilia: Serpentes). *Journal of Bombay Natural History Society* 63: 1–17.

Available from <https://www.biodiversitylibrary.org/page/47950335> [accessed 4 Apr. 2024].

Malsawmdawngiana, Boruah B., Patel N.G., Lalronunga S., Zosangliana I., Lalhmangaiha K. & Das A. 2022. An updated checklist of reptiles from Dampa Tiger Reserve, Mizoram, India, with sixteen new distribution records. *Journal of Threatened Taxa* 14 (10): 21946–21960.

<https://doi.org/10.11609/jott.8004.14.10.21946-21960>

Mathew R. 1995. Reptilia. In: Ghosh A.K. (ed.) *Fauna of Meghalaya. Part 1. Vertebrates*: 379–454. State Fauna Series 4, Zoological Survey of India, Calcutta.

Mathew R. & Meetei A.B. 2004. On the identity of *Amphiesma venningi* (Wall, 1910) reported from Meghalaya, India. *Hamadryad* 29 (1): 134–135.

Minh B.Q., Nguyen M.A.T. & von Haeseler A. 2013. Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30: 1188–1195. <https://doi.org/10.1093/molbev/mst024>

Muangsanga L., Laltlanhlui L.H., Biakzuala L., Rathee Y.S. & Lalremsanga H.T. 2021. Observations of feeding behavior and a note on clutch size in Wall's Keelback, *Herpetoreas xenura* (Wall 1907) (Squamata: Natricidae). *Reptiles & Amphibians* 28 (1): 82–83. <https://doi.org/10.17161/randa.v28i1.15324>

Nguyen L.T., Schmidt H.A., von Haeseler A. & Minh B.Q. 2015. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum likelihood phylogenies. *Molecular Biology and Evolution* 32: 268–274. <https://doi.org/10.1093/molbev/msu300>

Patel H., Thackeray T., Campbell P.D. & Mirza Z.A. 2023. Systematic assessment of *Hebius beddomei* (Günther, 1864) (Serpentes: Colubridae: Natricinae) with description of a new genus and a new allied species from the Western Ghats, India. *Taxonomy* 3: 415–434. <https://doi.org/10.3390/taxonomy3030024>

Pawar S. & Birand A. 2001. *A Survey of Amphibians, Reptiles, and Birds in Northeast India*. Centre for Ecological Research and Conservation, Pune, India.

Peng L.F., Huang S., Burbrink F.T., Zhang Y., Guo P. & Wu H.L. 2021. Morphological description of a new specimen of *Herpetoreas burbrinki* Guo et al. 2014 (Serpentes: Colubridae). *Zootaxa* 5039: 433–439. <https://doi.org/10.11646/zootaxa.5039.3.8>

Poyarkov N.A., Nguyen T.V., Popov E.S., Geissler P., Pawangkhanant P., Neang T., Suwannapoom C., Ananjeva N.B. & Orlov N.L. 2023. Recent progress in taxonomic studies, biogeographic analysis and revised checklist of Reptilians in Indochina. *Russian Journal of Herpetology* 30 (5): 255–476.

<https://doi.org/10.30906/1026-2296-2023-30-5-255-476>

- Puillandre N., Brouillet S. & Achaz G. 2021. ASAP: Assemble Species by Automatic Partitioning. *Molecular Ecology Resources* 21: 609–620. <https://doi.org/10.1111/1755-0998.13281>
- Rambaut A., Drummond A.J., Xie D., Baele G. & Suchard M.A. 2018. Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67: 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Ren J.L., Wang L., Nguyen T.T., Hoang C.V., Zhing G.H., Jiang K., Guo P. & Li J.-T. 2018. Taxonomic re-evaluation of the monotypic genus *Pararhabdophis* Bourret, 1934 (Squamata, Colubridae, Natricinae) with discovery of its type species, *P. chapaensis*, from China. *Zootaxa* 4486 (1): 31–56. <https://doi.org/10.11646/zootaxa.4486.1.2>
- Ren J.L., Jiang K., Huang J.J., David P. & Li J.T. 2022. Taxonomic review of the genus *Herpetoreas* (Serpentes: Natricidae), with the description of a new species from Tibet, China. *Diversity* 14: 79. <https://doi.org/10.3390/d14020079>
- Reza A. 2010. First record of *Amphiesma vennungi* (Wall, 1910) (Serpentes, Colubridae, Natricinae) from Bangladesh, with notes on its taxonomy, natural history, biogeography and other sympatric species. *Hamadryad* 35: 64–72.
- Romer J. 1945. A new record of a rare snake (*Natrix xenura*) from Assam. *The Journal of the Bombay Natural History Society* 45: 430–431.
Available from <https://www.biodiversitylibrary.org/page/48127182> [accessed 25 Apr. 2024].
- Ronquist F., Teslenko M., Van Der Mark P., Ayres D.L., Darling A., Höhna S., Larget B., Liu L., Suchard M.A. & Huelsenbeck J.P. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Smith M.A. 1943. *The Fauna of British India, Ceylon and Burma, including the Whole of the Indo-Chinese Subregion. Reptilia and Amphibia. Vol. III. Serpentes*. Taylor & Francis, London, UK.
- Tamura K., Stecher G. & Kumar S. 2021. MEGA11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution* 38: 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Uetz P., Freed P. & Hošek J. 2023. *The Reptile Database*. (September 2023). Electronic database. Available from <http://reptile-database.reptarium.cz/> [accessed 4 Apr. 2024].
- Wall F. 1907. Notes on snakes from the neighbourhood of Darjeeling. *The Journal of the Bombay Natural History Society* 19: 337–357. Available from <https://www.biodiversitylibrary.org/page/5449715> [accessed 4 Apr. 2024].
- Wall F. 1909. Some new Asian snakes. *The Journal of the Bombay Natural History Society* 17: 612–619. Available from <https://www.biodiversitylibrary.org/page/30119822> [accessed 4 Apr. 2024].
- Wallach V., Williams K.L. & Boundy J. 2014. *Snakes of the World. A Catalogue of Living and Extinct Species*. CRC Press, Boca Raton, FL, USA.
- Wogan G.O.U., Vindum J.V., Wilkinson J.A., Koo M.S., Slowinski J.B., Win H., Thin T., Kyi S.W., Oo S.L. & Lwin K.S. 2008. New country records and range extensions for Myanmar amphibians and reptiles. *Hamadryad* 33: 83–96.
- Wogan G.O.U., Ghosh A. & Khan M. 2021. *Hebius xenura*. *The IUCN Red List of Threatened Species* 2021: e.T192012A2027882. <https://doi.org/10.2305/IUCN.UK.2021-3.RLTS.T192012A2027882.en>
- Wu Y.H., Hou S.B., Yuan Z.Y., Jiang K., Huang R.Y., Wang K., Liu Q., Yu Z.B., Zhao H.P., Zhang B.L., Chen J.M., Wang L.J., Stuart B.L., Chambers E.A., Wang Y.F., Gao W., Zou D.H., Yan F., Zhao G.G., Fu Z.X., Wang S.N., Jiang M., Zhang L., Ren J.L., Wu Y.Y., Zhang L.Y., Yang D.C., Jin J.Q., Yin T.T.,

Li J.T., Zhao W.G., Murphy R.W., Huang S., Guo P., Zhang Y.P. & Che J. 2023. DNA barcoding of Chinese snakes reveals hidden diversity and conservation needs. *Molecular Ecology Resources* 23: 1124–1141. <https://doi.org/10.1111/17550998.13784>

Xu W., Dong W.J., Fu T.T., Gao W., Lu C.Q., Yan F., Wu Y.H., Jiang K., Jin J.Q., Chen H.M., Zhang Y.P., Hillis D.M. & Che J. 2021. Herpetological phylogeographic analyses support a Miocene focal point of Himalayan uplift and biological diversification. *National Science Review* 8 (9): nwaa263. <https://doi.org/10.1093/nsr/nwaa263>

Xu Y.H., Yang D.C., Gong Y.A., Wu J.X., Huang R.Y., Liu Y.J., Liang S.M., Huang T.Q. & Huang S. 2023. A new species of the genus *Hebius* Thompson, 1913 (Squamata: Colubridae) from Baise, Guangxi, China. *Zootaxa* 5319 (1): 76–90. <https://doi.org/10.11646/zootaxa.5319.1.5>

Yang D. & Rao D.Q. 2008 *Amphibia and Reptilia of Yunnan Kunming*. Yunnan Publishing Group Corporation, Yunnan Science. [In Chinese.]

Zhou Z., Sun Z., Qi S., Lu Y., Lyu Z., Wang Y., Li P. & Ma J. 2019. A new species of the genus *Hebius* (Squamata: Colubridae: Natricinae) from Hunan Province, China. *Zootaxa* 4674 (1): 68–82.

<https://doi.org/10.11646/zootaxa.4674.1.3>

Ziegler T., Luu V.Q., Nguyen T.T., Ha N.V., Ngo H.T., Le M.D. & Nguyen T.Q. 2019. Rediscovery of Andrea's keelback, *Hebius andreae* (Ziegler & Le, 2006): first country record for Laos and phylogenetic placement. *Revue suisse de Zoologie* 126 (1): 61–71. <https://doi.org/10.5281/zenodo.2619520>

Manuscript received: 4 September 2023

Manuscript accepted: 9 February 2024

Published on: 3 May 2024

Topic editor: Magalie Castelin

Section editor: Aurélien Miralles

Desk editor: Pepe Fernández

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic.

Supplementary files

Supp. file 1. Maximum Likelihood (ML) phylogenetic tree based on the fragment of mitochondrial Cytochrome b (cytb) gene among the genera *Amphiesma* Duméril, Bibron & Duméril, 1854, *Amphiesmoides* Malnate, 1961, *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023. The ultrafast bootstrap (UFB) support values are given at each branch. <https://doi.org/10.5852/ejt.2024.932.2519.11303>

Supp. file 2. *Herpetoreas* cf. *sieboldii* (Günther, 1860), specimen ♀ (BNHS 1567). **A.** Lateral view of the head, right side. **B.** Lateral view of the head, left side. **C.** Dorsal view of the head. **D.** Ventral view of the head. **E.** General dorsal view. **F.** General ventral view. Photos by O. Adhikari.

<https://doi.org/10.5852/ejt.2024.932.2519.11305>

Appendix 1 (continued on next page). List of examined specimens of *Hebius khasiensis* (Boulenger, 1890), *Herpetoreas sieboldii* (Günther, 1860), and *Her. platiceps* (Blyth, 1854).

No.	Species	Number	Location	Sex
1	<i>Hebius khasiensis</i>	NHMUK 1925.4.2.11	Putao, Kachin, Myanmar	M
2	<i>Hebius khasiensis</i>	NHMUK 1925.4.2.13	Putao, Kachin, Myanmar	M
3	<i>Hebius khasiensis</i>	NHMUK 1925.4.2.14	Putao, Kachin, Myanmar	M
4	<i>Hebius khasiensis</i>	NHMUK 1946.1.12.81 (syntypes of <i>Tropidonotus khasiensis</i>)	Khasi Hills, Meghalaya, India	M
5	<i>Hebius khasiensis</i>	NHMUK 1946.1.12.82 (syntypes of <i>Tropidonotus khasiensis</i>)	Khasi Hills, Meghalaya, India	M
6	<i>Hebius khasiensis</i>	NHMUK 1946.1.13.62 (holotype of <i>Natrix gilhodesi</i>)	Putao, Kachin, Myanmar	M
7	<i>Hebius khasiensis</i>	NHMUK 1946.1.13.63	Putao, Kachin, Myanmar	M
8	<i>Hebius khasiensis</i>	CAS 220023	Nat Ma Taung NP., Chin, Myanmar	M
9	<i>Hebius khasiensis</i>	CIB 2000 I 0009	Ruili, Dehong, Yunnan, China	M
10	<i>Hebius khasiensis</i>	FMNH 251781	Chiang Mai, Thailand	M
11	<i>Hebius khasiensis</i>	KSC 140	Kohima, Nagaland, India	M
12	<i>Hebius khasiensis</i>	NRCT 980504	Doi Inthanon NP., Chiang Mai, Thailand	M
13	<i>Hebius khasiensis</i>	378 P	Dibang, Arunachal Pradesh, India	M
14	<i>Hebius khasiensis</i>	NHMUK 1925.4.2.10	Putao, Kachin, Myanmar	F
15	<i>Hebius khasiensis</i>	NHMUK 1925.4.2.12	Putao, Kachin, Myanmar	F
16	<i>Hebius khasiensis</i>	NHMUK 1925.4.2.15	Putao, Kachin, Myanmar	F
17	<i>Hebius khasiensis</i>	NHMUK 1925.4.2.15A	Putao, Kachin, Myanmar	F
18	<i>Hebius khasiensis</i>	NHMUK 1946.1.12.80	Khasi Hills, Meghalaya, India	F
19	<i>Hebius khasiensis</i>	NHMUK 1946.1.13.45	Khasi Hills, Meghalaya, India	F
20	<i>Hebius khasiensis</i>	NHMUK 1974.884	Putao, Kachin, Myanmar	F
21	<i>Hebius khasiensis</i>	CAS 221504	Putao, Kachin, Myanmar	F
22	<i>Hebius khasiensis</i>	CAS 221525	Putao, Kachin, Myanmar	F
23	<i>Hebius khasiensis</i>	CAS 221543	Putao, Kachin, Myanmar	F
24	<i>Hebius khasiensis</i>	CAS 224654	Putao, Kachin, Myanmar	F
25	<i>Hebius khasiensis</i>	CAS 224694	Putao, Kachin, Myanmar	F
26	<i>Hebius khasiensis</i>	CIB Xi0089	Xichuangbanna, Yunnan, China	F
27	<i>Hebius khasiensis</i>	FMNH 251780	Chiang Mai, Thailand	F
28	<i>Hebius khasiensis</i>	MNHN 1893.0399	Karen Hills, Shan, Myanmar	F
29	<i>Hebius khasiensis</i>	MNHN 2004.0248	Boun Tay, Phongsaly, Laos	F
30	<i>Hebius khasiensis</i>	QSMI 0273	Phu Luang, Loei, Thailand	F
31	<i>Hebius khasiensis</i>	ZSI 23926	Changlang, Arunachal Pradesh, India	F
32	<i>Hebius khasiensis</i>	377 P	Dibang, Arunachal Pradesh, India	F
33	<i>Hebius khasiensis</i>	379 P	Dibang, Arunachal Pradesh, India	F
34	<i>Herpetoreas sieboldii</i>	NHMUK 1913.5.22.1	Maikola Valley, Nepal	M
35	<i>Herpetoreas sieboldii</i>	CAS 177672	Tibet, China	M
36	<i>Herpetoreas sieboldii</i>	CAS 177673	Tibet, China	M

Appendix 1 (continued). List of examined specimens of *Hebius khasiensis* (Boulenger, 1890), *Herpetoreas sieboldii* (Günther, 1860), and *Her. platyceps* (Blyth, 1854).

No.	Species	Number	Location	Sex
37	<i>Herpetoreas sieboldii</i>	CAS 90690	Nepal	M
38	<i>Herpetoreas sieboldii</i>	FMNH 204502	Nepal	M
39	<i>Herpetoreas sieboldii</i>	FMNH 204504	Nepal	M
40	<i>Herpetoreas sieboldii</i>	FMNH 190856	Nepal	M
41	<i>Herpetoreas sieboldii</i>	FMNH 131966	Nepal	M
42	<i>Herpetoreas sieboldii</i>	FMNH 204499	Nepal	M
43	<i>Herpetoreas sieboldii</i>	FMNH 204501	Nepal	M
44	<i>Herpetoreas sieboldii</i>	NHMW 22383:1	Darjeeling, West Bengal, India	M
45	<i>Herpetoreas sieboldii</i>	NHMW 22383:3	Darjeeling, West Bengal, India	M
46	<i>Herpetoreas sieboldii</i>	NHMW 22383:4	Darjeeling, West Bengal, India	M
47	<i>Herpetoreas sieboldii</i>	CAS 15973	Darjeeling, West Bengal, India	F
48	<i>Herpetoreas sieboldii</i>	CAS 177674	Tibet, China	F
49	<i>Herpetoreas sieboldii</i>	MNHG 1355.73	Nepal	F
50	<i>Herpetoreas sieboldii</i>	FMNH 109762	Gorkha, Nepal	F
51	<i>Herpetoreas sieboldii</i>	FMNH 204500	Nepal	F
52	<i>Herpetoreas sieboldii</i>	FMNH 204503	Nepal	F
53	<i>Herpetoreas platyceps</i>	NHMW 22383:5	Darjeeling, West Bengal, India	F
54	<i>Herpetoreas platyceps</i>	NHMW 18570:1 (syntypes of <i>Zamenis himalayanus</i>)	Shimla and Kullu, Himachal Pradesh, India	F
55	<i>Herpetoreas platyceps</i>	NHMW 18570:2 (syntypes of <i>Zamenis himalayanus</i>)	Shimla and Kullu, Himachal Pradesh, India	F
56	<i>Herpetoreas platyceps</i>	NHMW 18569	Himachal Pradesh, India	F
57	<i>Herpetoreas platyceps</i>	MNHN 4197	Himachal Pradesh, India	F
58	<i>Herpetoreas platyceps</i>	RMNH 20502	Landrang, Nepal	F

Appendix 2 (continued on next two pages). Sequences of the *Herpetoreas* Günther, 1860 and related taxa used in this study. Remark: “?” = request verification.

No	Previous identification	New identification	Revised distributions	Voucher number	Cytb	Locality	References
1	<i>Amphiesma monticolum</i>	<i>Amphiesma monticola</i>	India	SN 009	OK315803	Wayanad, India	Deepak <i>et al.</i> 2022
2	<i>Amphiesma stolatum</i>	<i>Amphiesma stolatum</i>	India, Nepal, Pakistan, Sri Lanka, Myanmar, China [?] , Taiwan [?] , Thailand [?] , Vietnam [?] , Laos [?] , Cambodia [?]	GP 2239	KJ1685694	Guangdong, China	Guo <i>et al.</i> 2014; Poyarkov <i>et al.</i> 2023; our data
3	<i>Amphiesmoides ornaticeps</i>	<i>Amphiesmoides ornaticeps</i>	China, Vietnam	ZMMU R16289	OK326949	Pu Hoat, Nghe An, Vietnam	Deepak <i>et al.</i> 2022; Poyarkov <i>et al.</i> 2023;
4	<i>Hebius andreae</i>	<i>Hebius andreae</i>	Vietnam, Laos	VNUF R.2017.25	MK253674	Khammouane, Laos	Ziegler <i>et al.</i> 2019; Poyarkov <i>et al.</i> 2023
5	<i>Hebius annamensis</i>	<i>Hebius annamensis</i>	Vietnam, Laos	FMNH 258637	OK315812	Kaleum, Xekong, Laos	Deepak <i>et al.</i> 2022; Poyarkov <i>et al.</i> 2023
6	<i>Hebius atemporalis</i>	<i>Hebius atemporalis</i>	Vietnam, China	ZMMU NAP07877	OK315813	Pu Hoat, Nghe An, Vietnam	Deepak <i>et al.</i> 2022; Poyarkov <i>et al.</i> 2023
7	<i>Hebius bitaeniatus</i>	<i>Hebius bitaeniatus</i>	Myanmar, Thailand, Vietnam, China [?]	AUP 00062	OK315816	Chiang Mai, Thailand	Deepak <i>et al.</i> 2022; Poyarkov <i>et al.</i> 2023
8	<i>Hebius boulongeri</i>	<i>Hebius boulongeri</i>	China, Vietnam, Thailand [?] , Laos [?] , Cambodia [?]	GP 1789	KJ1685684	Guangdong, China	Guo <i>et al.</i> 2014; Poyarkov <i>et al.</i> 2023
9	<i>Hebius chapaensis</i>	<i>Hebius chapaensis</i>	Vietnam, Laos, China	VNMN 06102	MHT778702	Sapa, Lao Cai, Vietnam	Ren <i>et al.</i> 2018; Poyarkov <i>et al.</i> 2023
10	<i>Amphiesma stolatum</i>	<i>Hebius clerki</i>	Myanmar, China	USNM 524073	KJ1685711	Chatthini, Sagaing, Myanmar	Guo <i>et al.</i> 2014
11	<i>Hebius concularis</i>	<i>Hebius concularis</i>	Japan	KUZ R20253	AB389268	Okinawa, Ryukyu, Japan	Kaito & Toda 2016
12	<i>Hebius craspedogaster</i>	<i>Hebius craspedogaster</i>	China	GP 1963	KJ1685689	Sichuan, China	Guo <i>et al.</i> 2014
13	<i>Hebius vennigeri</i>	<i>Hebius deschauenseei</i>	Thailand	GP 2468	KJ1685700	Chiang Mai, Thailand	Guo <i>et al.</i> 2014; Poyarkov <i>et al.</i> 2023
14	<i>Hebius</i> sp.	<i>Hebius cf. groundwateri</i>	Thailand	AUP NAP-06639	OK315825	Ratchaburi, Thailand	Deepak <i>et al.</i> 2022; Poyarkov <i>et al.</i> 2023
15	<i>Hebius ishigakiensis</i>	<i>Hebius ishigakiensis</i>	Japan	KUZ R33045	AB389296	Okinawa, Ryukyu, Japan	Kaito & Toda 2016
16	<i>Hebius octolineatus</i>	<i>Hebius johannii</i>	China	GP 1242	KJ1685673	Guizhou, China	Guo <i>et al.</i> 2014
17	<i>Hebius khasiensis</i>	<i>Hebius khasiensis</i>	Myanmar, China, Thailand	CAS 221504	KJ1685668	Putao, Kachin, China	Guo <i>et al.</i> 2014; Poyarkov <i>et al.</i> 2023
18	<i>Hebius leucomystax</i>	<i>Hebius leucomystax</i>	Vietnam, Laos, Cambodia, Thailand	ZMMU R.14807	OK315821	Quang Binh, Vietnam	Deepak <i>et al.</i> 2022; Poyarkov <i>et al.</i> 2023
19	<i>Amphiesma sauteri</i>	<i>Hebius maximus</i>	China	FMNH 232808	AF402905	Hongya, Sichuan, China	Alfaro & Arnold 2001; Li <i>et al.</i> 2022
20	<i>Hebius craspedogaster</i>	<i>Hebius meiusia</i>	China	GP 1712	KJ1685682	Baoxing, Sichuan, China	Guo <i>et al.</i> 2014

Appendix 2 (continued). Sequences of the *Herpetoreas* Günther, 1860 and related taxa used in this study. Remark: “?” = request verification.

No	Previous identification	New identification	Revised distributions	Voucher number	Cytb	Locality	References
21	<i>Hebius modestum</i>	<i>Hebius modestum</i>	India, Myanmar, China	CAS 234262	KJ685671	Dehong, Yunnan, China	Guo <i>et al.</i> 2014; David <i>et al.</i> 2021
22	<i>Hebius vennungi</i>	<i>Hebius nigritreter</i>	Myanmar, China	ZMMU NAP-08395	OK315829	Kachin, Myanmar	David <i>et al.</i> 2021; Deepak <i>et al.</i> 2022
23	<i>Hebius johannis</i>	<i>Hebius octolineatus</i>	China	GP 1569	KJ685678	Yunnan, China	Guo <i>et al.</i> 2014
24	<i>Hebius optatus</i>	<i>Hebius optatus</i>	China, Vietnam?	GP 1885	KJ685687	Libo, Guangxi, China	Guo <i>et al.</i> 2014; Poyarkov <i>et al.</i> 2023
25	<i>Hebius popei</i>	<i>Hebius popei</i>	China	GP 2169	KJ685692	Qiongzhang, Hainan, China	Guo <i>et al.</i> 2014
26	<i>Hebius pryeri</i>	<i>Hebius pryeri</i>	Japan	KUZ R67983	AB989102	Ryukyu, Japan	Kaito & Toda 2016
27	<i>Hebius</i> sp.	<i>Hebius</i> cf. <i>ruthveni</i>	China	GP 1766	KJ685683	Jiangxi, China	Guo <i>et al.</i> 2014; Kaito & Toda 2016
28	<i>Hebius sangzhiensis</i>	<i>Hebius sangzhiensis</i>	China	SYNU 08070350	MK340763	Sangzhi, Hunan, China	Zhou <i>et al.</i> 2019
29	<i>Hebius</i> sp.	<i>Hebius sauteri</i>	China, Taiwan	GP 1790	KJ685685	Guangdong, China	Guo <i>et al.</i> 2014; Li <i>et al.</i> 2022
30	<i>Hebius septentrinalis</i>	<i>Hebius septentrinalis</i>	China	KIZ 037706	MZ570485	Tengchong, Yunnan, China	Hou <i>et al.</i> 2021
31	<i>Hebius</i> sp.	<i>Hebius taronensis</i>	Myanmar, India	GP1618	KJ685679	Myanmar	Guo <i>et al.</i> 2014; David <i>et al.</i> 2021
32	<i>Hebius</i> sp.	<i>Hebius terrakarenorum</i>	Thailand	ZMMU NAP-06638	OK315824	Tak, Thailand	Deepak <i>et al.</i> 2022; Poyarkov <i>et al.</i> 2023
33	<i>Hebius vennungi</i>	<i>Hebius vennungi</i>	India, Myanmar	CAS 233206	KJ685670	Phalum, Chin, Myanmar	Guo <i>et al.</i> 2014; David <i>et al.</i> 2021
34	<i>Hebius vennungi</i>	<i>Hebius vennungi</i>	MZMU1313 (isolate A)	OR389974	Mizoram, India	This study	
35	<i>Hebius vennungi</i>	<i>Hebius vennungi</i>	MZMU1313 (isolate B)	OR389975	Mizoram, India		
36	<i>Hebius vibakari</i>	<i>Hebius vibakari</i>	Japan	KUZ R21587	AB989302	Kansai, Japan	Kaito & Toda 2016
37	<i>Hebius weixiensis</i>	<i>Hebius weixiensis</i>	China	KIZ 035776	MZ570490	Weixi, Yunnan, China	Hou <i>et al.</i> 2021
38	<i>Hebius yanbianensis</i>	<i>Hebius yanbianensis</i>	China	GP 4006	MH532291	Yanbian, Sichuan, China	Liu <i>et al.</i> 2018
39	<i>Hebius bitaeniatum</i>	<i>Hebius</i> sp.	China	GP 1940	KJ685688	Guangxi, China	Guo <i>et al.</i> 2014
40	<i>Hebius youjiangensis</i>	<i>Hebius youjiangensis</i>	China	HSR 22184	OQ085073	Baise, Guangxi, China	Xu <i>et al.</i> 2023
41	<i>Amphiesma craspedogaster</i>	<i>Herpetoreas burbrinki</i>	China	YBU 071128	GQ281781	Tibet, China	Guo <i>et al.</i> 2012
42	<i>Herpetoreas burbrinki</i>	<i>Herpetoreas burbrinki</i>	China	HSR 19252	MZ358098	Tibet, China	Peng & Huang unpubl. data; Ren <i>et al.</i> 2022
43	<i>Herpetoreas xenura</i>	<i>Herpetoreas davidi</i> sp. nov.	Myanmar	CAS 220256	OK315830	Gwa, Rakhine, Myanmar	Deepak <i>et al.</i> 2022

Appendix 2 (continued). Sequences of the *Herpetoreas* Günther, 1860 and related taxa used in this study. Remark: “?” = request verification.

No	Previous identification	New identification	Revised distributions	Voucher number	Cytb	Locality	References
44	<i>Herpetoreas murlen</i>	<i>Herpetoreas murlen</i>	India	MZMU 2473	ON204025	Mizoram, India	Lalrem sangha et al. 2022
45	<i>Herpetoreas pealii</i>	<i>Herpetoreas pealii</i>	India	WII ADR547	MT571586	Arunachal Pradesh, India	Das et al. 2020
46	<i>Herpetoreas platiceps</i>	<i>Herpetoreas platiceps</i>	India, Nepal, Pakistan, Bhutan, China, Bangladesh	KIZ YPX26317	MW111464	Gyirong, Tibet, China	Xu et al. 2021
47	<i>Herpetoreas platiceps</i>	<i>Herpetoreas cf. sieboldii</i>	Pakistan, India, Nepal, Bhutan, Bangladesh	WII AD R183	MT571587	Chamoli, Uttarakhand, India	Das et al. 2020
48	<i>Hebius parallelum</i>	<i>Herpetoreas ipser</i>	China	CHS 849	MK201567	Tibet, China	Li et al. 2020
49	<i>Hebius parallelum</i>	<i>Herpetoreas ipser</i>	China	KIZ 06681	MW111476	Tibet, China	Xu et al. 2021
50	<i>Herpetoreas ipser</i>	<i>Herpetoreas ipser</i>	China	CIB 107855	OM313292	Medog, Tibet, China	Ren et al. 2022
51	<i>Herpetoreas ipser</i>	<i>Herpetoreas ipser</i>	India, Bangladesh	CIB 118524	OM313293	Medog Tibet, China	Ren et al. 2022
52	<i>Herpetoreas xenura</i>	<i>Herpetoreas xenura</i>	India, Bangladesh	PUCZMX SL1	MN993850	Mizoram, India	Lalronunga et al. 2020
53	<i>Herpetoreas xenura</i>	<i>Herpetoreas xenura</i>	India	PUCZM X SL17	MN993852	Mizoram, India	Lalronunga et al. 2020
54	<i>Herpetoreas xenura</i>	<i>Herpetoreas xenura</i>	India	PUCZM X SL16	MN993851	Mizoram, India	Lalronunga et al. 2020
55	<i>Herpetoreas xenura</i>	<i>Herpetoreas xenura</i>	India	AD 633	MN993891	Assam, India	Lalronunga et al. 2020
56	<i>Sahyadriophis beddomei</i>	<i>Sahyadriophis beddomei</i>	India	SN 013	OK315814	Valparai, India	Deepak et al. 2022

Appendix 3. The best partitioning schemes and nucleotide substitution models selected for the partitioned Bayesian Inference and Maximum Likelihood phylogenies in the present study.

Partitions	Sites	For Bayesian Inference (Partition Finder)	For Maximum Likelihood (ModelFinder)
I	Cytbpos1	TIM+I+G	TN+F+R7
II	Cytbpos2, Cytbpos3	HKY+I+G	TPM2+F+I+G4

Appendix 4 (continued on next seven pages). Uncorrected p-distance estimated using Cytb gene sequence among the genera *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, *Amphiesma* Duméril, Bibron & Duméril, 1854, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023, with *Amphiesmoides* Malnate, 1961 as outgroup.

No.	Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1	OK315830 <i>Herpetoreas davidi</i> sp. nov																		
2	ON204025 <i>Herpetoreas muren</i>	0.192																	
3	MZ358098 <i>Herpetoreas burbrinki</i>	0.231	0.154																
4	GQ281781 <i>Herpetoreas burbrinki</i>	0.231	0.154	0.000															
5	MW111464 <i>Herpetoreas platiceps</i>	0.231	0.231	0.231	0.231														
6	MT571587 <i>Herpetoreas cf. sieboldii</i>	0.231	0.231	0.231	0.231	0.077													
7	MT571586 <i>Herpetoreas peali</i>	0.269	0.308	0.385	0.385	0.269	0.269												
8	MN993850 <i>Herpetoreas xenura</i>	0.269	0.308	0.308	0.231	0.231	0.154	0.308											
9	MN993852 <i>Herpetoreas xenura</i>	0.269	0.308	0.308	0.231	0.154	0.308	0.000											
10	MN993851 <i>Herpetoreas xenura</i>	0.269	0.308	0.308	0.231	0.154	0.308	0.000	0.000										
11	MN993891 <i>Herpetoreas xenura</i>	0.308	0.346	0.346	0.269	0.269	0.192	0.346	0.038	0.038	0.038								
12	MW111476 <i>Herpetoreas tpser</i>	0.308	0.231	0.269	0.269	0.154	0.231	0.231	0.346	0.346	0.346	0.385							
13	OM313293 <i>Herpetoreas tpser</i>	0.308	0.231	0.269	0.269	0.154	0.231	0.231	0.346	0.346	0.346	0.385	0.000						
14	OM313292 <i>Herpetoreas tpser</i>	0.308	0.231	0.269	0.269	0.154	0.231	0.231	0.346	0.346	0.346	0.385	0.000	0.000					
15	MK201567 <i>Herpetoreas tpser</i>	0.308	0.231	0.269	0.269	0.154	0.231	0.231	0.346	0.346	0.346	0.385	0.000	0.000	0.000				
16	MK253674 <i>Hebius andreae</i>	0.154	0.231	0.231	0.231	0.154	0.346	0.269	0.269	0.269	0.269	0.308	0.308	0.308	0.308	0.308	0.308	0.308	
17	OK315812 <i>Hebius annamensis</i>	0.192	0.192	0.269	0.269	0.192	0.192	0.269	0.308	0.308	0.308	0.346	0.192	0.192	0.192	0.192	0.192	0.115	
18	KJ685671 <i>Hebius modestum</i>	0.192	0.192	0.269	0.269	0.192	0.192	0.269	0.308	0.308	0.308	0.346	0.192	0.192	0.192	0.192	0.115	0.000	
19	KJ685684 <i>Hebius bouengeri</i>	0.192	0.231	0.308	0.308	0.231	0.231	0.308	0.269	0.269	0.269	0.308	0.231	0.231	0.231	0.231	0.231	0.115	
20	KJ685700 <i>Hebius deschauenseei</i>	0.192	0.192	0.269	0.269	0.192	0.192	0.269	0.308	0.308	0.308	0.346	0.192	0.192	0.192	0.192	0.192	0.000	
21	MH778702 <i>Hebius chapaensis</i>	0.192	0.192	0.269	0.269	0.192	0.192	0.269	0.308	0.308	0.308	0.346	0.192	0.192	0.192	0.192	0.115	0.000	
22	MN788503 <i>Hebius khasiensis</i>	0.231	0.231	0.308	0.308	0.231	0.231	0.308	0.346	0.346	0.346	0.385	0.231	0.231	0.231	0.231	0.231	0.154	
23	KJ685668 <i>Hebius khasiensis</i>	0.231	0.269	0.346	0.346	0.269	0.269	0.346	0.308	0.308	0.308	0.346	0.269	0.269	0.269	0.269	0.192	0.077	
24	KJ685683 <i>Hebius cf. ruthveni</i>	0.231	0.231	0.308	0.308	0.231	0.231	0.269	0.346	0.346	0.346	0.385	0.192	0.192	0.192	0.192	0.154	0.038	
25	OQ085073 <i>Hebius youjiangensis</i>	0.231	0.231	0.308	0.308	0.231	0.231	0.231	0.346	0.346	0.346	0.385	0.154	0.154	0.154	0.154	0.154	0.038	
26	KJ685711 <i>Hebius clerkii</i>	0.231	0.231	0.269	0.269	0.154	0.231	0.269	0.269	0.269	0.269	0.308	0.154	0.154	0.154	0.154	0.154	0.038	
27	AF402905 <i>Hebius maximus</i>	0.231	0.231	0.308	0.308	0.231	0.231	0.269	0.346	0.346	0.346	0.385	0.192	0.192	0.192	0.192	0.154	0.038	

Appendix 4 (continued). Uncorrected p-distance estimated using Cytb gene sequence among the genera *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, *Amphiesma* Duméril, Bibron & Duméril, 1854, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023, with *Amphiesmoides* Malnate, 1961 as outgroup.

No.	Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
28	KJ685687 <i>Hebius optatum</i>	0.231	0.269	0.346	0.269	0.269	0.308	0.308	0.308	0.346	0.231	0.231	0.231	0.192	0.077	0.077			
29	MZ570490 <i>Hebius weixiensis</i>	0.231	0.269	0.346	0.269	0.269	0.308	0.308	0.308	0.346	0.231	0.231	0.231	0.231	0.115	0.115			
30	OK315813 <i>Hebius atemporalis</i>	0.231	0.269	0.346	0.269	0.269	0.308	0.308	0.308	0.346	0.231	0.231	0.231	0.192	0.077	0.077			
31	KJ685673 <i>Hebius johannis</i>	0.231	0.231	0.308	0.231	0.231	0.269	0.346	0.346	0.385	0.192	0.192	0.192	0.192	0.154	0.038	0.038		
32	OK315824 <i>Hebius terrakarenorum</i>	0.231	0.192	0.269	0.231	0.231	0.308	0.346	0.346	0.385	0.231	0.231	0.231	0.231	0.154	0.038	0.038		
33	MZ570485 <i>Hebius septentrionalis</i>	0.231	0.269	0.346	0.269	0.269	0.308	0.308	0.308	0.346	0.231	0.231	0.231	0.231	0.115	0.115			
34	AB989302 <i>Hebius vibakari</i>	0.269	0.231	0.308	0.269	0.269	0.308	0.385	0.385	0.423	0.192	0.192	0.192	0.192	0.192	0.077	0.077		
35	KJ685678 <i>Hebius octolineatus</i>	0.269	0.269	0.346	0.269	0.269	0.192	0.308	0.308	0.346	0.231	0.231	0.231	0.231	0.115	0.077	0.077		
36	KJ685682 <i>Hebius metusia</i>	0.269	0.231	0.308	0.269	0.269	0.308	0.385	0.385	0.423	0.192	0.192	0.192	0.192	0.192	0.077	0.077		
37	KJ685688 <i>Hebius</i> sp.	0.269	0.231	0.308	0.269	0.269	0.308	0.385	0.385	0.423	0.192	0.192	0.192	0.192	0.192	0.077	0.077		
38	MH532291 <i>Hebius yanbianensis</i>	0.269	0.269	0.346	0.269	0.269	0.308	0.385	0.385	0.423	0.231	0.231	0.231	0.231	0.192	0.077	0.077		
39	OK315816 <i>Hebius bitaeniatus</i>	0.269	0.308	0.385	0.308	0.308	0.308	0.269	0.269	0.308	0.269	0.269	0.269	0.269	0.154	0.154			
40	KJ685689 <i>Hebius crassodogaster</i>	0.269	0.192	0.346	0.346	0.346	0.346	0.346	0.346	0.385	0.308	0.308	0.308	0.308	0.308	0.192	0.192		
41	MK340763 <i>Hebius sangzhiensis</i>	0.269	0.231	0.308	0.269	0.269	0.308	0.385	0.385	0.423	0.231	0.231	0.231	0.231	0.115	0.115			
42	OK315825 <i>Hebius groundwateri</i>	0.269	0.231	0.308	0.269	0.269	0.269	0.385	0.385	0.423	0.192	0.192	0.192	0.192	0.192	0.077	0.077		
43	KJ685670 <i>Hebius vennigi</i>	0.269	0.269	0.269	0.231	0.231	0.308	0.308	0.308	0.346	0.231	0.231	0.231	0.231	0.115	0.115			
44	OR389974 <i>Hebius vennigi</i>	0.308	0.308	0.269	0.269	0.192	0.269	0.269	0.269	0.308	0.192	0.192	0.192	0.192	0.269	0.154	0.154		
45	OR389975 <i>Hebius vennigi</i>	0.308	0.308	0.269	0.269	0.192	0.269	0.269	0.269	0.308	0.192	0.192	0.192	0.192	0.269	0.154	0.154		
46	KJ685701 <i>Hebius sauteri</i>	0.308	0.269	0.423	0.423	0.346	0.269	0.308	0.269	0.269	0.308	0.308	0.308	0.308	0.192	0.154	0.154		
47	AB989296 <i>Hebius ishigakiensis</i>	0.308	0.308	0.308	0.269	0.269	0.346	0.346	0.346	0.385	0.269	0.269	0.269	0.269	0.115	0.115			
48	KJ685692 <i>Hebius popei</i>	0.308	0.308	0.346	0.231	0.231	0.269	0.346	0.346	0.385	0.192	0.192	0.192	0.192	0.115	0.115			
49	KJ685679 <i>Hebius taronensis</i>	0.308	0.308	0.269	0.269	0.192	0.269	0.269	0.269	0.308	0.192	0.192	0.192	0.192	0.154	0.154			

Appendix 4 (continued). Uncorrected p-distance estimated using Cytb gene sequence among the genera *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, *Amphiesma* Duméril, Bibron & Duméril, 1854, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023, with *Amphiesmoides* Malnate, 1961 as outgroup.

No.	Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
50	OK315829 <i>Hebius nigrovirens</i>	0.308	0.308	0.269	0.192	0.269	0.269	0.269	0.269	0.269	0.308	0.192	0.192	0.192	0.269	0.154	0.154	0.154	
51	AB989268 <i>Hebius concularius</i>	0.346	0.269	0.308	0.269	0.269	0.308	0.385	0.385	0.385	0.423	0.231	0.231	0.231	0.231	0.308	0.192	0.192	
52	OK315821 <i>Hebius leucomystax</i>	0.346	0.385	0.462	0.462	0.385	0.308	0.423	0.269	0.269	0.269	0.308	0.385	0.385	0.385	0.385	0.269	0.231	0.231
53	AB989102 <i>Hebius pryeri</i>	0.385	0.385	0.385	0.269	0.269	0.308	0.308	0.308	0.308	0.346	0.269	0.269	0.269	0.269	0.346	0.231	0.231	
54	KJ685694 <i>Amphiesma stolatum</i>	0.269	0.192	0.308	0.308	0.231	0.231	0.269	0.308	0.308	0.346	0.231	0.231	0.231	0.231	0.308	0.192	0.192	
55	OK315803 <i>Amphiesma monticola</i>	0.269	0.231	0.231	0.231	0.269	0.308	0.423	0.346	0.346	0.385	0.269	0.269	0.269	0.269	0.346	0.231	0.231	
56	OK315814 <i>Sahyadriophis beddomei</i>	0.385	0.346	0.346	0.346	0.346	0.346	0.269	0.346	0.346	0.346	0.308	0.385	0.385	0.385	0.308	0.308	0.308	
57	OK326949 <i>Amphiesmoides ornatusiceps</i>	0.462	0.538	0.500	0.500	0.385	0.462	0.500	0.577	0.577	0.538	0.423	0.423	0.423	0.423	0.385	0.385	0.385	

Appendix 4 (continued). Uncorrected p-distance estimated using Cytb gene sequence among the genera *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, *Amphiesma* Duméril, Bibron & Duméril, 1854, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023, with *Amphiesmoides* Malnate, 1961 as outgroup.

No.	Species	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
1	OK315830 <i>Herpetoreas davidi</i> sp. nov																		
2	ON204025 <i>Herpetoreas murlen</i>																		
3	MZ358098 <i>Herpetoreas burbrinki</i>																		
4	GQ281781 <i>Herpetoreas burbrinki</i>																		
5	MW111464 <i>Herpetoreas platiceps</i>																		
6	MT571587 <i>Herpetoreas cf. sieboldii</i>																		
7	MT571586 <i>Herpetoreas pealii</i>																		
8	MN993850 <i>Herpetoreas xenura</i>																		
9	MN993852 <i>Herpetoreas xenura</i>																		
10	MN993851 <i>Herpetoreas xenura</i>																		
11	MN993891 <i>Herpetoreas xenura</i>																		
12	MW111476 <i>Herpetoreas ipser</i>																		
13	OM313293 <i>Herpetoreas ipser</i>																		
14	OM313292 <i>Herpetoreas ipser</i>																		
15	MK201567 <i>Herpetoreas ipser</i>																		
16	MK253674 <i>Hebius andreae</i>																		
17	OK315812 <i>Hebius annamensis</i>																		
18	KJ685671 <i>Hebius modestum</i>																		
19	KJ685684 <i>Hebius bouengeri</i>																		
20	KJ685700 <i>Hebius deschauenseei</i>																		
21	MH778702 <i>Hebius chapaensis</i>																		
22	MN788503 <i>Hebius khasiensis</i>																		
23	KJ685668 <i>Hebius khasiensis</i>																		
24	KJ685683 <i>Hebius cf. trutveni</i>																		
25	OQ085073 <i>Hebius youtiangensis</i>																		

Appendix 4 (continued). Uncorrected p-distance estimated using Cytb gene sequence among the genera *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, *Amphiesma* Duméril, Bibron & Duméril, 1854, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023, with *Amphiesmoides* Malnate, 1961 as outgroup.

No.	Species	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
26	KJ685711 <i>Hebius clerki</i>	0.154	0.038	0.038	0.077	0.115	0.077	0.077	0.115	0.000	0.038	0.077	0.077	0.115	0.154	0.154	0.154	0.154	0.154
27	AF402905 <i>Hebius maximus</i>	0.154	0.038	0.038	0.077	0.115	0.077	0.038	0.077	0.115	0.038	0.077	0.077	0.115	0.154	0.154	0.154	0.154	0.154
28	KJ685687 <i>Hebius opiatum</i>	0.115	0.077	0.077	0.115	0.154	0.115	0.115	0.115	0.154	0.115	0.077	0.077	0.115	0.154	0.154	0.154	0.154	0.154
29	MZ570490 <i>Hebius weixiensis</i>	0.115	0.115	0.115	0.154	0.115	0.115	0.115	0.115	0.154	0.115	0.077	0.077	0.115	0.154	0.154	0.154	0.154	0.154
30	OK315813 <i>Hebius atemporalis</i>	0.115	0.077	0.077	0.115	0.077	0.115	0.000	0.038	0.077	0.000	0.038	0.077	0.115	0.154	0.154	0.154	0.154	0.154
31	KJ685673 <i>Hebius johannis</i>	0.154	0.038	0.038	0.077	0.115	0.077	0.077	0.077	0.077	0.077	0.115	0.115	0.115	0.154	0.154	0.154	0.154	0.154
32	OK315824 <i>Hebius terrakarenorum</i>	0.154	0.038	0.038	0.077	0.115	0.077	0.115	0.154	0.077	0.038	0.038	0.077	0.115	0.154	0.154	0.154	0.154	0.154
33	MZ570485 <i>Hebius septentrionalis</i>	0.115	0.115	0.115	0.154	0.115	0.077	0.115	0.154	0.077	0.038	0.038	0.077	0.077	0.115	0.154	0.154	0.154	0.154
34	AB899302 <i>Hebius vibakari</i>	0.192	0.077	0.077	0.115	0.154	0.038	0.077	0.115	0.038	0.077	0.154	0.115	0.038	0.115	0.115	0.115	0.115	0.115
35	KJ685678 <i>Hebius octolineatus</i>	0.192	0.077	0.077	0.115	0.154	0.038	0.077	0.115	0.038	0.077	0.154	0.115	0.038	0.115	0.115	0.115	0.115	0.115
36	KJ685682 <i>Hebius menisia</i>	0.192	0.077	0.077	0.115	0.154	0.038	0.077	0.115	0.038	0.077	0.154	0.115	0.038	0.115	0.115	0.115	0.115	0.115
37	KJ685688 <i>Hebius</i> sp.	0.192	0.077	0.077	0.115	0.154	0.038	0.077	0.115	0.038	0.077	0.154	0.115	0.038	0.115	0.115	0.115	0.115	0.115
38	MH532291 <i>Hebius yanbianensis</i>	0.115	0.077	0.077	0.038	0.077	0.115	0.038	0.077	0.115	0.038	0.077	0.154	0.115	0.038	0.115	0.115	0.115	0.115
39	OK315816 <i>Hebius bitaenatus</i>	0.154	0.154	0.154	0.192	0.154	0.115	0.154	0.192	0.115	0.077	0.077	0.115	0.154	0.154	0.154	0.154	0.154	0.154
40	KJ685689 <i>Hebius craspedogaster</i>	0.192	0.192	0.192	0.231	0.192	0.154	0.192	0.231	0.154	0.115	0.115	0.154	0.154	0.154	0.154	0.154	0.154	0.154
41	MK340763 <i>Hebius sangzhensis</i>	0.192	0.115	0.115	0.154	0.192	0.077	0.115	0.154	0.077	0.115	0.115	0.154	0.077	0.154	0.154	0.154	0.154	0.154
42	OK315825 <i>Hebius groundwateri</i>	0.192	0.077	0.077	0.115	0.154	0.077	0.038	0.115	0.077	0.115	0.154	0.115	0.077	0.154	0.154	0.154	0.154	0.154
43	KJ685670 <i>Hebius vennigi</i>	0.192	0.115	0.115	0.154	0.192	0.077	0.115	0.154	0.077	0.115	0.154	0.115	0.077	0.154	0.154	0.154	0.154	0.154
44	OR389974 <i>Hebius vennigi</i>	0.231	0.154	0.154	0.192	0.231	0.115	0.154	0.115	0.115	0.154	0.154	0.115	0.154	0.192	0.115	0.154	0.154	0.154
45	OR389975 <i>Hebius vennigi</i>	0.231	0.154	0.154	0.192	0.231	0.115	0.154	0.115	0.154	0.154	0.154	0.115	0.192	0.115	0.154	0.154	0.154	0.154
46	KJ685701 <i>Hebius sauteri</i>	0.269	0.154	0.154	0.192	0.231	0.115	0.154	0.192	0.115	0.154	0.231	0.192	0.115	0.192	0.192	0.192	0.192	0.192
47	AB899296 <i>Hebius ishgakiensis</i>	0.154	0.115	0.115	0.077	0.115	0.077	0.115	0.154	0.077	0.115	0.154	0.192	0.154	0.077	0.154	0.154	0.154	0.154
48	KJ685692 <i>Hebius popei</i>	0.154	0.115	0.115	0.077	0.115	0.077	0.115	0.077	0.077	0.115	0.192	0.154	0.077	0.154	0.154	0.154	0.154	0.154
49	KJ685679 <i>Hebius taronensis</i>	0.231	0.154	0.154	0.192	0.231	0.115	0.154	0.115	0.154	0.154	0.154	0.192	0.115	0.154	0.154	0.154	0.154	0.154
50	OK315829 <i>Hebius nigritiver</i>	0.231	0.154	0.154	0.192	0.231	0.115	0.154	0.115	0.154	0.154	0.154	0.192	0.115	0.154	0.154	0.154	0.154	0.154
51	AB899268 <i>Hebius concularius</i>	0.192	0.192	0.192	0.154	0.192	0.154	0.192	0.154	0.192	0.192	0.231	0.154	0.192	0.192	0.154	0.192	0.192	0.192

Appendix 4 (continued). Uncorrected p-distance estimated using Cytb gene sequence among the genera *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, *Amphiesma* Duméril, Bibron & Duméril, 1854, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023, with *Amphiesmoides* Malnate, 1961 as outgroup.

No.	Species	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
52	OK315821 <i>Hebius leucomystax</i>	0.231	0.231	0.231	0.269	0.231	0.269	0.269	0.269	0.231	0.192	0.231	0.269	0.192	0.231	0.269	0.192	0.231	0.308
53	AB889102 <i>Hebius pryeri</i>	0.231	0.231	0.231	0.192	0.231	0.192	0.231	0.192	0.192	0.231	0.231	0.269	0.192	0.269	0.192	0.231	0.231	0.231
54	KJ685694 <i>Amphiesma stolatum</i>	0.154	0.192	0.192	0.154	0.192	0.231	0.231	0.154	0.231	0.154	0.269	0.269	0.231	0.192	0.269	0.269	0.269	0.269
55	OK315803 <i>Amphiesma monticola</i>	0.231	0.231	0.231	0.269	0.269	0.269	0.269	0.269	0.269	0.269	0.269	0.269	0.269	0.231	0.269	0.269	0.231	0.231
56	OK315814 <i>Sahyadriophis beddomei</i>	0.346	0.308	0.308	0.346	0.385	0.346	0.346	0.346	0.346	0.346	0.346	0.346	0.346	0.385	0.385	0.346	0.385	0.308
57	OK326949 <i>Amphiesmoides ornatusiceps</i>	0.500	0.385	0.385	0.423	0.462	0.423	0.423	0.346	0.423	0.423	0.500	0.462	0.423	0.423	0.500	0.462	0.462	0.462

Appendix 4 (continued). Uncorrected p-distance estimated using Cytb gene sequence among the genera *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, *Amphiesma* Duméril, Bibron & Duméril, 1854, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023, with *Amphiesmoides* Malnate, 1961 as outgroup.

No.	Species	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56
1	OK315830 <i>Herpetoreas davidi</i> sp. nov																				
2	ON204025 <i>Herpetoreas murlen</i>																				
3	MZ358098 <i>Herpetoreas burbrinki</i>																				
4	GQ281781 <i>Herpetoreas burbrinki</i>																				
5	MW111464 <i>Herpetoreas platiceps</i>																				
6	MT571587 <i>Herpetoreas cf. sieboldii</i>																				
7	MT571586 <i>Herpetoreas pealii</i>																				
8	MN993850 <i>Herpetoreas xenura</i>																				
9	MN993852 <i>Herpetoreas xenura</i>																				
10	MN993851 <i>Herpetoreas xenura</i>																				
11	MN993891 <i>Herpetoreas xenura</i>																				
12	MW111476 <i>Herpetoreas tpsr</i>																				
13	OM313293 <i>Herpetoreas tpsr</i>																				
14	OM313292 <i>Herpetoreas tpsr</i>																				
15	MK201567 <i>Herpetoreas tpsr</i>																				
16	MK253674 <i>Hebius andreae</i>																				
17	OK315812 <i>Hebius annamensis</i>																				
18	KJ685671 <i>Hebius modestum</i>																				
19	KJ685684 <i>Hebius bouengeri</i>																				
20	KJ685700 <i>Hebius deschauenseei</i>																				
21	MH778702 <i>Hebius chapaensis</i>																				
22	MN788503 <i>Hebius khasiensis</i>																				
23	KJ685668 <i>Hebius khasiensis</i>																				
24	KJ685683 <i>Hebius cf. ruthveni</i>																				
25	OQ083073 <i>Hebius yujiangensis</i>																				
26	KJ685711 <i>Hebius clerkii</i>																				
27	AF402905 <i>Hebius maximus</i>																				
28	KJ685687 <i>Hebius optatum</i>																				
29	MZ570490 <i>Hebius weixiensis</i>																				

Appendix 4 (continued). Uncorrected p-distance estimated using Cytb gene sequence among the genera *Hebius* Thompson, 1913, *Herpetoreas* Günther, 1860, *Amphiesma* Duméril, Bibron & Duméril, 1854, and *Sahyadriophis* Patel, Thackeray, Campbell & Mirza, 2023, with *Amphiesmoides* Malnate, 1961 as outgroup.

No.	Species	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56
30	OK315813 <i>Hebius aequalis</i>																				
31	KJ685673 <i>Hebius johannis</i>																				
32	OK315824 <i>Hebius terrakarenorum</i>																				
33	MZ570485 <i>Hebius septentrionalis</i>																				
34	AB989302 <i>Hebius vibakari</i>																				
35	KJ685678 <i>Hebius octolineatus</i>																				
36	KJ685682 <i>Hebius metusia</i>																				
37	KJ685688 <i>Hebius</i> sp.																				
38	MH532291 <i>Hebius yanbianensis</i>	0.077																			
39	OK315816 <i>Hebius bitaeniatus</i>	0.154	0.154																		
40	KJ685689 <i>Hebius craspedogaster</i>	0.192	0.192	0.115																	
41	MK340763 <i>Hebius sangzhiensis</i>	0.038	0.115	0.115	0.154																
42	OK315825 <i>Hebius groundwaei</i>	0.115	0.115	0.192	0.192	0.154															
43	KJ685670 <i>Hebius craspedogaster</i>	0.115	0.115	0.115	0.154	0.077	0.154														
44	OR389974 <i>Hebius sangzhiensis</i>	0.154	0.154	0.154	0.192	0.115	0.115	0.192	0.038												
45	OR389975 <i>Hebius vennigi</i>	0.154	0.154	0.154	0.192	0.115	0.192	0.192	0.038	0.000											
46	KJ685701 <i>Hebius vennigi</i>	0.154	0.154	0.154	0.192	0.192	0.192	0.192	0.192	0.231	0.231										
47	AB989296 <i>Hebius ishgakiensis</i>	0.115	0.038	0.192	0.231	0.154	0.154	0.154	0.077	0.115	0.115	0.192									
48	KJ685692 <i>Hebius popei</i>	0.115	0.038	0.192	0.231	0.154	0.154	0.154	0.154	0.115	0.115	0.192	0.077								
49	KJ685679 <i>Hebius taronensis</i>	0.154	0.154	0.154	0.192	0.115	0.192	0.038	0.000	0.000	0.000	0.231	0.115	0.115							
50	OK315829 <i>Hebius nigrovirens</i>	0.154	0.154	0.154	0.192	0.115	0.192	0.038	0.000	0.000	0.000	0.231	0.115	0.115	0.000						
51	AB989268 <i>Hebius concularius</i>	0.192	0.115	0.192	0.154	0.154	0.192	0.154	0.115	0.115	0.269	0.154	0.077	0.115	0.115						
52	OK315821 <i>Hebius leucomystax</i>	0.308	0.308	0.154	0.269	0.269	0.308	0.269	0.308	0.308	0.231	0.346	0.346	0.308	0.308	0.346					
53	AB989102 <i>Hebius pryeri</i>	0.231	0.154	0.154	0.269	0.192	0.269	0.154	0.115	0.115	0.231	0.154	0.115	0.115	0.115	0.115	0.115	0.115	0.115	0.308	
54	KJ685694 <i>Amphiesma stolatum</i>	0.269	0.192	0.308	0.231	0.269	0.192	0.269	0.231	0.231	0.269	0.231	0.154	0.231	0.231	0.154	0.385	0.231			
55	OK315803 <i>Amphiesma monticola</i>	0.231	0.308	0.308	0.269	0.231	0.269	0.231	0.269	0.385	0.269	0.346	0.269	0.346	0.308	0.308	0.385	0.385	0.308		
56	OK315814 <i>Sahyadriophis beddomii</i>	0.308	0.385	0.423	0.462	0.308	0.385	0.308	0.346	0.346	0.385	0.423	0.346	0.462	0.462	0.462	0.462	0.462	0.462	0.423	
57	OK320949 <i>Amphiesmoides ornaticeps</i>	0.462	0.462	0.538	0.577	0.500	0.462	0.500	0.462	0.538	0.500	0.423	0.462	0.462	0.500	0.577	0.538	0.500	0.577	0.538	

Appendix 5 (continued on next four pages). Literature used for the revised distribution of *Herpetoreas davidi* sp. nov., *H. xenura* (Wall, 1907), and *Hebius khasiensis* (Boulenger, 1890).

No.	Species	Number in map	Verified by molecular	Verified by morphology	Location	Sources
1	<i>Herpetoreas davidi</i> sp. nov.	1	yes	yes	Daung Stream, Rakhine Yoma Elephant Sanctuary, Rakhine, Myanmar (type locality)	our data
2	<i>Herpetoreas davidi</i> sp. nov.	2	no	yes	Kyat Stream, Rakhine Yoma Elephant Sanctuary, Rakhine, Myanmar	our data
3	<i>Herpetoreas xenura</i>	1	no	yes	Cherrapunji, East Khasi Hills, Meghalaya, India (type locality)	Wall 1907; Ren et al. 2022
4	<i>Herpetoreas xenura</i>	2	no	yes	Tura, West Garo Hills, Meghalaya, India	Mathew 1995; Ren et al. 2022
5	<i>Herpetoreas xenura</i>	3	no	yes	Umroi, Meghalaya, India	Muansanga et al. 2021
6	<i>Herpetoreas xenura</i>	4	no	yes	Narpuh RF, Jaintia Hills, Meghalaya, India	Mathew & Meetei 2004; Ren et al. 2022
7	<i>Herpetoreas xenura</i>	5	yes	no	Menam Mikir Pumjee, Assam, India	Lalronunga et al. 2020
8	<i>Herpetoreas xenura</i>	6	no	yes	Kohima, Nagaland, India	Romer 1945; NHMUK 1956.1.12.79
9	<i>Herpetoreas xenura</i>	7	no	yes	Naga Hills, Nagaland, India	Romer 1945; Ren et al. 2022
10	<i>Herpetoreas xenura</i>	8	no	yes	North Hlumen, Kolasib Dist., Mizoram, India	Muansanga et al. 2021
11	<i>Herpetoreas xenura</i>	9	no	yes	Silphur, Aizawl Dist., Mizoram, India	Muansanga et al. 2021
12	<i>Herpetoreas xenura</i>	10	no	yes	Reiek Community Reserved Forest, Manit, Mizoram, India	Hmar et al. 2020
13	<i>Herpetoreas xenura</i>	11	no	yes	Dampa Tiger Reserve, Manit Dist., Mizoram, India	Malsawndawnglana et al. 2022
14	<i>Herpetoreas xenura</i>	12	no	yes	Pala Lake Area, Mizoram, India	Pawar & Birand 2001; Ren et al. 2022
15	<i>Herpetoreas xenura</i>	13	no	yes	Tripura, India	Samuel Laltronunga unpubl. data
16	<i>Herpetoreas xenura</i>	14	no	yes	Kaptai NP, Rangamati, Bangladesh	Reza 2010; Ren et al. 2022
17	<i>Herpetoreas xenura</i>	15	no	yes	Bandarban, Keokradong, Bangladesh	Ahsan et al. 2009; Ren et al. 2022

Appendix 5 (continued). Literature used for the revised distribution of *Herpetoreas davidi* sp. nov., *H. xenura* (Wall, 1907), and *Hebius khasiensis* (Boulenger, 1890).

No.	Species	Number in map	Verified by molecular	Verified by morphology	Location	Sources
18	<i>Herpetoreas xenura</i>	16	no	yes	Bilkhawhlir, Kolasib Dist., Mizoram, India	our data
19	<i>Herpetoreas xenura</i>	17	no	yes	Sawleng, Aizawl Dist., Mizoram, India	our data
20	<i>Herpetoreas xenura</i>	18	no	yes	Kawnpui, Kolasib Dist., Mizoram, India	our data
21	<i>Herpetoreas xenura</i>	19	no	yes	Dampa Tiger Reserve, Mamit Dist., Mizoram, India	our data
22	<i>Herpetoreas xenura</i>	20	no	yes	Tamdi national Wetland, Saitual Dist., Mizoram, India	our data
23	<i>Herpetoreas xenura</i>	21	no	yes	Saitual, Saitual Dist., Mizoram, India	our data
24	<i>Herpetoreas xenura</i>	22	no	yes	Hmuifang Community Reserved Forest, Aizawl Dist., Mizoram, India	our data
25	<i>Herpetoreas xenura</i>	23	no	yes	Thenzawl, Serchhip Dist., Mizoram, India	our data
26	<i>Herpetoreas xenura</i>	24	no	yes	Lunglei, Lunglei Dist., Mizoram, India	our data
27	<i>Herpetoreas xenura</i>	Not shown	no	yes	Mizoram University Campus, Aizawl Dist., Mizoram, India	our data
28	<i>Herpetoreas xenura</i>	Not shown	no	yes	Luangmual, Aizawl Dist., Mizoram, India	our data
29	<i>Herpetoreas xenura</i>	Not shown	no	yes	Durtlang, Aizawl, Dist., Mizoram, India	our data
30	<i>Herpetoreas xenura</i>	Not shown	no	yes	Chandmari West, Aizawl Dist., Mizoram, India	our data
31	<i>Herpetoreas xenura</i>	Not shown	no	yes	Ramrikawn, Aizawl Dist., Mizoram, India	our data
32	<i>Herpetoreas xenura</i>	Not shown	no	yes	Tanhril, Aizawl Dist., Mizoram, India	our data
33	<i>Herpetoreas xenura</i>	Not shown	no	yes	Sihhmui, Aizawl Dist., Mizoram, India	our data

Appendix 5 (continued). Literature used for the revised distribution of *Herpetoreas davidi* sp. nov., *H. xenura* (Wall, 1907), and *Hebius khasiensis* (Boulenger, 1890).

No.	Species	Number in map	Verified by molecular	Verified by morphology	Location	Sources
34	<i>Herpetoreas xenura</i>	Not shown	no	yes	Mission vengthlang, Aizawl Dist., Mizoram, India	our data
35	<i>Herpetoreas xenura</i>	Not shown	no	yes	Tuirial, Aizawl Dist., Mizoram, India	our data
36	<i>Herpetoreas xenura</i>	Not shown	no	yes	Seling, Aizawl Dist., Mizoram, India	our data
37	<i>Herpetoreas xenura</i>	Not shown	no	yes	College veng, Aizawl Dist., Mizoram, India	our data
38	<i>Hebius khasiensis</i>	1	no	yes	Khasi Hills, Meghalaya, India (type locality)	David <i>et al.</i> 2013; our data
39	<i>Hebius khasiensis</i>	2	no	yes	Dibang, Arunachal Pradesh, India	David <i>et al.</i> 2013; our data
40	<i>Hebius khasiensis</i>	3	no	yes	Namdapha Tiger Reserve, Changlang, Arunachal Pradesh	https://www.inaturalist.org/observations/29628011
41	<i>Hebius khasiensis</i>	4	no	yes	Changlang, Arunachal Pradesh, India	David <i>et al.</i> 2013; our data
42	<i>Hebius khasiensis</i>	5	no	yes	Zunheboto, Nagaland, India	David <i>et al.</i> 2013; our data
43	<i>Hebius khasiensis</i>	6	no	yes	Kohima, Nagaland, India	David <i>et al.</i> 2013; our data
44	<i>Hebius khasiensis</i>	7	no	yes	Reiek Community RF, Mizoram, India	Hmar <i>et al.</i> 2020
45	<i>Hebius khasiensis</i>	8	no	yes	Dampa Tiger Reserve, Mizoram, India	Malsawmdawnglana <i>et al.</i> 2022
46	<i>Hebius khasiensis</i>	9	yes	yes	Putao, Kachin, Myanmar	David <i>et al.</i> 2013; our data
47	<i>Hebius khasiensis</i>	10	no	yes	Hutung, Bhamo, Kachin, Myanmar (type locality of <i>Natrix gilhoensis</i>)	David <i>et al.</i> 2013; our data
48	<i>Hebius khasiensis</i>	11	no	yes	Bhamo, Kachin, Myanmar	https://www.inaturalist.org/observations/119392371
49	<i>Hebius khasiensis</i>	12	no	yes	Nat Ma Taung, Chin, Myanmar	David <i>et al.</i> 2013; our data
50	<i>Hebius khasiensis</i>	13	no	yes	Mindat, Chin, Myanmar	https://www.inaturalist.org/observations/144728979
51	<i>Hebius khasiensis</i>	14	no	yes	Karen Hills, Shan, Myanmar	David <i>et al.</i> 2013; our data

Appendix 5 (continued). Literature used for the revised distribution of *Herpetoreas davidi* sp. nov., *H. xenura* (Wall, 1907), and *Hebius khasiensis* (Boulenger, 1890).

No.	Species	Number in map	Verified by molecular	Verified by morphology	Location	Sources
52	<i>Hebius khasiensis</i>	15	no	yes	Medog, Nyingchi, Tibet, China	David <i>et al.</i> 2013; Che <i>et al.</i> 2020
53	<i>Hebius khasiensis</i>	16	no	yes	Htamanthi WS, Sagaing, Myanmar	our data
54	<i>Hebius khasiensis</i>	17	yes	yes	Yingjiang, Yunnan, China	Yang & Rao 2008; Wu <i>et al.</i> 2023
55	<i>Hebius khasiensis</i>	18	no	yes	Ruili, Dehong, Yunnan, China	David <i>et al.</i> 2013; our data
56	<i>Hebius khasiensis</i>	19	no	yes	Longling, Baoshan, Yunnan, China	Yang & Rao 2008
57	<i>Hebius khasiensis</i>	20	no	yes	Menglian, Puer, Yunnan, China	Yang & Rao 2008
58	<i>Hebius khasiensis</i>	21	no	yes	Xichuangbanna, Yunnan, China	David <i>et al.</i> 2013; our data
59	<i>Hebius khasiensis</i>	22	no	yes	Boun Tay, Phongsaly, Laos	David <i>et al.</i> 2013; our data
60	<i>Hebius khasiensis</i>	23	no	yes	Doi Tung Mt., Chiang Rai, Thailand	https://www.inaturalist.org/observations/12865317
61	<i>Hebius khasiensis</i>	24	no	yes	Doi Luang NP, Chiang Rai, Thailand	https://www.inaturalist.org/observations/86044420
62	<i>Hebius khasiensis</i>	25	no	yes	Chae Son NP, Lampang, Thailand	Chan-ard <i>et al.</i> 2015
63	<i>Hebius khasiensis</i>	26	no	yes	Doi Phu Kha NP, Nan, Thailand	https://www.inaturalist.org/observations/165474307
64	<i>Hebius khasiensis</i>	27	no	yes	Doi Suthep-Pui NP, Chiang Mai, Thailand	Chan-ard <i>et al.</i> 2015
65	<i>Hebius khasiensis</i>	28	no	yes	Doi Inthanon NP, Chiang Mai, Thailand	our data; https://www.inaturalist.org/observations/123678122
66	<i>Hebius khasiensis</i>	29	no	yes	Mae Tho NP, Chiang Mai, Thailand	Chan-ard <i>et al.</i> 2015; our data
67	<i>Hebius khasiensis</i>	30	no	yes	Huai Nam Dang NP, Mae Hong Son, Thailand	Chan-ard <i>et al.</i> 2015; our data
68	<i>Hebius khasiensis</i>	31	no	yes	Lan Sang NP, Tak, Thailand	Chan-ard <i>et al.</i> 2015; our data
69	<i>Hebius khasiensis</i>	32	no	yes	Namtok Pha Charoen NP, Tak, Thailand	Chan-ard <i>et al.</i> 2015; our data
70	<i>Hebius khasiensis</i>	33	no	yes	Khlong Wang Chao NP, Kamphaeng Phet, Thailand	Chan-ard <i>et al.</i> 2015; our data

Appendix 5 (continued). Literature used for the revised distribution of *Herpetoreas davidi* sp. nov., *H. xenura* (Wall, 1907), and *Hebius khasiensis* (Boulenger, 1890).

No.	Species	Number in map	Verified by molecular	Verified by morphology	Location	Sources
71	<i>Hebius khasiensis</i>	34	no	yes	Umphang WS, Tak, Thailand	David <i>et al.</i> 2013; our data https://www.inaturalist.org/observations/53232442
72	<i>Hebius khasiensis</i>	35	no	yes	Phu Suan Sai NP, Loei, Thailand	our data, https://www.inaturalist.org/observations/105954833 https://www.inaturalist.org/observations/53232442
73	<i>Hebius khasiensis</i>	36	no	yes	Phu Luang WS, Loei, Thailand	our data, https://www.inaturalist.org/observations/105954833 https://www.inaturalist.org/observations/53232442
74	<i>Hebius khasiensis</i>	37	no	yes	Phu Suan Sai NP, Loei, Thailand	our data, https://www.inaturalist.org/observations/105954833 https://www.inaturalist.org/observations/53232442
75	<i>Hebius khasiensis</i>	38	no	yes	Nam Nao NP, Phetchabun, Thailand	https://www.inaturalist.org/observations/37859050