



This work is licensed under a Creative Commons Attribution License (CC BY 4.0).

Corrigendum

urn:lsid:zoobank.org:pub:3ACB94F7-F020-4038-BD00-EE79061AFC17

A revision of the Afrotropical hover fly genus *Afroxanthandrus* Kassebeer, 2000 (Diptera, Syrphidae), with the description of two new species and one new synonymy – Corrigendum

John M. MIDGLEY¹, Georg GOERGEN² & Kurt JORDAENS³

¹KwaZulu-Natal Museum, Department of Natural Sciences, 237 Jabu Ndlovu Street, Pietermaritzburg, 3201, South Africa.

¹Rhodes University, Department of Zoology and Entomology, P.O. Box 94, Makhanda, 6140, South Africa.

²International Institute of Tropical Agriculture, Biodiversity Centre, 08 BP 0932 Tri Postal, Cotonou, Benin.

³Royal Museum for Central Africa, Department of Biology and JEMU, Leuvensesteenweg 13, B-3080 Tervuren, Belgium.

*Corresponding author: jmidgley@nmsa.org.za

²Email: G.Goergen@cgiar.org

³Email: kurt.jordaens@africamuseum.be

¹urn:lsid:zoobank.org:author:FBAB428E-8A94-42AF-A489-F5131CF05A87

²urn:lsid:zoobank.org:author:11B63B43-550B-46BE-B9BF-BC1F0A803FAA

³urn:lsid:zoobank.org:author:9E6F237E-BC90-4F8F-891F-798347E9A3F9

Midgley J.M., Goergen G. & Jordaens K. 2024. A revision of the Afrotropical hover fly genus *Afroxanthandrus* Kassebeer, 2000 (Diptera, Syrphidae), with the description of two new species and one new synonymy – Corrigendum. *European Journal of Taxonomy* 968: 295–296. https://doi.org/10.5852/ejt.2024.968.2741

The present corrigendum corrects figure numbers that were incorrect in Midgley *et al.* (2024).

Page 63, the key should read:

Key to species

1. Tergite 4 entirely dark. Yellow fascia on tergite 2 with curved or triangular posterior border (Figs 14, 16–17) 2
- Tergite 4 with broad yellow fascia. Yellow fascia on tergite 2 with straight posterior border (Figs 13, 15) 3

2. Tergite 3 with broad fascia, about $\frac{3}{4}$ of the length of the tergite (Figs 14, 16); scutal pile short, half as long or less as length of scape (Figs 8, 10); male genitalia: surstyli short, about 2.5 times as long as wide (Figs 20–21) *A. congensis* (Curran, 1938)

- Tergite 3 with narrow fascia, about $\frac{1}{3}$ of the length of the tergite (Fig. 17); scutal pile long, as long as or slightly longer than length of scape (Fig. 11)
..... *A. longipilus* Kassebeer, 2000
- 3. Tergite 4 with broad fascia, similar in shape to fasciae on tergites 2 and 3 (Fig. 13); male genitalia: surstyli short, about twice as long as wide (Figs 18–19) *A. comorosensis* sp. nov.
- Tergite 4 with narrow fascia, clearly narrower than fasciae on tergites 2 and 3 (Fig. 15); male genitalia: surstyli long, about eight times as long as wide (Figs 22–23) *A. conopeum* sp. nov.

Page 74, in Remarks, 7th line: “...(Figs 19, 22)...” should read “...(Figs 20–21)...”.

Page 76, the figure legend should read: **Figs 18–23**. Male genitalia of species of *Afroxanthandrus* Kassebeer, 2000. **18–19**. *A. comorosensis* sp. nov., holotype (RMCA ENT 000056900). **18**. Dorsal view. **19**. Lateral view. **20–21**. *A. congensis* (Curran, 1938) (RMCA ENT 000028396). **20**. Dorsal view. **21**. Lateral view. **22–23**. *A. conopeum* sp. nov., paratype (NMSA-Dip 086809). **22**. Dorsal view. **23**. Lateral view. Scale bars = 0.5 mm.

Reference

Midgley J.M., Goergen G. & Jordaens K. 2024. A revision of the Afrotropical hover fly genus *Afroxanthandrus* Kassebeer, 2000 (Diptera, Syrphidae), with the description of two new species and one new synonymy. *European Journal of Taxonomy* 968: 60–85. <https://doi.org/10.5852/ejt.2024.968.2717>

Published on: 19 November 2024

Printed versions of all papers are deposited in the libraries of four of the institutes that are members of the EJT consortium: Muséum national d’Histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium. The other members of the consortium are: Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic; The Steinhardt Museum of Natural History, Tel Aviv, Israël.