

Received: 26 September 2024 • Accepted: 27 November 2025 • Published: 23 February 2026

Topic editor: Magalie Castellin • Section editor: Felipe Ottoni • Desk editor: Thomas Guyomard

Monograph

urn:lsid:zoobank.org:pub:C56D842B-1FEB-44AA-A788-3B0714DC7CE1

Freshwater fishes (Petromyzonti, Elasmobranchii, and Actinopteri) of Bosnia and Herzegovina: an updated inventory on the biodiversity and distribution

Pero TUTMAN¹  , Marko ČALETA^{2,*}  , Zoran MARČIĆ³  , Ivana BUJ⁴  ,
 Adem HAMZIĆ⁵  , Belma KALAMUJIĆ STROIL⁶  , Dragojla GOLUB⁷  ,
 Radek ŠANDA⁸  , Avdul ADROVIĆ⁹  , Pavle SPASOJEVIĆ¹⁰  , Jerko PAVLIČEVIĆ¹¹  ,
 Irena ROZIĆ¹²  , Nikola ZOVKO¹³  , Edhem HASKOVIĆ¹⁴  , Rifat ŠKRIJELJ¹⁵  ,
 Amra KAZIĆ¹⁶  , Radoslav DEKIĆ¹⁷  , Jakov DULČIĆ¹⁸  , Perica MUSTAFIĆ¹⁹  ,
 Davor ZANELLA²⁰  , Azra BAKRAČ²¹  , Subha AVDIĆ²²  ,
 Avdo SOFRADŽIJA²³   & Branko GLAMUZINA²⁴  

¹Institute of Oceanography and Fisheries, Šetalište Ivana Meštrovića 63, 21000 Split, Croatia.

²University of Zagreb, Faculty of Teacher Education, Savska 77, 10000 Zagreb, Croatia.

^{3,4}University of Zagreb, Faculty of Science, Rooseveltov trg 6, 10000 Zagreb, Croatia.

⁵Zagrebačka 2b, 71000 Sarajevo, Bosnia and Herzegovina.

⁶University of Sarajevo, Institute for Genetic Engineering and Biotechnology,
Zmaja od Bosne 8, 71000 Sarajevo, Bosnia and Herzegovina.

⁷University of Banja Luka, Faculty of Natural Sciences and Mathematics,
Mladena Stojanovića 2, 78000 Banja Luka, Republic of Srpska, Bosnia and Herzegovina.

⁸National Museum of the Czech Republic, Václavské Náměstí 68, 115 79 Prague, Czech Republic.

⁹University of Tuzla, Faculty of Science, Univerzitetska 4, 75000 Tuzla, Bosnia and Herzegovina.

¹⁰Society for Genetic Conservation of Bosnia and Herzegovina Endemic and Autochthonous
Resources GENOFOND, Zmaja od Bosne 8, 71000 Sarajevo, Bosnia and Herzegovina.

¹¹University of Mostar, Faculty of Agronomy and Food Technology,
Biskupa Čule bb 88000 Mostar, Bosnia and Herzegovina.

^{12,13}Hutovo Blato Nature Park Public Institution, Karaotok bb, 88307 Višići, Bosnia and Herzegovina.

^{14,15}University of Sarajevo, Faculty of Science, Zmaja od Bosne 33-35,
71000 Sarajevo, Bosnia and Herzegovina.

¹⁶University of Sarajevo, Institute for Genetic Engineering and Biotechnology,
Zmaja od Bosne 8, 71000 Sarajevo, Bosnia and Herzegovina.

¹⁷Deceased [23 May 2025]. Former adress: University of Banja Luka, Faculty of Natural Sciences and
Mathematics, Mladena Stojanovića 2, 78000 Banja Luka, Republic of Srpska, Bosnia and Herzegovina.

¹⁸Institute of Oceanography and Fisheries, Šetalište Ivana Meštrovića 63, 21000 Split, Croatia.

^{19,20}University of Zagreb, Faculty of Science, Rooseveltov trg 6, 10000 Zagreb, Croatia.

^{21,22}University of Bihać, Biotechnical Faculty,

Luke Marjanovića bb, 77 000 Bihać, Bosnia and Herzegovina.

²³Zmaja od Bosne 70, 71000 Sarajevo, Bosnia and Herzegovina.

²⁴University of Dubrovnik, Department of Applied Ecology, Ćire Carića 4, 20000 Dubrovnik, Croatia.

*Corresponding author: marko.caleta@ufzg.hr

¹Email: tutman@izor.hr

³Email: zoran.marcic@biol.pmf.hr

⁴Email: ivana.buj@biol.pmf.hr

⁵Email: aadem@hotmail.com

⁶Email: belma.kalamujic@ingeb.unsa.ba

⁷Email: dragojla.golub@pmf.unibl.org

⁸Email: radek.sanda@nm.cz

⁹Email: avdul.adrovic@untz.ba

¹⁰Email: pavle.spasojevic7@gmail.com

¹¹Email: pavlicevicj@gmail.com

¹²Email: irenar.2010@gmail.com

¹³Email: nzovko.g@gmail.com

¹⁴Email: e.haskovic@pmf.unsa.ba

¹⁵Email: rskrijelj@pmf.unsa.ba

¹⁶Email: amrakazic@hotmail.com

¹⁸Email: dulcic@izor.hr

¹⁹Email: perica.mustafic@biol.pmf.hr

²⁰Email: davor.zanella@biol.pmf.hr

²¹Email: azra_bakrac_beciraj@yahoo.com

²²Email: subhaavdic8@gmail.com

²⁴Email: branko.glamuzina@unidu.hr

Abstract. In terms of ichthyology, Bosnia and Herzegovina (BiH) is one of the most interesting parts of Southeast Europe, due to its rich biodiversity and high level of endemism. Despite its relevance, the entire territory has been poorly explored. Here, we provide an updated inventory of the current state of knowledge on fishes, including lampreys, from the freshwaters of BiH by hydrographic basin, with recent distributional data and updated taxonomic status reviewed and compared with previous lists. The checklist was compiled based on the existing scientific and grey literature, technical reports, scientific congresses, academic dissertations, and unpublished/personal observations. In total, 123 species including diadromous and euryhaline fishes have been documented in BiH freshwaters to date. Of these, 110 are primarily freshwater. In comparison to the last published monography (Sofradžija 2009), we present a 9% increase in species number (11 species), resulting mainly from taxonomic re-evaluations of existing taxa on the basis of new information and the adoption of a new changes in the taxonomic status of several species. Among the valid primarily freshwater species, 87 are native and 23 are non-native. A total of 38 endemic species have restricted distribution, and are threatened by numerous anthropogenic pressures. Four species are considered endemic only to BiH: *Cobitis herzegoviniensis* Buj & Šanda, 2014; *Phoxinellus pseudalepidotus* Bogutskaya & Zupančič, 2003; *Telestes dabar* Bogutskaya, Zupančič, Bogut & Naseka, 2012; and *T. metohiensis* (Steindachner, 1901). In total, 75 genera and 34 families are represented: Leuciscidae is represented by 37 species, the Salmonidae by 13, followed by the Cyprinidae, Cobitidae and Percidae, each with eight species. The native species richness follows a pattern similar to that observed in other southern European countries. A national list of endangered species has not yet been proposed to BiH and management strategies for their protection or conservation are also not implemented. Hopefully, this updated checklist will serve as a basis for future research aimed at understanding the origin and status of conservation of the BiH fishes diversity, and supporting effective management and conservation programmes.

Keywords. Adriatic Basin, Danube River Basin, endemic species, ichthyofauna, Southeast Europe.

Tutman P., Čaleta M., Marčić Z., Buj I., Hamzić A., Kalamujić Stroil B., Golub D., Šanda R., Adrović A., Spasojević P., Pavličević J., Rozić I., Zovko N., Hasković E., Škrijelj R., Kazić A., Dekić R., Dulčić J., Mustafić P.,

Zanella D., Bakrač-Bećiraj A., Avdić S., Sofradžija A. & Glamuzina B. 2026. Freshwater fishes (Petromyzonti, Elasmobranchii, and Actinopteri) of Bosnia and Herzegovina: an updated inventory on the biodiversity and distribution. *European Journal of Taxonomy* 1042: 1–166. <https://doi.org/10.5852/ejt.2026.1042.3201>

Introduction

Southeast Europe hosts a unique freshwater ichthyofauna strikingly different from the rest of Europe. The majority of species are endemic to small sections of the Mediterranean catchments, while most of the species native to the non-Mediterranean basins of Europe are not present in Mediterranean freshwaters (Bianco 1990; Economidis & Bănărescu 1991; Bănărescu 1992; Bohlen & Ráb 2001; Bobori & Economidis 2006; Kottelat & Freyhof 2007; Geiger *et al.* 2014; Oikonomou *et al.* 2014). The geographic isolation of the Southeast Europe region from other European basins by the Alps and Dinaric Mountain range, together with catchment discreteness, climatic conditions, and the complex geologic history of the area appear to be the main reasons for this distinctiveness (Bianco 1990, 1995; Bănărescu 1992; Hewitt 1999; Šlechtová *et al.* 2004). One country that stands out at the crossroads of the Danube and Adriatic Basins is Bosnia and Herzegovina (hereinafter referred to as BiH). In BiH, like in several other circum-Mediterranean countries, an exceptional level of freshwater fishes fauna richness and diversity is present, with many endemic species with restricted distributional ranges (Vuković 1977a; Vuković & Sofradžija 1987; Crivelli 1996; Smith & Darwall 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Darwall *et al.* 2014; Tutman *et al.* 2017a, 2017b, 2020a, 2020b), largely due to hydrographic isolation, and the occurrence of refuge areas during Pleistocene glaciations (Skoulikidis *et al.* 2022). However, despite the remarkable biodiversity present in both the Danube and Adriatic Basins, the BiH lamprey and other fishes fauna remains surprisingly poorly known and in need of a thorough revision (Tutman *et al.* 2009a, 2012b, 2013, 2017b, 2020a, 2020b).

Indeed, the state of knowledge of freshwater fishes of BiH lags far behind that of other countries of the Mediterranean Sea Basin (Koutsikos *et al.* 2012; Bianco 2014; Barbieri *et al.* 2015; Povž *et al.* 2015; Čaleta *et al.* 2019; Marić 2019). Furthermore, zoogeographic analyses of the freshwater ichthyofauna are still incomplete, with large areas not covered by intensive surveys, and therefore the actual distribution ranges for many species remain to be determined properly (Tutman *et al.* 2017a, 2020a, 2020b). Even though some groups have been thoroughly revised (Tutman *et al.* 2017a, 2020a, 2020b), there has been a strong need for a thorough revision of the overall diversity of the ichthyofauna of the region, especially given recent advances in molecular tools applied to taxonomy (Buj *et al.* 2010, 2014, 2015a; Palandačić *et al.* 2010, 2015, 2017). However, with the exception of Dekić *et al.* (2024), no general review of the actual taxonomic status and geographic distribution of many species in BiH is available. Therefore, the knowledge of the distribution and current status of populations, including the precise taxonomic identity of species in particular regions is still insufficient (Tutman *et al.* 2009a, 2012b, 2013, 2017b, 2020a, 2020b). This is particularly significant since the advent of biomolecular research and biogeographical theories about the origins, especially of peri-Mediterranean endemics, have led to great shifts in many taxonomic aspects of European and BiH freshwater fishes over the past 20 years (e.g., Palandačić *et al.* 2010, 2015, 2017; Perea *et al.* 2010; Vucić *et al.* 2017; Sabolić *et al.* 2021; Buj *et al.* 2022; Lukač *et al.* 2023). These data provide the basic information for conservation management, while also contributing to more detailed knowledge of fishes distribution at the continental scale. Such a compilation is needed, since the general distribution patterns of species can profoundly impact conservation at both the national (BiH) and European scales (Geiger *et al.* 2014).

The need of a thorough revision of the ichthyofauna of BiH is especially noticeable considering significant advancements in recent research related to the phylogeny and distribution of species (Razpet *et al.* 2007; Snoj *et al.* 2002a, 2002b, 2010; Palandačić *et al.* 2015, 2024; Vucić *et al.* 2017; Sabolić *et al.* 2021; Tougard *et al.* 2021; Buj *et al.* 2022). One of the key tasks for national conservation and

management should be ensuring detailed knowledge of the lamprey and fishes diversity according to river catchments. Consequently, the periodic updating of checklists according to newly recorded changes in species composition and distribution is a necessary tool for national and regional fisheries management (Tutman *et al.* 2017b, 2020a, 2020b; Tutman & Glamuzina 2021).

Previous general checklists of freshwater lampreys and other fishes in BiH include several monographs: Vuković (1963a, 1977a), Vuković & Ivanović (1971), Bogut *et al.* (2006), Sofradžija (2009), and Hamzić (2024). The foundations of contemporary ichthyological knowledge in BiH were laid with the early works of Vuković (1963a), who listed 103 species and subspecies in 22 families and 45 genera, and Vuković (1977a), who reported 106 species and subspecies in 26 families and 64 genera. According to the two most recent reviews of the freshwater ichthyofauna of BiH (Sofradžija 2009; Hamzić 2024), 118 and 137 species were listed as the total number in the national freshwater ichthyofauna, respectively. With a more detailed analysis of both recent reviews, this discrepancy in the total number of species refers to numerous changes in systematics, taxonomy and nomenclature, as well as different interpretations on the validity of certain species. For example, Sofradžija (2009) did not fully accepted the proposed changes in the taxonomic status of certain species presented by Kottelat & Freyhof (2007). In recent years, several new distributional records, discoveries and descriptions of new species have been reported, increasing the knowledge about the biodiversity of freshwater fishes, including lampreys found in BiH (Šanda & Kovačić 2009; Buj *et al.* 2010, 2014; Tutman *et al.* 2009a, 2012b, 2013, 2017a, 2017b; Čolić *et al.* 2018). The lack of consensus over the exact number of freshwaters species of fish in BiH also highlights the need for a systematic review (Tutman *et al.* 2013, 2017a, 2017b).

History of ichthyological research in Bosnia and Herzegovina

Research on the ichthyofauna of BiH has a very long tradition that began in the mid-19th century (Heckel & Kner 1858). Since then, ichthyological research has been constant (Heckel & Kner 1858; Steindachner 1882a, 1882b, 1882c, 1883, 1901; Karaman 1928, 1938, 1963, 1972; Taler 1944, 1945a, 1945b, 1953a, 1953b; Vuković 1963a, 1977a, 1982b; Vuković & Ivanović 1971; Sofradžija 1977, 2009), although with varying intensity throughout the years.

The first list of freshwater species of fish of BiH was published by Heckel & Kner (1858), with the description of 11 new species. Later, Steindachner (1882a, 1882b, 1882c, 1883, 1901) and then Trgovčević (1905), Kolombatović (1907) and Heintz (1908) continued to explore the BiH freshwater ichthyofauna. The work of Heintz (1908) on local salmonids is particularly noteworthy. Research on the systematics of certain species of fishes of BiH, which was then a part of Yugoslavia, was also advanced by Karaman (1926, 1927, 1938), who described nine species, mostly salmonids. However, the taxonomic status of most of the species described by him is still controversial (Kottelat & Freyhof 2007, Georgiev 2011). In the same period, there were reports on the fishes from Sarajevo and its surroundings (Zaplata & Taler 1932), and on the Neretva River salmonids by Čurčić (1938) and by Taler (1944, 1945a, 1945b).

After the Second World War, research intensified. Jedlička (1947) published a list of species of fish, which was soon followed by a monograph by Taler (1953a) entitled “*Distribution and List of Freshwater Fishes of Yugoslavia*”. This study was particularly significant because it not only listed the species of fish but also considered the distribution and systematic relationships of freshwater fishes of the region. In the second half of the 20th century, especially in the 1960s and 1970s, the number of scientists engaged in the topic increased, resulting in a substantial advancement and increase of the ichthyologic knowledge in BiH (see references in Sofradžija 2009). The most notable were researchers of the period were Drs. Tonko Šoljan and Tihomir Vuković. Three published monographs, “*Fishes of Bosnia and Herzegovina*” in two editions (Vuković 1963a, 1977a) and “*Freshwater Fishes of Yugoslavia*” (Vuković & Ivanović 1971) presented detailed descriptions of the species and subspecies of freshwater fishes, including lampreys found in BiH.

In the 1970s and 1980s, several studies were published on freshwater fishes in BiH, including Aganović & Kapetanović (1970, 1971, 1978), Aganović & Vuković (1971), Aganović *et al.* (1974, 1976), Kosorić (1969, 1972a, 1972b, 1974, 1976, 1977a, 1977b, 1978, 1981), Sofradžija (1977, 1979, 1982a, 1982b), Mikavica (1987a, 1987b, 1988a, 1988b), Guzina & Vuković (1972, 1977, 1978, 1987), Berberović (1967, 1980), Vuković & Sofradžija (1987). Although the war in BiH (1992–1995) significantly hindered scientific ichthyological efforts, later studies resulted in remarkable progresses on the knowledge of the field (e.g., Korjenić 1998, 1999, 2003; Škrijelj & Mašović 2001; Škrijelj 2002; Hamzić 2002; Ivanković *et al.* 2011, 2017; Francuski *et al.* 2019; Markotić *et al.* 2018; Kalamujić Stroil *et al.* 2019, 2020, 2021; Pilić *et al.* 2020; Žujo-Zekić *et al.* 2020; Bajrić *et al.* 2021a). Particularly notable are monographs presented by Hamzić (2002) on aquaculture in BiH, Bogut *et al.* (2006), and Sofradžija (2009) and also Glamuzina *et al.* (2013), who revised the list of the ichthyofauna in BiH and adapted it to updated views on the freshwater lamprey and fishes systematics at that time. The publication provided by Drešković *et al.* (2011) supported government institutions in preparing BiH for the Natura 2000 network, and included 50 species of fish listed in Annex II of the Bern directive. Tutman *et al.* (2009a, 2012b, 2013) recorded new changes in the composition of the lower reaches of the Neretva River in BiH, also presenting a detailed inventory of the taxonomic status and distribution of loaches (Tutman *et al.* 2017a, 2017b), lampreys (Tutman *et al.* 2020a) and freshwater gobies (Tutman *et al.* 2020b). In a recent article, Dekić *et al.* (2024) emphasized the need for a thorough revision of the taxonomical state of the BiH ichthyofauna, as well as the need for more field expeditions and the use of molecular tools that might help to solve many taxonomic doubts and likely identification mistakes in previous studies present (Bogut *et al.* 2006; Sofradžija 2009). Dekić *et al.* (2024) also provide a taxonomic summary overview of the freshwater ichthyofauna of BiH but, in some cases, they did not consider the results of modern genetic, phylogenetic or phylogeographic studies.

Current state of the knowledge on the taxonomy and geographic distribution

The contemporary fishes fauna composition in BiH, especially endemic primary or near-primary freshwater species, is the result of long-term speciation related to numerous processes that took place during the Miocene to Pliocene (Perea *et al.* 2010; Lukač *et al.* 2023). The current distribution patterns and genetic diversity of many species in BiH have been contextualized in biogeographical hypotheses by Bănărescu (1989, 1992), and Bianco (1990), reflecting findings on evolutionary histories of fishes in neighbouring Croatia (Ćaleta *et al.* 2019). This valuable biodiversity is now under significant anthropogenic threat (e.g., pollution, habitat destruction, dams, introduction of non-native species) which has affected the fishes fauna (Hamzić 2024).

In terms of ichthyology, BiH is one of the most interesting parts of Southeast Europe due to the high number of endemic species. Despite its remarkable ichthyofaunistic importance, its fishes fauna is still poorly explored and known. The zoogeographical analysis of the BiH freshwater ichthyofauna is partial and incomplete, with large areas not covered by intensive surveys, and the actual distribution ranges for many species have not yet been clearly determined. The taxonomic status of several species remains disputed as they are currently the subject of research (e.g., Kottelat & Freyhof 2007), particularly within the genera *Salmo* Linnaeus, 1758 (Razpet *et al.* 2007; Snoj *et al.* 2010; Georgiev 2011); *Romanogobio* Bănărescu, 1961; *Scardinius* Bonaparte, 1837 (Freyhof *et al.* 2006; Sabolić *et al.* 2021); *Alburnus* Linnaeus, 1758 (Buj *et al.* 2010; Vucić *et al.* 2017); *Phoxinus* Rafinesque, 1820 (Bianco & De Bonis 2015; Palandačić *et al.* 2015; Vucić *et al.* 2017); *Squalius* Bonaparte, 1837 (Buj *et al.* 2020a); *Telestes* Bonaparte, 1837 (Buj *et al.* 2017, 2022); *Knipowitschia* Iljin, 1927 (Šanda & Kovačić 2009; Tougard *et al.* 2021). These genera encompass species with a narrow distribution range that are often under strong anthropogenic pressures and, consequently, high environmental risk. Therefore, elucidating their taxonomic status should be a primary task, especially considering their legal protection and associated potential risks of endangerment. Consequently, it is necessary to comprehensively collect and analyse literature data on freshwater fishes, including lampreys of BiH, revise some of the proposed taxonomic changes, and update regional lists with recently introduced species, while considering the latest scientific achievements.

The survival of the native freshwater fishes fauna in BiH and their natural habitats is threatened by a wide range of increasing anthropogenic factors, such as dam construction, pollution, habitat destruction, and the introduction of non-native species (Drešković *et al.* 2011; Tutman *et al.* 2012a; Hamzić 2024; Dekić *et al.* 2024). Therefore, to ensure the survival of endemic species, it is necessary to implement measures to protect and preserve the habitats and ecosystems where they live.

This study is a continuous effort to elucidate the taxonomic status of the freshwater fishes, including lampreys, in BiH that started in previous monographs and studies by Vuković (1963a, 1977a), Vuković & Ivanović (1971), Bogut *et al.* (2006), Sofradžija (2009), and Hamzić (2024). Furthermore, this contribution is complementary to Dekić *et al.* (2024), who presented a valuable overview of the freshwater ichthyofauna of BiH, but left somehow unsolved the taxonomic position of several species, particularly those with problems in taxonomy and systematics, which we tried to elucidate in this study.

Here, we present an updated annotated checklist of the currently known freshwater fishes, including lampreys, in BiH freshwaters by hydrographic catchments with clarified nomenclature and taxonomic status. This species list includes all recent data on distribution and taxonomic status, discussion about some questionable previously reported and recently described species. Considering the current state of ichthyological knowledge in BiH, the purpose is to critically analyse the distribution of previously reported species and clarify the current knowledge on them. This information will provide a scientific baseline for decision-makers and broader public interest regarding conservation requirements for these species in BiH.

Material and methods

Sources

The updated checklist of freshwater fishes of BiH is compiled based on data published in the recent scientific and grey literature, technical reports, conference proceedings from scientific congresses, academic dissertations, unpublished/personal observations and field sampling of all major catchments conducted since 1997 to present. Classification of orders and families follow Van der Laan *et al.* (2025).

Within each family and genus, species are listed in alphabetic order. Only those species considered as valid in the recent scientific literature are included. The designation of species in the checklist is arranged in a way that the ordinal numbers (1., 2., ...), are used to designate only exclusively freshwater species. In addition to the exclusively freshwater species, data on euryhaline species that inhabit brackish and transitional waters and migratory species of fish are also provided. Species such as the mullets, seabasses and seabreams were not included in the list but were considered as occasional, with the abbreviation ME (Marine Euryhaline), with corresponding numbers (ME1, ME2,...). The list also includes species that are considered (RE) with corresponding numbers (RE1, RE2,...), and one marine species (M) with a very unusual record (*Raja miraletus* Linnaeus, 1758). Furthermore, the taxonomic status of some species mentioned in earlier lists (Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009; Hamzić 2024) was re-examined, if their taxonomic status had changed.

After the scientific name, a common local name from Vuković (1977a) and Sofradžija (2009) is provided, followed by the common English name. After the common name, the distributional range for each species is provided, together with the source for considering the species as endemic, native or non-native. Endemic species are particularly highlighted. The occurrence in catchments (Danube River and Adriatic Basin) of each species is based on literature reviews and field knowledge. To complement this distribution and diversity data, the precise distribution of each endemic species regarding their known area is also provided, considering: Adriatic Basin Endemic (sensu lato, i.e., present in BiH, and also distributed in Italy, Slovenia, Croatia (CRO) or Montenegro) – (ABE); Neretva River Catchment Endemic (sensu stricto, i.e., distributed only in the Neretva River system in BiH and Croatia) – (NCE); and Danube

River Basin Endemic (DBE) and Cetina River Catchment Endemic (distributed only in the Cetina River system in BiH and Croatia) – (CCE). Species endemic only to BiH are marked with the abbreviation BiH, followed by the catchment distribution information. Several records on the checklist represent expansions of the known range or new records of the species in BiH. We considered a species as a new record in BiH when it was not listed either in Bogut *et al.* (2006) or Sofradžija (2009).

A list of all freshwater fish species recorded in the BiH according to literature to date is also presented in Supp. file 1. Their distribution across the basins and catchments according to literature and knowledge of the authors are presented in Table 1, and information on species conservation status according to IUCN is presented in Table 2 and in the species treatments, whenever appropriate.

Study area

Although its area of 51 209.20 km² makes BiH a territorially small country, the rivers of two large European catchments flow through it, the larger Danube (also called the Black Sea) (38 719 km² or 75.7% of total BiH surface area) and the smaller Adriatic Basin (12 410 km² or 24.3%), divided by the Dinaric Mountain range (Fig. 1). The Dinaric Mountains comprise the largest continuous karst area in Europe. They are parallel to the eastern Adriatic coast and have typical karst conditions, such as dolines and karst poljes above ground, sinkholes, caves, and underground streams below the ground, forming a complex water network connecting surface and underground water bodies into which certain species enters during the dry season (Palandačić *et al.* 2024).

The majority of the Danube River Basin is drained by the Sava River and its tributaries as far as the Croatian border to the north, while the Adriatic Basin is mostly drained by the Neretva River and its tributaries that drain into the Adriatic Sea in Croatia. Each catchment is characterised by a different ichthyofaunal composition: in general, the Danube River Basin has species present in other parts




Fig. 1. Map of Bosnia and Herzegovina showing the two main basins, the Danube River Basin (green shaded) and the Adriatic Basin (blue shaded).

Table 1 (continued on the next 5 pages). List of all lampreys and other species of fish recorded in freshwaters of Bosnia and Herzegovina as well as their distribution across the basins and catchments according to literature and this study. Abbreviation: ME = Marine Euryhaline.

Family	No.	Species	Danube River Basin										Adriatic Sea Basin				
			Rivers catchments										Neretva	Trebišnjica	Cetina		
			Sava	Una	Vrba	Bosna	Drina	Ukrina	Korana	Glina							
Petromyzontidae	1.	<i>Eudontomyzon vladkykovi</i>	+	+	+	+	+	+	+	+	+						
	2.	<i>Lampetra soljani</i>													+		
	ME1	<i>Petromyzon marinus</i>													+		
Rajidae	M1	<i>Raja miraletus</i>													+		
Acipenseridae	3.	<i>Huso ruthenus</i>	+	+	+	+	+	+									
Anguillidae	4.	<i>Anguilla anguilla</i>	+												+		+
Clupeidae	ME2	<i>Alosa fallax</i>													+		
Cobitidae	5.	<i>Cobitis elongata</i>	+	+	+	+	+	+							+		
	6.	<i>Cobitis elongatooides</i>	+												+		
	7.	<i>Cobitis herzegoviniensis</i>													+		
	8.	<i>Cobitis illyrica</i>													+		
	9.	<i>Cobitis narentana</i>													+		
	10.	<i>Misgurnus fossilis</i>	+	+	+	+	+	+								+	
	11.	<i>Sabanejewia balcanica</i>	+	+	+	+	+	+								+	
	12.	<i>Sabanejewia bulgarica</i>	+	+	+	+	+	+									
Nemacheilidae	13.	<i>Barbatula barbatula</i>	+	+	+	+	+	+							+		
Cyprinidae	14.	<i>Aulopyge huegelii</i>		+											+		+
	15.	<i>Barbus balcanicus</i>	+	+	+	+	+	+									
	16.	<i>Barbus barbus</i>	+	+	+	+	+	+									
	17.	<i>Carassius auratus</i>	+	+	+	+	+	+							+		
	18.	<i>Carassius carassius</i>	+	+	+	+	+	+							+		+
	19.	<i>Carassius gibelio</i>	+	+	+	+	+	+							+		+
	20.	<i>Carassius langsdorfi</i>													+		
	21.	<i>Cyprinus carpio</i>	+	+	+	+	+	+							+		+

Table 1 (continued). List of all lampreys and other species of fish recorded in freshwaters of Bosnia and Herzegovina as well as their distribution across the basins and catchments according to literature and this study. Abbreviation: ME = Marine Euryhaline.

Family	No.	Species	Danube River Basin										Adriatic Sea Basin			
			Sava	Una	Vrbaš	Bosna	Drina	Rivers catchments				Neretva	Trebišnjica	Cetina		
								Ukrina	Korana and Glina							
Xenocyprididae	22.	<i>Ctenopharyngodon idella</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	23.	<i>Hypophthalmichthys molitrix</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	24.	<i>Hypophthalmichthys nobilis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Tincidae	25.	<i>Tinca tinca</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Achieleognathidae	26.	<i>Rhodeus amarus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Gobionidae	27.	<i>Gobio obtusirostris</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	28.	<i>Pseudorasbora parva</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	29.	<i>Romanogobio carpathorossicus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	30.	<i>Romanogobio uranoscopus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	31.	<i>Romanogobio vladkykovi</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Leuciscidae	32.	<i>Abramis brama</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	33.	<i>Alburnoides bipunctatus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	34.	<i>Alburnus alburnus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	35.	<i>Alburnus neretvae</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	36.	<i>Alburnus sava</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	37.	<i>Ballerus ballerus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	38.	<i>Ballerus sapa</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	39.	<i>Blicca bjoerkna</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	40.	<i>Chondrostoma knerii</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	41.	<i>Chondrostoma nasus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+

Table 1 (continued). List of all lampreys and other species of fish recorded in freshwaters of Bosnia and Herzegovina as well as their distribution across the basins and catchments according to literature and this study. Abbreviation: ME = Marine Euryhaline.

Family	No.	Species	Danube River Basin										Adriatic Sea Basin				
			Rivers catchments										Neretva	Trebišnjica	Cetina		
			Sava	Una	Vrba	Bosna	Drina	Ukrina	Korana and Glina								
Odontobutidae	88.	<i>Perccottus glenii</i>	+														
Oxudereidae	89.	<i>Knipowitschia radovici</i>															+
	90.	<i>Ninnigobius canestrinii</i>															+
	91.	<i>Orsinigobius croaticus</i>															+
Gobiidae	92.	<i>Babka gymnotrachelus</i>	+					+									
	93.	<i>Neogobius fluviatilis</i>	+	+	+	+											
	94.	<i>Neogobius melanostomus</i>	+	+	+	+											
	95.	<i>Ponticola kessleri</i>	+	+													
	96.	<i>Proterorhinus semilunaris</i>	+														
		ME4	<i>Platichthys flesus</i>														
Pleuronectidae	ME5	<i>Atherina boyeri</i>															+
Poeciliidae	97.	<i>Gambusia hoolbroki</i>															+
	ME6	<i>Chelon auratus</i>															+
Mugilidae	ME7	<i>Chelon labrosus</i>															+
	ME8	<i>Chelon ramada</i>															+
	ME9	<i>Chelon saliens</i>															+
	ME10	<i>Mugil cephalus</i>															+
Bleniidae	98.	<i>Salariaopsis fluviatilis</i>															+

Table 1 (continued). List of all lampreys and other species of fish recorded in freshwaters of Bosnia and Herzegovina as well as their distribution across the basins and catchments according to literature and this study. Abbreviation: ME = Marine Euryhaline.

Family	No.	Species	Danube River Basin												Adriatic Sea Basin			
			Sava	Una	Vrba	Bosna	Drina	Rivers catchments				Neretva	Trebišnjica	Cetina				
								Ukrina	Korana	and	Glina							
Percidae	99.	<i>Gymnocephalus baloni</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	100.	<i>Gymnocephalus cernua</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	101.	<i>Gymnocephalus schraetser</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	102.	<i>Perca fluviatilis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	103.	<i>Sander lucioperca</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	104.	<i>Sander volgensis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	105.	<i>Zingel streber</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	106.	<i>Zingel zingel</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Gasterosteidae	107.	<i>Gasterosteus aculeatus</i>																+
Cottidae	108.	<i>Cottus gobio</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Centrarchidae	109.	<i>Lepomis gibbosus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	110.	<i>Micropterus salmoides</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Moronidae	ME11	<i>Dicentrarchus labrax</i>																+
Sparidae	ME12	<i>Sparus aurata</i>																+
	123	Total number of reported species (freshwater + euryhaline + marine)	71	65	65	62	66	66	57	50	76	40	22	83	77			

of Central Europe, while the Adriatic Basin area is characterised by a number of endemic forms (Vuković 1977a; Kottelat & Freyhof 2007; Sofradžija 2009; Buj *et al.* 2015a, 2017, 2020b).

The Sava River is the most prominent river connecting Croatia, BiH, and Serbia. Together with its tributaries, it constitutes one of the major catchments of Southeast Europe, covering a total area of approximately 97 713.20 km². The Sava River is also one of the largest catchments of the Danube River Basin, accounting for 12% of its catchment area in BiH (Jukić 2008). With the exception of the Neretva River, all the major rivers in BiH flow into the Sava River, the largest tributary of the Danube. The Drina River is the Sava River's largest and most important tributary. The Drina River catchments extend into four countries: Montenegro, BiH, Serbia, and a very small part extends into Albania. Other important catchments include the: Una, Vrbas, Bosna, Ukrina, Korana, and Glina Rivers. The Adriatic Basin area of BiH, in turn, consists of three catchments: Neretva, Trebišnjica and Cetina Rivers (Fig. 2). The Neretva River catchment has an area of 10 100 km² or 81.4% of the total Adriatic Basin area of BiH, the Trebišnjica River catchment area has 2250 km², while the Cetina River catchment spreads over 2310 km² (18.6% of the total Adriatic Basin area of BiH) (Hamzić 2024). The Neretva and Cetina rivers flow into the Adriatic Sea, while the Trebišnjica River



Fig. 2. Map of Bosnia and Herzegovina showing the main catchments. From: Hamzić A. 2024. *Biodiverzitet slatkovodnih riba Bosne i Hercegovine*. Buybook, Sarajevo.

is a complex of underground and above-ground watercourses that sinks in the western part of Popovo Polje in BiH, thus connecting with the Neretva River. In the Herzegovina region, there is a massive karst area (covering more than 4000 km²) that contain numerous flowing underground rivers and streams, that rise as springs.

The freshwater fauna of the Adriatic Basin in BiH is separated from the Danube River Basin by the Dinaric Mountains and thus contributing to the isolation of the Adriatic freshwater fauna which contains many endemic forms and a few non-endemic species (Economidis & Bănărescu 1991; Oikonomou *et al.* 2014; Ludoški *et al.* 2021). In the overall structure of Southeast Europe and the ichthyogeographic provinces presented by Economidis & Bănărescu (1991), BiH is considered as part of the Dalmatian subdivision that extends from northern Slovenia to southern Montenegro. This area is a hotspot of ichthyological diversity for several cyprinid genera, like the unique *Delminichthys* Freyhof, Lieckfeldt, Bogutskaya, Pitra & Ludwig, 2006, followed by *Telestes*, *Squalius* or *Scardinius* (Kottelat & Freyhof 2007), which are primarily associated with the typical karst areas of groundwater. The Neretva River catchment deserves special attention, as this catchment contains a high number of unique endemic freshwater fishes (Mrakovčić *et al.* 1995; Economidis & Bănărescu 1991; Bănărescu & Herzig-Straschil 1998; Bogutskaya & Zupančič 1999; Zupančič & Bogutskaya 2002; Neuburg *et al.* 2023). It is noteworthy that several new species of fish have been described from this catchment in recent decades (Mrakovčić *et al.* 1996; Bogutskaya & Zupančič 2003; Kovačić 2005; Buj *et al.* 2010; Bogutskaya *et al.* 2012; Bianco & De Bonis 2015; Tutman *et al.* 2017a).

Results

Species account of the fishes, including lampreys, reported in freshwaters of BiH

Class Petromyzonti

Order Petromyzontiformes Berg, 1940

Family Petromyzontidae Bonaparte, 1831

Based on the available literature, lampreys are the least studied and most poorly known group of freshwater vertebrates in BiH, though the composition and geographical distribution of the group is better known following a recent revision (Tutman *et al.* 2020a). While eight species were previously reported as present in BiH (*Eudontomyzon danfordi* Regan, 1911; *E. mariae* (Berg, 1931); *E. vladykovi* Oliva & Zandrea, 1959; *Lampetra fluviatilis* (Linnaeus, 1758); *L. planeri* (Bloch, 1784); *L. soljani* Tutman, Freyhof, Dulčić, Glamuzina & Geiger, 2017; *Lethenteron zanandreae* Vladykov, 1955; *Petromyzon marinus* Linnaeus, 1758) (Vuković 1977a; Sofradžija 2009; Tutman *et al.* 2009a), only three species (*E. vladykovi*, *L. soljani*, *P. marinus*) have recently been confirmed, with records in both the Danube and Adriatic Basins (Tutman *et al.* 2020a). Consequently, five species previously reported in BiH are not included in this checklist, since their occurrence in the region is dubious or not based on confirmed records (*E. danfordi*, *E. mariae*, *L. fluviatilis*, *L. planeri*, and *L. zanandreae*).

Genus *Eudontomyzon* Regan, 1911

Eudontomyzon danfordi Regan, 1911

Vernacular names

Dunavska paklara. Carpathian lamprey.

Previous names used for the species

Lampetra danfordi (Regan, 1911); *Lampetra danfordi danfordi* (Regan, 1911).

Remarks

Distribution of this species is limited to Tisza and Timis drainages in Romania, Slovakia, Ukraine and Hungary (Kottelat & Freyhof 2007). There are no reliable scientific records in BiH freshwaters (Tutman *et al.* 2020a), and all earlier literature that mention this species (Vuković 1963a, 1977a; Kosorić 1981; Šorić 1998) are most likely related to *Eudontomyzon vladykovi*. Therefore, this species is not included in the checklist.

Literature

See *Eudontomyzon vladykovi*.

Eudontomyzon mariae (Berg, 1931)

Vernacular names

Ukrajinska paklara. Ukrainian brook lamprey.

Previous names used for the species

Lampetra mariae Berg, 1931.

Remarks

This species seems to be restricted to Don and Volga basins (Levin *et al.* 2016). There is no scientific confirmation of its presence in BiH waters (Tutman *et al.* 2020a), and all earlier literature data that mention this species (Vuković 1963a, 1977a; Kosorić 1981; Šorić 1998) are most likely related to *E. vladykovi*. Therefore, it is not included in the checklist.

Literature

See *Eudontomyzon vladykovi*.

1. *Eudontomyzon vladykovi* Oliva & Zanandrea, 1959

Vernacular names

Dunavska paklara. Danube brook lamprey.

Previous names used for the species

Eudontomyzon danfordi vladykovi (Zanandrea, 1959).

Origin and geographic range in BiH

Endemic (DBE). Danube River Basins.

Remarks

Under the name of *Eudontomyzon vladykovi*, this species was recorded from several localities in BiH, scattered within the river systems of the Una (Šanda & Vukić 2009), Bosna (Golob *et al.* 2015) and Sava River (Golub *et al.* 2018a, 2018b; Čaleta & Marčić, *unpublished data*). Recently, it was recorded in streams on Mt. Vranica (Golob *et al.* 2015) and in the Lašva River (Mustafić *et al.* 2015). Tutman *et al.* (2020a) presented an overview of its taxonomic status and distribution, confirming its presence in the BiH.

Literature

Anonymous 1886; Glowacki 1896; Ćurčić 1910; Zaplata & Taler 1932; Zanandrea 1959; Vuković 1963a, 1977a; Kosorić 1981; Šorić 1998; Holčik & Delić 2000; Mrakovčić *et al.* 2006; Skenderović *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Šanda & Vukić 2009; Drešković *et al.* 2011; Čičić-Močić 2014; Golob *et al.* 2015; Milanović *et al.* 2015; Mustafić *et al.* 2015; Simonović *et al.* 2015; Lukić *et al.* 2016; Adrović 2018; Golub *et al.* 2018a, 2018b; Memić 2018; Tutman *et al.* 2020a; Dekić *et al.* 2024; Hamzić 2024; Pleše & Buj 2024.

Genus *Lampetra* Gray, 1851

Lampetra fluviatilis (Linnaeus, 1758)

Vernacular names

Riječna paklara. European river lamprey.

Previous names used for the species

Petromyzon fluviatilis Linnaeus, 1758; *Petromyzon branchialis* Linnaeus, 1758.

Remarks

Reports of its presence in BiH appear to be wrong identification of specimens (Tutman *et al.* 2020a). It is unlikely that *Lampetra fluviatilis* inhabits BiH because its distribution range is restricted to northern and western Europe with extirpated populations in the western Mediterranean (Kottelat & Freyhof 2007). Previous reports of the species inhabiting the Adriatic Basin in BiH (Vuković 1963a, 1977a; Sofradžija 2009) are most likely misidentifications of *L. soljani* (Tutman *et al.* 2017a, 2020a). Therefore, it is not included in the checklist.

Literature

See *Lampetra soljani*.

Lampetra planeri (Bloch, 1784)

Vernacular names

Potočna paklara. European brook lamprey.

Previous names used for the species

Petromyzon planeri Bloch, 1784.

Remarks

Reports on its occurrence in BiH appear to be wrong identification of specimens (Tutman *et al.* 2020a). It is unlikely that *Lampetra planeri* inhabits BiH because its distribution range is restricted to northern and western Europe, with isolated populations in the western Mediterranean (Kottelat & Freyhof 2007). Previous reports from the Adriatic Basin in BiH (Vuković 1963a, 1977a; Sofradžija 2009) are most likely related to *L. soljani* (Tutman *et al.* 2017a), while those from Danube Basin (Vuković 1963a, 1977a; Kosorić 1976; Sofradžija 2009) are likely misidentifications of *E. vladikovi*. Therefore, it is not included in the checklist.

Literature

See *Eudontomyzon vladikovi* and *Lampetra soljani*.

2. *Lampetra soljani* Tutman, Freyhof, Dulčić, Glamuzina & Geiger, 2017

Vernacular names

Primorska paklara. Šoljan's brook lamprey.

Previous names used for the species

Lampetra zanandreae Vladykov, 1955; *Lethenteron zanandreae* (Vladykov, 1955).

Origin and geographic range in BiH

Endemic (ABE). Adriatic Basin.

Remarks

Lampetra soljani was described from the lower Neretva River in BiH and Croatia (Adriatic Basin) (Tutman *et al.* 2017a), with populations of this species previously regarded as *Lethenteron zanandreae* (Tutman *et al.* 2009a). This non-parasitic species is endemic with a geographic range restricted to the lower section of the Neretva River, from its mouth to about 30 km upstream to the Buna River and in the Neretva River tributaries: Tihaljina/Trebižat River catchment, Krupa River, Bregava River and the Hutovo Blato wetland (Tutman *et al.* 2017a, 2017b, 2020a).

Literature

Kottelat & Freyhof 2007; Tutman *et al.* 2009a, 2009b, 2012a, 2017a, 2020a; Drešković *et al.* 2011; Geiger *et al.* 2014; Milanović *et al.* 2015; Freyhof *et al.* 2020; Pereira *et al.* 2021; Rüber *et al.* 2023.

Lampetra zanandreae Vladykov, 1955

Vernacular names

Po brook lamprey.

Previous names used for the species

Lethenteron zanandreae (Vladykov, 1955).

Remarks

Lampetra zanandreae was considered as distributed in streams in the northern and western Adriatic basin with two disjunct distributional areas: one in streams of the Italian and Slovenian side of the Adriatic basin, and a second in the headwaters of the Neretva and Morača drainages in Croatia, BiH and Montenegro (Holčík & Mrakovčić 1997; Kottelat & Freyhof 2007; Tutman *et al.* 2009a). Recently, populations from the second distributional area were described as a different species, *L. soljani* (Tutman *et al.* 2017a), previously identified as *Lethenteron zanandreae* (Tutman *et al.* 2009a). Therefore, *L. zanandreae* is not included in the list.

Literature

See *Lampetra soljani*.

Genus *Petromyzon* Linnaeus, 1758

ME1 *Petromyzon marinus* Linnaeus, 1758.

Vernacular names

Morska paklara. Atlantic sea lamprey.

Previous names used for the species

Petromyzon fluviatilis Linnaeus, 1758; *Petromyzon lampetra* Pallas, 1814.

Origin and geographic range in BiH

Native, migratory, anadromous. Adriatic Basin.

Remarks

Vuković (1963a, 1977a) and Sofradžija (2009) reported the occurrence of this species in the lower reaches of the Neretva River. Nevertheless, only four records of adult *Petromyzon marinus* have been reported to date, one at Žitomislići approx. 50 km upstream (Holčik *et al.* 2004), one in 2013 near Gabela (city of Čapljina) approx. 25 km upstream, and another in 2014 in the same area (Bregava River, a tributary of the Neretva), approx. 27 km upstream (Tutman *et al.* 2020a), and the last one near Gabela (Glamuzina *et al.* 2019). The latter was a ripe adult male, the first mature individual caught in the eastern Adriatic Sea, indicating potential spawning activity not previously recorded (Holčik *et al.* 2004).

Literature

Vuković 1963a, 1977a; Holčik *et al.* 2004; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Tutman *et al.* 2012a, 2020a; Milanović *et al.* 2015; Glamuzina *et al.* 2019.

2. Class Elasmobranchii Bonaparte, 1838
Order Rajiformes Berg, 1940
Family Rajidae de Blainville, 1816
Genus *Raja* Linnaeus, 1758

M1 *Raja miraletus* Linnaeus, 1758

Vernacular names

Raža modropjega. Brown ray.

Origin and geographic range in BiH

Adriatic Basin.

Remarks

One specimen of *Raja miraletus* was caught in mid-March 2020 in the Neretva River about 25 km upstream of its mouth (Tutman *et al.* 2021). This collection is unusual since it is the first record of this marine species so far inland in the Neretva River, and a rare record in the Mediterranean Sea overall. This could indicate the expansion of saltwater intrusion deeper into the Neretva River. It may be linked to climate change or human activity (water level lowering in the upper areas of the Neretva River).

Literature

Tutman *et al.* 2021.

3. Class Actinopteri Cope, 1871
Order Acipenseriformes Berg, 1940

Family **Acipenseridae** Bonaparte, 1831

Recent literature data (Brownstein & Thomas 2025) suggests that according to molecular data, most European sturgeon species belong to genus *Huso*, except *Acipenser sturio* and *A. oxyrinchus*.

Genus *Acipenser* Linnaeus, 1758

RE1 *Acipenser sturio* Linnaeus, 1758

Vernacular names

Atlantska jesetra. Atlantic sturgeon.

Previous names used for the species

Sturio vulgaris Rafinesque, 1810.

Origin and geographic range in BiH

Native, migratory, anadromous. Danube and Adriatic Basins. Extinct (EX).

Remarks

Atlantic sturgeon used to occur in both the Danube River Basin and in the Adriatic Sea (Vuković 1963a, 1977a). Today, it is considered a regionally extinct species and all recent references (Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009) are based on historical literature.

Literature

Ćurčić 1910; Vuković 1963a, 1977a; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Freyhof *et al.* 2020; Dekić *et al.* 2024.

Genus *Huso* Brandt & Ratzeburg, 1833

RE2 *Huso gueldenstaedtii* Brandt & Ratzeburg, 1833

Vernacular names

Jesetra. Russian sturgeon.

Previous names used for the species

Acipenser güldenstädti Brandt, 1833; *Acipenser gueldenstaedtii* Brandt & Ratzeburg, 1833; *Acipenser aculeatus* Lovetsky, 1834; *Acipenser güldenstädti colchicus* Marti, 1940.

Origin and geographic range in BiH

Native, migratory, anadromous. Danube River Basin. Extinct (EX).

Remarks

Vuković (1963a, 1977a) mentioned the occurrence of the species in BiH because it was common in the rivers of the Danube River Basin in the past. Recently, however, there are no confirmed records of the species in the rivers of BiH.

Literature

Anonymous 1886; Glowacki 1896; Ćurčić 1910; Vuković 1963a, 1977a; Vlasenko *et al.* 1989; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Freyhof *et al.* 2020; Dekić *et al.* 2024.

RE3 *Huso huso* (Linnaeus, 1758)

Vernacular names

Moruna. Beluga.

Previous names used for the species

Acipenser huso Linnaeus, 1758; *Acipenser beluga* Forster, 1767.

Origin and geographic range in BiH

Native, migratory, anadromous. Adriatic and Danube River Basins. Extinct (EX).

Remarks

Beluga used to occur in both the Danube River Basin and the Adriatic Sea (Vuković 1963a, 1977a). All recent references (Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009) are based on historical literature .

Literature

Anonymous 1886, 1909; Jurinac 1887; Glowacki 1896; Hirc 1896; Hawlitschek 1898; Medić 1901; Brusina 1902; Katuriće 1903; Ćurčić 1910; Plančić 1923b; Taler 1953a; Vuković 1963a; Tadić 1966; Sabioncello 1967; Sofradžija & Hadžiselimović 1981b; Jardas 1996; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Freyhof *et al.* 2020; Dekić *et al.* 2024.

RE4 *Huso naccarii* Bonaparte, 1836

Vernacular names

Jadranska jesetra. Adriatic sturgeon.

Previous names used for the species

Acipenser aculeatus Lovetsky, 1834; *Acipenser naccarii* Bonaparte, 1836; *Acipenser nardoi* Heckel, 1851.

Origin and geographic range in BiH

Endemic (ABE), migratory, anadromous. Adriatic Basin. Extinct (EX).

Remarks

In the past, the species was present in the Adriatic Sea and occasionally entered the Neretva River (Vuković 1963a, 1977a; Vuković & Ivanović 1971). Currently, populations in the entire distribution area are extremely small and there is no accurate data on its presence along the eastern coast of the Adriatic Sea. All recent literature data (Crivelli 1996; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011) are based on old reports.

Table 2 (continued on the next 4 pages). List of all lampreys and other fish species reported from freshwaters of Bosnia and Herzegovina with IUCN categories. Abbreviations: CR = Critically Endangered; DD = Data Deficient; EN = Endangered; FbIH = Federation of Bosnia and Herzegovina; LC = Least Concern; ME = marine euryhaline; NE = Not Evaluated; NT = Near Threatened; RS = Republic of Srpska; * = listed by Regulation on the Red List of protected species of flora and fauna of the Republic of Srpska; VU = Vulnerable.

Family	No.	Species	IUCN assessment	Status in BiH		Protection on international level	
				FBiH	RS	Habitat directive	Bern Convention
Petromyzontidae	1.	<i>Eudontomyzon vladkykovi</i>	LC	NE		Annex II	Appendix III
	2.	<i>Lampetra soljani</i>	NT	NE		Annex II, V	Appendix II, III
Acipenseridae	ME1	<i>Petromyzon marinus</i>	LC	DD		Annex II	Appendix III
	3.	<i>Huso ruthenus</i>	EN	VU	*	Annex V	Appendix III
Anguillidae	4.	<i>Anguilla anguilla</i>	CR	LC			
Clupeidae	ME2	<i>Alosa fallax</i>	NT	NE		Annex II, V	Appendix III
	5.	<i>Cobitis elongata</i>	LC	LC	*	Annex II	Appendix III
Cobitidae	6.	<i>Cobitis elongatoides</i>	LC	LC	*	Annex II	
	7.	<i>Cobitis herzegoviniensis</i>	EN	NE			
	8.	<i>Cobitis illyrica</i>	EN	NE			
	9.	<i>Cobitis narentana</i>	NT	VU			
	10.	<i>Misgurnus fossilis</i>	LC	LC	*	Annex II	Appendix III
	11.	<i>Sabanejewia balcanica</i>	LC	DD	*	Annex II	
Nemacheilidae	12.	<i>Sabanejewia bulgarica</i>	LC	NE			
	13.	<i>Barbatula barbatula</i>	LC	LC			
Cyprinidae	14.	<i>Aulopyge huegelii</i>	EN	EN		Annex II	
	15.	<i>Barbus balcanicus</i>	LC	LC	*	Annex II, V	Appendix III
	16.	<i>Barbus barbus</i>	LC	LC		Annex V	
	17.	<i>Carassius auratus</i>	LC	LC			
	18.	<i>Carassius carassius</i>	LC	LC	*		
	19.	<i>Carassius gibelio</i>	NE	LC			
	20.	<i>Carassius langsdorffi</i>	LC	NE			
	21.	<i>Cyprinus carpio</i>	NT	LC	*		
Xenocypridae	22.	<i>Ttenopharyngodon idella</i>	LC	DD			
	23.	<i>Hypophthalmichthys molitrix</i>	NT	LC			

Table 2 (continued). List of all lampreys and other fish species reported from freshwaters of Bosnia in Herzegovina with IUCN categories. Abbreviations: ME = marine euryhaline; FbiH = Federation of Bosnia and Herzegovina, RS = Republic of Srpska; * = listed at Regulation on the Red List of protected species of flora and fauna of the Republic of Srpska); CR = Critically Endangered; DD = Data Deficient; EN = Endangered; LC = Least Concern; NT = Near Threatened; NE = Not Evaluated; VU = Vulnerable.

Family	No.	Species	IUCN assessment	Status in BiH		Protection on international level	
				FbIH	RS	Habitat directive	Bern Convention
Tincidae	24.	<i>Hypophthalmichthys nobilis</i>	DD	LC			
	25.	<i>Tinca tinca</i>	LC	LC	*		
Acheilognathidae	26.	<i>Rhodeus amarus</i>	LC	LC	*	Annex II	Appendix III
	27.	<i>Gobio obtusirostris</i>	LC	LC	*		
Leuciscidae	28.	<i>Pseudorasbora parva</i>	LC	LC			
	29.	<i>Romanogobio carpathorossicus</i>	LC	EN	*	Annex II	Appendix III
	30.	<i>Romanogobio uranoscopus</i>	LC	EN	*	Annex II	Appendix III
	31.	<i>Romanogobio vladkyovi</i>	LC	NE		Annex II	Appendix III
	32.	<i>Abramis brama</i>	LC	LC			
	33.	<i>Alburnoides bipunctatus</i>	LC	LC	*		Appendix III
	34.	<i>Alburnus alburnus</i>	LC	LC			
	35.	<i>Alburnus neretvae</i>	NT	NE			
	36.	<i>Alburnus sava</i>	LC	NE	*	Annex II	Appendix III
	37.	<i>Ballerus ballerus</i>	LC	LC	*		Appendix III
38.	<i>Ballerus sapa</i>	LC	LC	*		Appendix III	
39.	<i>Blicca bjoerkna</i>	LC	LC				
40.	<i>Chondrostoma knerii</i>	EN	EN		Annex II	Appendix III	
41.	<i>Chondrostoma nasus</i>	NT	LC	*		Appendix III	
42.	<i>Chondrostoma phoxinus</i>	EN	CR		Annex II	Appendix III	
43.	<i>Delminichthys adspersus</i>	EN	EN			Appendix III	
44.	<i>Delminichthys ghetaldii</i>	VU	EN	*			
45.	<i>Leucaspis delineatus</i>	LC	VU	*		Appendix III	
46.	<i>Leuciscus aspius</i>	LC	LC	*	Annex II, V	Appendix III	
47.	<i>Leuciscus idus</i>	LC	LC	*		Appendix III	
48.	<i>Leuciscus leuciscus</i>	LC	LC				

Table 2 (continued). List of all lampreys and other fish species reported from freshwaters of Bosnia in Herzegovina with IUCN categories. Abbreviations: ME = marine euryhaline; FbiH = Federation of Bosnia and Herzegovina, RS = Republic of Srpska; * = listed at Regulation on the Red List of protected species of flora and fauna of the Republic of Srpska); CR = Critically Endangered; DD = Data Deficient; EN = Endangered; LC = Least Concern; NT = Near Threatened; NE = Not Evaluated; VU = Vulnerable.

Family	No.	Species	IUCN assessment	Status in BiH		Protection on international level	
				FbH	RS	Habitat directive	Bern Convention
	49.	<i>Leucos basak</i>	LC	EN		Annex II	
	50.	<i>Pelecus cultratus</i>	LC	LC	*	Annex II, V	Appendix III
	51.	<i>Phoxinellus alepidotus</i>	EN	EN		Annex II	
	52.	<i>Phoxinellus pseudalepidotus</i>	EN	NE		Annex II	
	53.	<i>Phoxinus karsticus</i>	NT	NE			
	54.	<i>Phoxinus lumaireul</i>	LC	LC			
	55.	<i>Rutilus rutilus</i>	LC	LC			
	56.	<i>Rutilus virgo</i>	LC	DD	*	Annex II, V	Appendix III
	57.	<i>Scardinius dergle</i>	NT	DD			
	58.	<i>Scardinius erythrophthalmus</i>	LC	LC			
	59.	<i>Scardinius plotizza</i>	NT	DD			
	60.	<i>Squalius cephalus</i>	LC	LC			
	61.	<i>Squalius microlepis</i>	EN	CR		Annex II	Appendix III
	62.	<i>Squalius squalus</i>	LC	LC	*		
	63.	<i>Squalius svallize</i>	NT	VU	*	Annex II	Appendix III
	64.	<i>Squalius tenellus</i>	EN	CR			Appendix III
	65.	<i>Telestes dabar</i>	EN	NE			
	66.	<i>Telestes metohiensis</i>	EN	CR	*		
	67.	<i>Telestes souffia</i>	LC	LC	*	Annex II	Appendix III
	68.	<i>Vimba vimba</i>	LC	LC	*		Appendix III
Siluridae	69.	<i>Silurus glanis</i>	LC	LC	*		Appendix III
Ictaluridae	70.	<i>Ameiurus melas</i>	LC	NE			
	71.	<i>Ameiurus nebulosus</i>	LC	LC			
Esocidae	72.	<i>Esox lucius</i>	LC	LC			
Umbridae	73.	<i>Umbrakrameri</i>	VU	EN	*	Annex II	Appendix II

Table 2 (continued). List of all lampreys and other fish species reported from freshwaters of Bosnia in Herzegovina with IUCN categories. Abbreviations: ME = marine euryhaline; FbiH = Federation of Bosnia and Herzegovina, RS = Republic of Srpska; * = listed at Regulation on the Red List of protected species of flora and fauna of the Republic of Srpska); CR = Critically Endangered; DD = Data Deficient; EN = Endangered; LC = Least Concern; NT = Near Threatened; NE = Not Evaluated; VU = Vulnerable.

Family	No.	Species	IUCN assessment	Status in BiH		Protection on international level		
				FbH	RS	Habitat directive	Bern Convention	
Salmonidae	74.	<i>Coregonus peled</i>	LC	NE		Annex V	Appendix III	
	75.	<i>Hucho hucho</i>	VU	EN	*	Annex II, V	Appendix III	
	76.	<i>Oncorhynchus mykiss</i>	LC	LC				
	77.	<i>Salmo farioides</i>	NT	NE				
	78.	<i>Salmo labrax</i>	LC	NE				
	79.	<i>Salmo letnica</i>	EN	NE				
	80.	<i>Salmo marmoratus</i>	VU	CR		Annex II		
	81.	<i>Salmo montenigrinus</i>	NE	NE				
	82.	<i>Salmo obtusirostris</i>	VU	CR	*	Annex II		
	83.	<i>Salmo trutta</i>	LC	LC	*			
	84.	<i>Salvelinus alpinus</i>	LC	LC				
	85.	<i>Salvelinus fontinalis</i>	LC	LC				
	86.	<i>Thymallus thymallus</i>	LC	LC	*	Annex V	Appendix III	
	87.	<i>Lota lota</i>	LC	LC	*			
	Syngnathidae	ME3	<i>Syngnathus abaster</i>	LC	DD			Appendix III
	Odontobutidae	88.	<i>Perccottus glenii</i>	LC	NE			
89.		<i>Knipowitschia radovici</i>	EN	NE				
Oxudercidae	90.	<i>Ninnigobius canestrinii</i>	LC	LC		Annex II	Appendix II	
	91.	<i>Orsinigobius croaticus</i>	VU	NE				
Gobiidae	92.	<i>Babka gymnotrachelus</i>	LC	NE				
	93.	<i>Neogobius fluviatilis</i>	LC	LC			Appendix III	
	94.	<i>Neogobius melanostomus</i>	LC	LC				
	95.	<i>Ponticola kessleri</i>	LC	DD			Appendix III	
Pleuronectidae	96.	<i>Proterorhinus semilunaris</i>	LC	LC				
	ME4	<i>Platichthys flesus</i>	LC	LC				

Table 2 (continued). List of all lampreys and other fish species reported from freshwaters of Bosnia in Herzegovina with IUCN categories. Abbreviations: ME = marine euryhaline; FbIH = Federation of Bosnia and Herzegovina, RS = Republic of Srpska; * = listed at Regulation on the Red List of protected species of flora and fauna of the Republic of Srpska); CR = Critically Endangered; DD = Data Deficient; EN = Endangered; LC = Least Concern; NT = Near Threatened; NE = Not Evaluated; VU = Vulnerable.

Family	No.	Species	IUCN assessment	Status in BiH		Protection on international level	
				FbIH	RS	Habitat directive	Bern Convention
Atherinidae	ME5	<i>Atherina boyeri</i>	LC	LC			
Poecilidae	97.	<i>Gambusia holbrooki</i>	LC	DD			
Mugilidae	ME6	<i>Chelon auratus</i>	LC	LC			
	ME7	<i>Chelon labrosus</i>	LC	LC			
	ME8	<i>Chelon ramada</i>	LC	LC			
	ME9	<i>Chelon saliens</i>	LC	LC			
	ME10	<i>Mugil cephalus</i>	LC	LC			
Bleniidae	98.	<i>Salaria fluviatilis</i>	LC	VU			Appendix III
Percidae	99.	<i>Gymnocephalus baloni</i>	LC	NE	*	Annex II, IV	Appendix III
	100.	<i>Gymnocephalus cernua</i>	LC	LC			
	101.	<i>Gymnocephalus schraetser</i>	LC	LC	*	Annex II, V	Appendix III
	102.	<i>Perca fluviatilis</i>	LC	LC			
	103.	<i>Sander lucioperca</i>	LC	LC			
	104.	<i>Sander volgensis</i>	LC	LC	*		Appendix III
	105.	<i>Zingel streber</i>	LC	VU	*	Annex II	Appendix III
	106.	<i>Zingel zingel</i>	LC	VU	*	Annex V	Appendix III
Gasterosteidae	107.	<i>Gasterosteus aculeatus</i>	LC	EN			
Cottidae	108.	<i>Cottus gobio</i>	LC	LC			Annex II
Centrarchidae	109.	<i>Lepomis gibbosus</i>	LC	NE			
	110.	<i>Micropterus salmoides</i>	LC	NE			
Moronidae	ME11	<i>Dicentrarchus labrax</i>	NT	DD			
Sparidae	ME12	<i>Sparus aurata</i>	LC	NE			

Literature

Vuković 1963a, 1977a; Vuković & Ivanović 1971; Crivelli 1996; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Freyhof *et al.* 2020; Dekić *et al.* 2024.

RE5 *Huso nudiventris* Lovetsky, 1828

Vernacular names

Sim. Ship sturgeon.

Previous names used for the species

Acipenser schypa Gldenstdt, 1772; *Acipenser nudiventris* Lovetsky, 1828; *Acipenser glaber* Fitzinger, 1832.

Origin and geographic range in BiH

Native, migratory, anadromous. Danube River Basin. Extinct (EX).

Remarks

In the past, the species inhabited the Danube River and its tributaries in former Yugoslavia, including BiH (Vuković 1963a, 1977a). However, today it is not present in the rivers of BiH and all recent references (Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009) are based on historical data.

Literature

Heckel & Kner 1858; Anonymous 1886; Brusina 1892; Glowacki 1896; Langhoffer 1904; Ćurčić 1910; Plančić 1923b; Tadić 1932; Vuković 1963a, 1977a; Sofradžija & Hadžiselimović 1981b; Sokolov & Vasil'ev 1989b; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Freyhof *et al.* 2020; Dekić *et al.* 2024.

3. *Huso ruthenus* (Linnaeus, 1758)

Vernacular names

Kečiga. Sterlet.

Previous names used for the species

Acipenser ruthenus Linnaeus, 1758; *Acipenser pygmaeus* Pallas, 1814; *Acipenser gmelini* Fitzinger, 1836; *Acipenser ruthenicus* Brusina, 1902.

Origin and geographic range in BiH

Native, migratory, potamodromous. Danube River Basin.

Remarks

Sterlet is the only species of sturgeon still present in the rivers of the Danube River Basin. Although its populations are small, it occurs in the Sava River and its tributaries (Hamzić 2024). Most literature references (Vuković 1963a, 1977a; Kosorić *et al.* 1980; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009) are based on catches by commercial and recreational fishermen.

Literature

Anonymous 1886, 1898; Ćurčić 1910; Plančić 1923a; Jedlička 1935; Taler 1948; Ristić 1963; Vuković 1963a, 1977a; Čanković 1964; Kiškarolj 1965; Čanković *et al.* 1968a, 1968b, 1968c; Munjko 1972; Kosorić *et al.* 1980; Bojčić *et al.* 1982; Obratil 1982; Veljović 1982, 1985; Delić 1984; Sokolov & Vasil'ev 1989a; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Suić *et al.* 2008; Sofradžija 2009; Freyhof 2012; Simonović *et al.* 2015; Kovačević *et al.* 2018; Freyhof *et al.* 2020; Skenderović *et al.* 2020b; Dekić *et al.* 2024.

RE6 *Huso stellatus* Pallas, 1771

Vernacular names

Pastruga. Stellate sturgeon.

Previous names used for the species

Acipenser stellatus Pallas, 1771; *Acipenser rostratus* Brandt & Ratzeburg, 1833.

Origin and geographic range in BiH

Native, migratory, anadromous. Danube River Basin. Extinct (EX).

Remarks

In BiH, its occurrence was reported in the Danube River and its tributaries (Vuković 1963a, 1977a). However, today it is not present in the rivers of BiH and all references for the region (Jardas 1996; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009) are based on historical data.

Literature

Anonymous 1886; Glowacki 1896; Medić 1896, 1898; Mojsisovics 1897; Horvat 1901; Plančić 1923b; Tadić 1931, 1932; Vuković 1963a, 1977a; Shubina *et al.* 1989; Jardas 1996; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Freyhof *et al.* 2020; Dekić *et al.* 2024.

Order Anguilliformes Goodrich, 1909

Family Anguillidae Rafinesque, 1810

Genus *Anguilla* Garsault, 1764

4. *Anguilla anguilla* (Linnaeus, 1758)

Vernacular names

Jegulja. European eel.

Previous names used for the species

Anguilla vulgaris Shaw, 1803; *Anguilla fluviatilis* Heckel & Kner, 1858.

Origin and geographic range in BiH

Native migratory, catadromous. Adriatic and Danube Basin.

Remarks

Native to the Adriatic Basin, in 2004 it was found in Sava River (Danube River Basin) (Golub, unpublished data).

Literature

Horvat 1901; Anonymous 1914, 1965; Aganović 1952a; Vuković 1963a, 1977a; Kosorić & Vuković 1966a; Kačanski *et al.* 1977; Gvozdenović *et al.* 1978; Kosorić 1978; Veljović 1982; Kosorić *et al.* 1983, 1989; Jardas 1996; Simonović 2001; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Tutman *et al.* 2007, 2012a; Glamuzina *et al.* 2008a, 2008b; Has-Schön *et al.* 2008; Muhamedagić *et al.* 2008; Šanda *et al.* 2008a, 2008b, 2009; Sofradžija 2009; Pavličević *et al.* 2016; Freyhof *et al.* 2020; Dekić *et al.* 2024.

Order Clupeiformes Goodrich, 1909
Family Clupeidae Cuvier, 1817
Genus *Alosa* Garsault, 1764

ME2 *Alosa fallax* (Lacepède, 1803)

Vernacular names

Čepa. Twaite shad.

Previous names used for the species

Clupea fallax Lacepède, 1803; *Alosa fallax nilotica* Geoffroy, 1827.

Origin and geographic range in BiH

Native, migratory, anadromous. Adriatic Basin.

Remarks

Alosa fallax is a migratory species that seasonally enters the Neretva River and its downstream tributaries about 45–50 km to the confluence with the River Buna (locality Bunski Kanali). Every year the species enters the Krupa River (connecting Hutovo Blato wetland with the Neretva River) for spawning in April and May.

Literature

Karaman 1928; Vuković 1961, 1962, 1963a, 1977a; Vuković & Ivanović 1971; Kosorić 1978; Kosorić *et al.* 1983, 1989; Jardas 1996; Mikavica 1998; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Muhamedagić *et al.* 2008; Sofradžija 2009; Drešković *et al.* 2011; Tutman *et al.* 2012a.

RE7 *Alosa immaculata* Bennet, 1835

Vernacular names

Crnomorska haringa. Pontic shad.

Previous names used for the species

Alosa pontica (Eichwald, 1838); *Alosa caspia* (Eichwald, 1838); *Alosa caspia nordmanni* (Antipa, 1906).

Origin and geographic range in BiH

Native, migratory, anadromous. Danube River Basin. Extinct (EX).

Remarks

Although considered as regionally extinct (Ćaleta *et al.* 2019), the western population of *Alosa immaculata* migrates from the Black Sea into the Danube River for spawning, with fish occasionally found near Đerdap in Serbia and Romania (Višnjić *et al.* 2010). Despite that, all recent references for

BiH are based on historical data, and therefore require confirmation. Sofradžija (2009) listed *A. pontica* (Eichwald, 1838), but this is a synonym of *A. immaculata*.

Literature

Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Milanović *et al.* 2015; Lukić *et al.* 2016; Dekić *et al.* 2024.

Order Cypriniformes Goodrich, 1909

Family **Cobitidae** Swainson, 1838

Tutman *et al.* (2017b) critically assessed the current taxonomic status of this family and its geographic distribution. In the Danube River Basin, four species (*Cobitis elongata* Heckel & Kner, 1858; *C. elongatoides* Băcescu & Mayer, 1969; *Misgurnus fossilis* (Linnaeus, 1758), *Sabanejewia balcanica* (Karaman, 1922)) have been recorded, while in the Adriatic Basin three species (*C. narentana* Karaman, 1928, *C. illyrica* Freyhof & Stelbrink, 2007, *C. herzegoviniensis* (Buj & Šanda, 2014)) were recorded. *Cobitis elongatoides* is the most common and widest distributed across northern BiH, i.e., in the Sava, Drina, Bosna, Vrbas, and Una rivers (Danube River Basin). Cobitid species from the Adriatic Basin, in turn, are endemic, each with a very small distribution range. Based on available data, it is clear that reports of *C. taenia* Linnaeus, 1758 in BiH are the result of a misidentification, because its distribution range is restricted to northern Europe (Nalbant *ets al.* 2001). Therefore, it is not included in the BiH species checklist.

Genus *Cobitis* Linnaeus, 1758

Cobitis dalmatina Karaman, 1928

Remarks

Listed in Hamzić (2024) with the remark that its presence is yet to be confirmed. Considering that the species occurs only in the Cetina River catchment in Croatia and is not recorded in BiH (Buj *et al.* 2014, 2015a; Čaleta *et al.* 2019), it is not included in the BiH species checklist.

5. ***Cobitis elongata*** Heckel & Kner, 1858

Vernacular names

Veliki vijun. Balkan spined loach.

Previous names used for the species

Cobitis taenia elongata Heckel & Kner, 1858.

Origin and geographic range in BiH

Endemic (DBE). Danube River Basin.

Literature

Brusina 1892; Medić 1901; Bănărescu & Nalbant 1957, 1998; Mrakovčić *et al.* 2000, 2002, 2006; Bogut *et al.* 2006; Adrović 2007; Kottelat & Freyhof 2007; Buj *et al.* 2008; Kopjar *et al.* 2008; Mrakovčić *et al.* 2008; Pavlović *et al.* 2009; Sofradžija 2009; Popović 2010; Drešković *et al.* 2011; Adrović 2012; Adrović *et al.* 2012; Jakovlić *et al.* 2013; Škrijelj *et al.* 2013; Čičić-Močić 2014, 2016, 2018; Delić *et al.* 2014; Pavličević *et al.* 2014; Hamzić *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Lukić *et al.* 2016; Tutman *et al.* 2017b; Kovačević *et al.* 2018; Memić 2018.

6. *Cobitis elongatoides* Băcescu & Mayer, 1969

Vernacular names

Vijun, badelj. Danube spined loach.

Previous names used for the species

Cobitis taenia var. *elongatoides* Băcescu, 1962; *Cobitis taenia elongatoides* Băcescu & Mayer, 1969; *Cobitis taenia danubialis* Nalbant, 1993.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin (Hamzić, unpublished data).

Remarks

In the older literature this species was mentioned as *C. taenia* (Vuković 1963a, 1977a; Vuković & Ivanović 1971; Sofradžija & Berberović 1978; Kosorić 1981; Kosorić *et al.* 1983; Korjenić 2004b) for the Sava River and its tributaries. Under the name of *C. taenia danubialis* it was mentioned by Mrakovčić *et al.* (2000) and Schneider *et al.* (2000a), while in Delić *et al.* (2003), Mrakovčić *et al.* (2008) and Dumbović *et al.* (2009) it is identified as *C. elongatoides*.

Literature

Anonymous 1886; Glowacki 1896; Ćurčić 1910; Zaplata & Taler 1932; Vuković 1963a, 1977a; Sofradžija & Berberović 1978; Kosorić *et al.* 1980; Habeković *et al.* 1981; Kosorić 1981; Veljović 1982; Sofradžija *et al.* 1984; Mikavica 1987b; Mikavica *et al.* 1991; Babić & Mikavica 1994; Mikavica & Savić 1999; Mrakovčić *et al.* 2000, 2002, 2006, 2008; Korjenić 2004b; Bogut *et al.* 2006; Skenderović *et al.* 2006, 2020a; Kottelat & Freyhof 2007; Ovčina 2007; Adrović *et al.* 2008, 2012; Vuković *et al.* 2008; Sofradžija 2009; Zrnčić *et al.* 2009; Memić & Adrović 2010; Golub *et al.* 2012; Čičić-Močić 2014, 2019; Delić *et al.* 2014; Nedić *et al.* 2014a, 2014c, 2014d; Hamzić *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Golub *et al.* 2016b, 2017c, 2018a, 2018b; Škrijelj *et al.* 2013; Dekić *et al.* 2017b, 2020a, 2020b, 2024; Tutman *et al.* 2017b; Keško 2018; Kovačević *et al.* 2018; Memić 2018, 2020; Lolić *et al.* 2019; Cvijić *et al.* 2020.

7. *Cobitis herzegoviniensis* (Buj & Šanda, 2014)

Vernacular names

Hercegovački vijun. Herzegovinian spined loach.

Origin and geographic range in BiH

Endemic (NCE, BiH). Adriatic Basin.

Remarks

The species, tentatively classified as *Cobitis* sp., was firstly reported by Šanda *et al.* (2008a) in the tributary of the Neretva River (Lištica) and surrounding karstic field (Mostarsko Blato) from where spined loaches were not previously reported (Kosorić & Vuković 1966a; Aganović 1969; Kosorić 1974, 1977a, 1978; Kosorić *et al.* 1989). Later taxonomic analysis revealed that the population distributed in the Mostarsko Blato karstic field should be considered as a distinct species that has not been described to date (Buj *et al.* 2014). The population is critically endangered by the construction of the Mostarsko Blato hydropower plant, where a large part of the habitat was irreversibly destroyed (dried out), and non-native, invasive and predatory species (pumpkinseed, brown bullhead, and rainbow trout) were introduced into the habitat.

Literature

Šanda *et al.* 2008a, 2008b, 2009; Glamuzina *et al.* 2013; Buj *et al.* 2014, 2015a, 2015b; Tutman *et al.* 2017b; Freyhof *et al.* 2020.

8. *Cobitis illyrica* Freyhof & Stelbrink, 2007

Vernacular names

Ilirski vijun. Illyric spined loach.

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

Another relatively recently described species with a disjunctive distribution, endemic to both Croatia and BiH in the Neretva River catchment. It was first recorded from Prološko Blato lake, in the Imotsko Polje field in Croatia (Freyhof & Stelbrink 2007), while Šanda *et al.* (2008b) noted the presence of a possibly undescribed species in Lake Krenica, Bekijsko Polje field. Later, Buj *et al.* (2014) confirmed both findings, as the first record of *C. illyrica* in BiH. This species inhabits Prološko Blato lake, Baćinska Lakes and the Matica River in neighbouring Croatia, as well as Krenica in the Bekijsko Polje field in BiH.

Literature

Freyhof & Stelbrink 2007; Šanda *et al.* 2008a, 2008b; Glamuzina *et al.* 2013; Buj *et al.* 2014, 2015a, 2015b; Darwall *et al.* 2014; Geiger *et al.* 2014; Čaleta *et al.* 2015; Tutman *et al.* 2017b; Freyhof *et al.* 2020.

9. *Cobitis narentana* Karaman 1928

Vernacular names

Neretvanski vijun. Neretva spined loach.

Previous names used for the species

Cobitis taenia narentana Karaman, 1928.

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

Endemic species distributed in the lower reaches of the Neretva River, its tributaries and channels and Lake Modro Oko in Croatia (Schneider *et al.* 2000b; Zanella *et al.* 2003; Buj *et al.* 2014, 2015a), as well as in the Trebišnjica River and Hutovo Blato wetland in BiH (Tutman *et al.* 2006, 2012a; Buj *et al.* 2014, 2015a, 2015b). It was first described as a subspecies (*C. taenia narentana* Karaman, 1928); later data include only general notes from the Neretva River catchment (Vuković 1963a, 1977a). Schneider *et al.* (2000b) redescribed it as a distinct species, *C. narentana*. Sofradžija (2009) considered the lower Neretva River from the town Čapljina as its distribution area in BiH, while Tutman *et al.* (2006, 2012a), Šanda *et al.* (2008a, 2008b), and Buj *et al.* (2015a, 2015b) found the species on the Hutovo Blato wetland area, in the Bregava and the Trebišnjica River, all in BiH.

Literature

Karaman 1928; Vuković 1963a, 1977a; Kottelat 1997; Mikavica 1998; Mrakovčić *et al.* 2000, 2006, 2008; Schneider *et al.* 2000b; Bogut *et al.* 2006; Tutman *et al.* 2006, 2012a, 2017b; Buj *et al.* 2007, 2014, 2015a, 2015b; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008a, 2008b; Zupančič 2008; Dulčić *et al.* 2009; Sofradžija 2009; Zanella *et al.* 2009; Glamuzina *et al.* 2008a, 2013; Buj *et al.* 2014, 2015a, 2015b; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Čaleta *et al.* 2015; Milanović *et al.* 2015; Dekić *et al.* 2024.

Genus *Misgurnus* Lacepède, 1803

10. *Misgurnus fossilis* (Linnaeus, 1758)

Vernacular names

Čikov. Weather loach.

Previous names used for the species

Cobitis fossilis Linnaeus, 1758.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Historical data on its distribution in BiH are very scarce (Vuković 1963a, 1977a; Bogut *et al.* 2006); therefore, its actual distribution is dubious. According to the available data, it occurs in the Sava River with its tributaries of northern BiH (Sofradžija 2009). A report from the Adriatic Basin (Trebišnjica River catchment) indicates that it may have been translocated from the Danube River Basin (Šukalo *et al.* 2018). Therefore, additional research is necessary.

Literature

Anonymous 1886, 1909; Brusina 1892; Glowacki 1896; Ćurčić 1910; Vuković 1963a, 1977a; Veljović 1982; Mikavica *et al.* 1991; Mikavica & Savić 1999; Mrakovčić *et al.* 2000, 2006, 2008; Radević 2000; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Vuković *et al.* 2008; Sofradžija 2009; Drešković *et al.* 2011; Jakovlić *et al.* 2013; Muhamedagić *et al.* 2013; Hamzić *et al.* 2015; Simonović *et al.* 2015; Lukić *et al.* 2016; Tutman *et al.* 2017b; Dekić *et al.* 2024.

Genus *Sabanejewia* Vladykov, 1929

11. *Sabanejewia balcanica* (Karaman 1922)

Vernacular names

Balkanski zlatni vijun. Balkan golden loach.

Previous names used for the species

Cobitis aurata (Filippi, 1865); *Cobitis aurata balcanica* Karaman 1922; *Cobitis aurata bosniaca* Karaman, 1963; *Sabanejewia aurata balcanica* (Karaman, 1922).

Origin and geographic range in BiH

Native. Danube River Basin.

Remarks

Sabanejewia balcanica was previously considered as a subspecies of *Cobitis aurata*, *C. aurata balcanica* Karaman, 1922, reported from waters of the Danube River Basin, whereas *C. aurata bosniaca* Karaman, 1963 was restricted to the Vrbas River and its tributaries Suturlija and Široka (Sava River catchment) (Karaman 1963; Vuković 1963a, 1977a; Sofradžija 2009). Later authors (Mrakovčić *et al.* 2000, 2006, 2008; Bogut *et al.* 2006; Sofradžija 2009; Drešković *et al.* 2011) simply followed previous reports without new, distributional data.

Literature

Karaman 1922, 1963; Zaplata & Taler 1932; Vuković 1963a, 1977a, Kottelat 1997; Mikavica 1998; Mrakovčić *et al.* 2000, 2006, 2008; Radević 2000; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Marešová *et al.* 2011; Adrović *et al.* 2012; Muhamedagić *et al.* 2013; Delić *et al.* 2014; Hamzić *et al.* 2015; Milanović *et al.* 2015; Simonović *et al.* 2015; Čičić-Močić 2016; Golub *et al.* 2016b; Tutman *et al.* 2017b; Adrović 2018; Memić 2018, 2020; Bajrić *et al.* 2018a, 2018b, 2021a, 2021b; Skenderović *et al.* 2020a; Dekić *et al.* 2024.

12. *Sabanejewia bulgarica* (Drensky, 1928)

Vernacular names

Dunavski zlatni vijun. Bulgarian golden loach.

Origin and geographic range in BiH

Endemic (DBE). Danube River Basin.

Remarks

Fedorčák *et al.* (2023) recently confirmed the presence of *Sabanejewia bulgarica* in several localities in BiH, in habitats with specific ecological characteristics.

Literature

Fedorčák *et al.* 2023.

Family Nemacheilidae Regan, 1911

Genus *Barbatula* Linck, 1790

13. *Barbatula barbatula* (Linnaeus, 1758)

Vernacular names

Brkica. Stone loach.

Previous names used for the species

Barbatula barbatulus (Linnaeus, 1758); *Cobitis barbatula* Linnaeus, 1758; *Nemacheilus barbatulus barbatulus* Linnaeus, 1758; *Nemachilus barbatulus* (Linnaeus, 1758); *Orthrias barbatulus barbatulus* (Linnaeus, 1758).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

First mentioned as *Cobitis barbatula* by Zaplata & Taler (1932) in the Bosna River, later as *Nemachilus barbatulus* (Vuković 1963a, 1977a) and *N. barbatulus* (Sofradžija 2009) in the small rivers of the Sava River watershed. Reported as *Barbatula barbatula* (Bogut *et al.* 2006) in the waters of the Danube River Basin without precise locations. The record in the upper Neretva River, Adriatic Basin, by Vegara *et al.* (2009) seems to be the result of accidental introduction during stocking of other fishes (Hamzić 2024). Calegari *et al.* (2025) designated neotype for *B. barbatula* and restricted the distribution of this species to a small region in the Mediterranean France, while many populations, including those from BiH, are currently of unknown taxonomic status, and will require further careful evaluation.

Literature

Heckel & Kner 1858; Anonymous 1886; Glowacki 1896; Zaplata & Taler 1932; Simić 1963; Vuković 1963a, 1977a; Sofradžija & Vuković 1979; Sofradžija *et al.* 1984; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Vegara *et al.* 2009; Golub *et al.* 2012, 2017c, 2018a, 2018b; Jakovlić *et al.* 2013; Čičić-Močić 2014, 2016, 2018; Nedić *et al.* 2014a, 2014c, 2014d; Hamzić *et al.* 2015; Simonović *et al.* 2015; Dekić *et al.* 2017b, 2020b; Tutman *et al.* 2017b; Adrović 2018; Memić 2018; Lolić *et al.* 2019; Neuburg *et al.* 2023; Calegari *et al.* (2025).

Family Cyprinidae Rafinesque, 1815

Genus *Aulopyge* Heckel, 1841

14. *Aulopyge huegelii* Heckel, 1843

Vernacular names

Oštrulj. Dalmatian barbel gudgeon.

Previous names used for the species

Aulopyge hugeli Heckel, 1841; *Aulopyge hügelii* Heckel, 1843.

Origin and geographic range in BiH

Endemic (ABE). Native to the Adriatic Basin, translocated to the Danube River Basin.

Remarks

Found in BiH and Croatia, where it inhabits the karst watercourses; occurs in some creeks in the Livanjsko Polje field, in the Cetina and Krka Rivers in Croatia, and was successfully introduced to Lake Blidinje (Ćurčić 1916), Buško Blato (Žujo-Zekić 2021) and Mostarsko Blato (Šanda *et al.* 2009) in the Adriatic Basin and Šator Lake (Delić *et al.* 2005) in the Danube River Basin (Una River catchment).

Literature

Heckel & Kner 1858; Ćurčić 1916; Karaman 1923, 1928; Protić 1924; Taler 1951a, 1951b, 1953a; Vuković 1963a, 1964, 1977a, 1982a; Aganović & Vuković 1966, 1967, 1968, 1971; Kapetanović *et al.* 1966; Pipić 1966; Berberović 1967; Vuković & žnidaršić-Krzyk 1969; Aganović 1969; Berberović *et al.* 1969c, 1971a; Guzina *et al.* 1971; Kaluđerčić *et al.* 1971; Miladinović *et al.* 1971; Vuković & Ivanović 1971; Winterhalter *et al.* 1971; Bozja & Švob 1974; Đurović & Vuković 1975; Korić 1975; Guzina & Vuković 1977, 1987; Kačanski *et al.* 1978; Mijatović *et al.* 1982; Kekić *et al.* 1983; Jojić 1984; Volić 1984; Gvozdenović *et al.* 1985; Guzina *et al.* 1986; Howes 1987; Pešić 1987; Vuković & Sofradžija 1987; Sofradžija & Kundurović 1988; Ivanc *et al.* 1989; Povž *et al.* 1990; Economidis & Bănărescu 1991; Crivelli 1996; Freyhof 1997; Mikavica 1998; Šorić & Bănărescu 1999;

Delić *et al.* 2005; Guzina 2000, 2005; Bogut *et al.* 2006; Freyhof *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Muhamedagić *et al.* 2008; Nuhanović 1989; Šanda *et al.* 2008a, 2008b, 2009; Zupančič 2008; Sofradžija 2009; Čaleta *et al.* 2009a; Hasković *et al.* 2010; Drešković *et al.* 2011; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Milanović *et al.* 2015; Mušović 2016; Dekić *et al.* 2016a, 2024; Benovics *et al.* 2017, 2019; Mustafić & Mrakovčić 2017; Mušović *et al.* 2018, 2020; Kalamujić Stroil *et al.* 2019; Freyhof *et al.* 2020; Buj *et al.* 2021; Ludoški *et al.* 2021; Žujo-Zekić 2021; Avdić *et al.* 2023; Lukač *et al.* 2023; Carosi *et al.* 2024.

Genus *Barbus* Cuvier & Cloquet, 1816

15. *Barbus balcanicus* Kotlík, Tsigenopoulos, Ráb & Berrebi, 2002

Vernacular names

Potočna mrena. Danube barbel.

Previous names used for the species

Barbus meridionalis petenyi Risso, 1826; *Barbus meridionalis* Risso, 1827; *Barbus caninus* (Bonaparte, 1839); *Barbus peloponnesius* Valenciennes, 1842; *Barbus peloponnesius petenyi* Heckel, 1852; *Barbus petenyi* Heckel, 1852.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to the Adriatic Basin (Drežanka River), as bait fish for pikeperch angling, also recorded in the Salakovac reservoir (Spasojević, unpublished data). All references of occurrences of *Barbus meridionalis petenyi* Heckel, 1852 (Vuković 1963a; Sofradžija 2009), *B. meridionalis* Riso, 1827 and *B. petenyi* Heckel, 1852 are erroneous since those species do not occur in BiH (Kottelat & Freyhof 2007).

Literature

Glowacki 1896; Karaman 1924; Zaplata & Taler 1932; Smlatić 1961; Vuković 1963a, 1963c, 1966a, 1977a; Šenk 1963; Šenk & Telalbašić 1963; Berberović 1967; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968, 1969; Šenk 1969; Aganović & Kapetanović 1973; Sofradžija & Berberović 1973; Kosorić *et al.* 1975, 1980; Berberović 1980; Omeragić *et al.* 1980; Kosorić 1981; Šolaja & Pocrnjić 1983a; Sofradžija *et al.* 1984; Mikavica 1987b, 1988a; Šorić & Janković 1989; Mikavica *et al.* 1991; Babić & Mikavica 1994; Proha 1997; Mikavica & Savić 1999; Radević 2000, 2002; Machordom & Doadrio 2001; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Skenderović *et al.* 2006, 2020a, 2020b; Škrijelj *et al.* 2006; Adrović 2007; Kottelat & Freyhof 2007; Skenderović & Adrović 2008b; Dekić *et al.* 2009a, 2013c, 2020a, 2020b, 2024; Kaltak 2009; Sofradžija 2009; Marková *et al.* 2010; Drešković *et al.* 2011; Ivanc *et al.* 2011; Adrović 2012, 2018; Golub *et al.* 2012, 2016a, 2017c, 2018a, 2018b; Ivanc & Dekić 2012; Čičić-Močić 2014, 2016, 2018, 2019; Delić *et al.* 2014; Milanović *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Adrović *et al.* 2018; Keško 2018; Kovačević *et al.* 2018; Lolić *et al.* 2019; Memić 2018, 2020; Cvijić *et al.* 2020.

16. *Barbus barbuis* (Linnaeus, 1758)

Vernacular names

Mrena. Barbel.

Previous names used for the species

Cyprinus barbuis Linnaeus, 1758; *Barbus barbuis barbuis* (Linnaeus, 1758); *Barbus vulgaris* Fleming, 1828; *Barbus fluviatilis* Fitzinger, 1832.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Ćurčić 1910; Plančić 1923b; Zaplata & Taler 1932; Bek 1935; Aganović & Milošević 1959; Vuković 1963a, 1977a; Čanković 1967; Čanković *et al.* 1968a, 1968b, 1968c; Kiškarolj & Čanković 1968, 1969; Šenk & Aganović 1968; Šenk 1969; Aganović & Kapetanović 1973; Kosorić *et al.* 1975, 1980; Kosorić 1981; Habeković *et al.* 1981; Kaćanski *et al.* 1981; Obratil 1982; Veljović 1982, 1985; Delić 1984, Mikavica 1987b, 1988a; Mikavica *et al.* 1991, 1997; Babić & Mikavica 1994; Mikavica & Savić 1997, 1999; Radević 2000; Machordom & Doadrio 2001; Mrakovčić *et al.* 2002; Bogut *et al.* 2006; Korjenić *et al.* 2006; Škrijelj *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Ivanc & Dekić 2012; Muhamedagić & Habibović 2013; Čičić-Močić 2014, 2018, 2019; Delić *et al.* 2014; Simonović *et al.* 2015; Bećiraj *et al.* 2016; Golub *et al.* 2016a, 2017c, 2018a, 2018b; Nedić *et al.* 2016a, 2018; Kovačević *et al.* 2018; Lolić *et al.* 2019; Skenderović & Adrović 2019; Zuliani *et al.* 2019; Cvijić *et al.* 2020; Dekić *et al.* 2020a, 2024; Skenderović *et al.* 2020a, 2020b, 2021.

Barbus plebejus Bonaparte, 1839

Previous names used for the species

Barbus plebejus plebejus Bonaparte, 1839; *Barbus plebeius* Valenciennes, 1842.

Remarks

Listed in Vuković & Ivanović (1971) and Drešković *et al.* (2011), but as a result of taxonomic misidentification of specimens since the southernmost finding of the species in the Adriatic Basin is the Krka River in Croatia (Ćaleta *et al.* 2019). Therefore, it is not included in the checklist.

Literature

Vuković & Ivanović (1971); Drešković *et al.* (2011).

Carassius Jarocki, 1822

17. *Carassius auratus* (Linnaeus, 1758)

Vernacular names

Zlatni karaš. Goldfish.

Previous names used for the species

Carassius auratus auratus (Linnaeus, 1758).

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Non-native species from Asia. It is possible that some records of *Carassius auratus* result from misidentifications of specimens of *C. gibelio*.

Literature

Vuković 1963a, 1977a; Vuković & Ivanović 1971; Kosorić *et al.* 1980; Bogut *et al.* 2006; Adrović & Skenderović 2007; Glamuzina *et al.* 2008a; Sofradžija 2009; Mrdak & Milošević 2017.

18. *Carassius carassius* (Linnaeus, 1758)

Vernacular names

Karaš. Crucian carp.

Previous names used for the species

Cyprinus carassius Linnaeus, 1758; *Carassius vulgaris* Nordmann, 1840.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to the Adriatic Basin (Hamzić, unpublished data).

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Ćurčić 1910; Plančić 1923b; Aganović 1957, 1967; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Žitnan *et al.* 1969; Vuković & Ivanović 1971; Pikula 1977; Sofradžija *et al.* 1977, 1978a; Berberović 1980; Guzina *et al.* 1980; Nadaždin *et al.* 1980; Vuković *et al.* 1980; Obratil 1982; Veljović 1982, 1985; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Adrović 2007; Kottelat & Freyhof 2007; Sofradžija 2009; Adrović 2012; Čičić-Močić 2014; Simonović *et al.* 2015; Skenderović 2015; Kovačević *et al.* 2018; Memić 2018; Skenderović *et al.* 2020b; Dekić *et al.* 2024.

19. *Carassius gibelio* (Bloch, 1782)

Vernacular names

Babuška. Prussian carp.

Previous names used for the species

Carassius auratus gibelio (Bloch, 1783).

Origin and geographic range in BiH

Non-native. Danube and Adriatic Basin.

Remarks

Non-native species that originally occurs in Asia. It is a widespread species present in both Danube and Adriatic basins with a high impact on native species of fish (Glamuzina *et al.* 2017; Hamzić 2024).

Literature

Vuković 1963a, 1977a; Plančić 1967; Čanković *et al.* 1968a; Žitnan *et al.* 1969; Vuković & Ivanović 1971; Pikula 1977; Sofradžija *et al.* 1977, 1978a; Kosorić 1978; Vuković *et al.* 1979, 1980; Nadaždin *et al.* 1980; Delić 1984, 1985; Škrijelj 1991; Safin 1997; Leiner 1998; Radević 2000, 2002; Mrakovčić *et al.* 2002; Škrijelj *et al.* 2005a, 2005b; Bogut *et al.* 2006; Adrović 2007; Kottelat & Freyhof 2007; Has-Schön *et al.* 2008; Vuković *et al.* 2008; Muhamedagić *et al.* 2008; Šanda *et al.* 2008a, 2008b; Sofradžija 2009; Skenderović *et al.* 2010, 2011a, 2011b, 2020b, 2021; Ivanc *et al.* 2011; Adrović 2012, 2018; Tutman *et al.* 2012a; Dekić *et al.* 2013d, 2020a, 2024; Golub *et al.* 2013, 2017b, 2018a, 2018b; Mikavica *et al.* 2013; Rylková *et al.* 2013; Čičić-Močić 2014, 2016; Nedić *et al.* 2014a, 2014c, 2014d, 2016b, 2018; Riđanović *et al.* 2015; Hajdarević *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Čulibrk *et al.* 2016; Pavličević *et al.* 2016; Glamuzina *et al.* 2017; Kovačević *et al.* 2018; Memić 2018; Mušović *et al.* 2018, 2020; Žujo-Zekić 2021; Žujo-Zekić *et al.* 2018; Skenderović & Adrović 2019; Adrović *et al.* 2021; Halilović & Dekić 2022; Carosi *et al.* 2024.

20. *Carassius langsdorfi* Temminck & Schlegel, 1846

Vernacular names

Japanski karaš. Japanese silver crucian carp.

Origin and geographic range in BiH

Non-native. Adriatic Basin.

Remarks

Non-native species that originally occurs in Asia; several specimens were recorded in the Neretva River catchment based on molecular data (Kalous *et al.* 2012).

Literature

Kalous *et al.* 2012; Rylková *et al.* 2013; Piria *et al.* 2018.

Cyprinus Linnaeus, 1758

21. *Cyprinus carpio* Linnaeus, 1758

Vernacular names

Šaran. Common Carp.

Previous names used for the species

Cyprinus carpio carpio Linnaeus, 1758.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to several lakes and reservoirs in the Adriatic Basin (Glamuzina *et al.* 2017), where it is regularly stocked (Spasojević, unpublished data).

Literature

Anonymous 1886, 1909, 1947a, 1947b, 1948, 1959; Brusina 1892; Glowacki 1896; Ćurčić 1910; Plančić 1923b; Trojanović 1934; Jedlička 1935; Taler 1951b; Aganović 1952a, 1957, 1967; Vuković 1963a, 1977a; Kosorić 1965, 1978; Čanković *et al.* 1968a, 1968b, 1968c, 1976; Aganović & Kapetanović 1973; Kačanski *et al.* 1977; Vuković & Kosorić 1978; Berberović 1980; Grbelja *et al.* 1980; Kosorić *et al.* 1980, 1983, 1989; Nadaždin *et al.* 1980; Vuković *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Veljović 1982; Delić 1984; Ratković & Mikavica 1984; Mikavica *et al.* 1991; Pažur 1993; Leiner 1998; Mikavica & Savić 1999; Radević 2000, 2002; Korjenić *et al.* 2003; Mikavica & Vuković 2003; Škrijelj *et al.* 2003, 2005a, 2005b; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Adrović 2007; Glamuzina *et al.* 2008a, 2017; Kottelat & Freyhof 2007; Has-Schön *et al.* 2008; Muhamedagić *et al.* 2008; Suić *et al.* 2008; Šanda *et al.* 2008a, 2008b, 2009; Vuković *et al.* 2008; Sofradžija 2009; Skenderović *et al.* 2010, 2011b, 2015, 2020b, 2021; Ivanc *et al.* 2011; Tomljanović *et al.* 2011, 2013; Adrović 2012; Freyhof 2012; Skenderović & Adrović 2012, 2016; Tutman *et al.* 2012a; Golub *et al.* 2013, 2017c; Čičić-Močić 2014; Delić *et al.* 2014; Nedić *et al.* 2014c, 2016a, 2016b, 2018; Riđanović *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Pavličević *et al.* 2016; Piria 2016b; Džafić *et al.* 2018; Kovačević *et al.* 2018; Mušović *et al.* 2018, 2020; Piria *et al.* 2018; Žujo-Zekić *et al.* 2018; Žujo-Zekić 2021; Adrović *et al.* 2021; Dekić *et al.* 2024.

Family Xenocyprididae Günther, 1868

Ctenopharyngodon Steindachner, 1866

22. *Ctenopharyngodon idella* (Valenciennes, 1844)

Vernacular names

Bijeli amur. Grass carp.

Previous names used for the species

Ctenopharingodon idella (Valenciennes, 1844); *Ctenopharingodon idellus* (Valenciennes, 1844),.

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Non-native species that originally occurs in Asia, introduced for biomanipulation and control of algae and aquatic plants.

Literature

Vuković 1963a, 1977a; Žitnan *et al.* 1969; Veljović 1982; Radević 2000; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Muhamedagić *et al.* 2008; Sofradžija 2009; Tutman *et al.* 2012a; Nedić *et al.* 2014a, 2014c, 2014d, 2016b; Riđanović *et al.* 2015; Glamuzina *et al.* 2017; Golub *et al.* 2017c; Kovačević *et al.* 2018; Skenderović *et al.* 2020b; Dekić *et al.* 2024.

Hypophthalmichthys Bleeker, 1860

23. *Hypophthalmichthys molitrix* (Valenciennes, 1844)

Vernacular names

Bijeli glavaš, bijeli tolstobik. Silver carp.

Previous names used for the species

Aristichthys molitrix (Richardson, 1845).

Origin and geographic range in BiH

Non-native. Danube and the Adriatic Basin.

Remarks

Non-native species that originally occurs in Asia, introduced for biomanipulation and control of planktonic organisms.

Literature

Vuković 1963a, 1977a; Radević 2000; Bogut *et al.* 2006; Sofradžija 2009; Tutman *et al.* 2012a; Simonović *et al.* 2015; Glamuzina *et al.* 2017; Mušović *et al.* 2020; Žujo-Zekić 2021; Dekić *et al.* 2024.

24. *Hypophthalmichthys nobilis* (Richardson, 1845)

Vernacular names

Šareni glavaš, šareni tolstobik. Bighead carp.

Origin and geographic range in BiH

Non-native. Danube and the Adriatic Basin.

Previous names used for the species

Aristichthys nobilis (Richardson, 1845).

Remarks

Non-native species from Asia introduced for biomanipulation and control of planktonic organisms.

Literature

Vuković 1963a, 1977a; Radević 2000; Sofradžija 2009; Glamuzina *et al.* 2017; Dekić *et al.* 2024.

Family Tincidae Jordan, 1878

3.4.5.1 Genus *Tinca* Cuvier, 1816

25. *Tinca tinca* (Linnaeus, 1758)

Vernacular names

Linjak. Tench.

Previous names used for the species

Cyprinus tinca Linnaeus, 1758; *Tinca vulgaris* Fleming, 1828.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to the Adriatic Basin (Glamuzina *et al.* 2017).

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Ćurčić 1910; Plančić 1923b; Aganović 1952a, 1957, 1967; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Žitnan *et al.* 1969; Vuković & Ivanović 1971; Juretić *et al.* 1977; Berberović *et al.* 1978; Kosorić 1978; Vuković & Kosorić 1978; Nadaždin *et al.* 1980; Vuković *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Veljović 1982; Mikavica 1987b; Mikavica *et al.* 1991; Leiner 1998; Mikavica & Savić 1999; Radević 2000; Škrijelj *et al.* 2005a, 2005b; Bogut *et al.* 2006; Adrović 2007; Hamzić & Lelo 2007; Kottelat & Freyhof 2007; Has-Schön *et al.* 2008; Suić *et al.* 2008; Šanda *et al.* 2008a, 2008b; Vuković *et al.* 2008; Sofradžija 2009; Perea *et al.* 2010; Skenderović *et al.* 2010, 2011b, 2020a, 2020b; Adrović 2012, 2018; Tutman *et al.* 2012a; Hasković *et al.* 2013; Čičić-Močić 2014, 2019; Geiger *et al.* 2014; Simonović *et al.* 2015; Skenderović 2015; Glamuzina *et al.* 2008a, 2017; Mrdak & Milošević 2017; Kovačević *et al.* 2018; Memić 2018; Mušović *et al.* 2018, 2020; Nedić *et al.* 2018; Žujo-Zekić *et al.* 2018; Adrović *et al.* 2021, Dekić *et al.* 2024.

Family Acheilognathidae Bleeker, 1863

Genus *Rhodeus* Agassiz, 1832

26. *Rhodeus amarus* (Bloch, 1782)

Vernacular names

Gavčica. European Bitterling.

Previous names used for the species

Rhodeus sericeus (Pallas, 1776); *Cyprinus amarus* Bloch, 1782; *Rhodeus sericeus amarus* (Bloch, 1782).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Literature

Glowacki 1896; Jankelić 1963; Vuković 1963a, 1977a; Sofradžija *et al.* 1975; Kosorić *et al.* 1983, 1989; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002; Bogut *et al.* 2006; Skenderović *et al.* 2006; Adrović 2007; Kottelat & Freyhof 2007; Skenderović & Adrović 2008b; Vuković *et al.* 2008; Sofradžija 2009; Drešković *et al.* 2011; Adrović 2012, 2018; Bănăduc & Curtean-Bănăduc 2014; Čičić-Močić 2014, 2016, 2019; Delić *et al.* 2014; Simonović *et al.* 2015; Skenderović 2015; Lukić *et al.* 2016; Glamuzina *et al.* 2017; Golub *et al.* 2017c, 2018a, 2018b; Kovačević *et al.* 2018; Memić 2018; Lolić *et al.* 2019; Cvijić *et al.* 2020; Dekić *et al.* 2020a, 2020b, 2024; Skenderović *et al.* 2020a.

Family Gobionidae Bleeker, 1863

Genus *Gobio* Cuvier, 1816

Taxonomic status of the gudgeons of the genus *Gobio* from the Danube River Basin requires thorough revision (Takács *et al.* 2014; Zangl *et al.* 2020).

27. *Gobio obtusirostris* Valenciennes, 1842

Vernacular names

Obična ili Dunavska Krkuš. Danube gudgeon.

Previous names used for the species

Gobio gobio (Linnaeus, 1758); *Gobio vulgaris* Heckel, 1837; *Gobio fluviatilis* Cuvier, 1842; *Gobio gobio obtusirostris* Valenciennes, 1844.

Origin and geographic range in BiH

Endemic (DBE). Native to the Danube River, translocated to the Adriatic Basin.

Remarks

Previous literature (Vuković 1963a, 1977a) mentioned it as *Gobio gobio*, which was a result of misidentification of specimens, and likely refer to *G. obtusirostris* (Takács *et al.* 2014; Zangl *et al.* 2020).

Literature

Głowacki 1896; Medić 1896; Zaplata & Taler 1932; Aganović & Milošević 1959; Vuković 1963a, 1966a, 1977a, Šenk 1969; Šenk & Kaluđerčić 1963; Vuković & Seratlić-Savić 1970; Vuković & Ivanović 1971; Duran 1975; Kosorić *et al.* 1975, 1980; Medić-Žuljević 1975; Sofradžija & Berberović 1975; Uršić 1975; Habeković *et al.* 1981; Kapetina 1981; Kosorić 1981; Obratil 1982; Veljović 1982, 1985; Sofradžija *et al.* 1984; Mikavica 1987b; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002, 2006; Memić & Adrović 2005; Mustafić *et al.* 2005; Zorlak 2005; Bogut *et al.* 2006; Memić 2006; Skenderović *et al.* 2006, 2020a; Adrović 2007; Kottelat & Freyhof 2007; Muhamedagić *et al.* 2008; Skenderović & Adrović 2008b; Vuković *et al.* 2008; Sofradžija 2009; Adrović 2012, 2018; Golub *et al.* 2012, 2016a, 2017c, 2018a, 2018b; Ivanc & Dekić 2012; Čičić-Močić 2014, 2016, 2018, 2019; Simonović *et al.* 2015; Skenderović 2015; Dekić *et al.* 2017b, 2020a, 2020b; Keško 2018; Kovačević *et al.* 2018; Memić 2018, 2020; Paspalj *et al.* 2018; Lolić *et al.* 2019; Muhamedagić *et al.* 2019; Cvijić *et al.* 2020.

Genus *Pseudorasbora* Bleeker, 1860

28. *Pseudorasbora parva* (Temminck & Schlegel, 1846)

Vernacular names

Bezribica. Topmouth gudgeon.

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Non-native species that originally occurs in eastern Asia, spreading rapidly in BiH lakes and rivers; invasive and harmful alien species. According to EU Regulation, it is considered an invasive species of EU concern.

Literature

Bogut *et al.* 2006; Adrović & Skenderović 2007; Šanda *et al.* 2008a, 2008b; Vuković *et al.* 2008; Sofradžija 2009; Tutman *et al.* 2012a; Simonović *et al.* 2015; Adrović *et al.* 2017; Glamuzina *et al.* 2017; Golub *et al.* 2018b; Mušović *et al.* 2018, 2020.

Genus *Romanogobio* Bănărescu, 1961

29. *Romanogobio carpathorossicus* (Vladykov, 1925)

Vernacular names

Istočna govedarka, istočna krkuša. Romanian gudgeon.

Previous names used for the species

Gobio kessleri Dybowski, 1862; *Gobio kessleri kessleri* Dybowski, 1862.

Origin and geographic range in BiH

Native. Danube River Basin.

Remarks

Previously reported as *Gobio (Romanogobio) kessleri*, but according to new literature data (Friedrich *et al.* 2018; Freyhof 2024), *R. kesslerii* is distributed in the lower Danube River Basin and the Baltic Basin, while in the upper and middle Danube River Basin *Romanogobio carpathorossicus* (Vladykov, 1925) is present.

Literature

Vuković 1963a, 1977a; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Milanović *et al.* 2015; Simonović *et al.* 2015; Friedrich *et al.* 2018; Kovačević *et al.* 2018; Lolić *et al.* 2019; Dekić *et al.* 2024; Freyhof 2024.

30. *Romanogobio uranoscopus* (Agassiz, 1828)

Vernacular names

Tankorepa krkuša. Stone gudgeon.

Previous names used for the species

Gobio uranoscopus (Agassiz, 1828).

Origin and geographic range in BiH

Endemic (DBE). Danube River Basin.

Literature

Seeley 1886; Glowacki 1896; Vuković 1963a, 1977a; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Golub *et al.* 2012; Milanović *et al.* 2015; Simonović *et al.* 2015; Dekić *et al.* 2024.

31. *Romanogobio vladykovi* (Fang, 1943)

Vernacular names

Dunavska bjeloperajna krkuša. Danube whitefin gudgeon.

Previous names used for the species

Gobio vladykovi Fang, 1943; *Romanogobio albipinnatus vladykovi* (Lukasch, 1933).

Origin and geographic range in BiH

Endemic (DBE). Danube River Basin.

Remarks

Mentioned by Sofradžija (2009) as *Romanogobio albipinnatus* (Lukasch, 1933), although this is possibly a result of misidentification, likely referring to specimens of *R. vladykovi*.

Literature

Naseka *et al.* 1999; Mrakovčić *et al.* 2002, 2006; Mustafić *et al.* 2005; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Milanović *et al.* 2015; Simonović *et al.* 2015; Dekić *et al.* 2024.

Family Leuciscidae Bonaparte, 1835

Genus *Abramis* Cuvier, 1816

32. *Abramis brama* (Linnaeus, 1758)

Vernacular names

Deverika. Bream.

Previous names used for the species

Cyprinus brama Linnaeus, 1758; *Abramis vulgaris* Mauduyt, 1849; *Abramis brama danubii* Pavlov, 1956.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to the Adriatic Basin (Glamuzina *et al.* 2017).

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Ćurčić 1910; Plančić 1923b; Zaplata & Taler 1932; Vuković 1963a, 1977a; Čanković *et al.* 1968a, 1968b; Ristić 1972; Berberović 1980; Kosorić *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Veljović 1982, 1985; Delić 1984; Mikavica 1987b; Mikavica *et al.* 1991; Pažur 1993; Leiner 1998; Mikavica & Savić 1999; Radević 2000, 2002;

Simonović 2001; Mrakovčić *et al.* 2002; Đonko 2005; Škrijelj *et al.* 2005a, 2005b; Bogut *et al.* 2006; Skenderović *et al.* 2006, 2010, 2011a, 2011b, 2015, 2020a, 2020b, 2021; Adrović 2007, 2012; Kottelat & Freyhof 2007; Skenderović & Adrović 2008a, 2008b, 2019; Suić *et al.* 2008; Vuković *et al.* 2008; Adrović *et al.* 2009a, 2009b; 2021; Sofradžija 2009; Čemer 2010; Bartulović *et al.* 2011; Hajdarević 2011; Tutman *et al.* 2012a; Čičić-Močić 2014; Riđanović *et al.* 2015; Simonović *et al.* 2015; Bajrić *et al.* 2016; Čulibrk *et al.* 2016; Nedić *et al.* 2014a, 2014c, 2014d; 2016a, 2018; Glamuzina *et al.* 2017; Golub *et al.* 2017c; Kovačević *et al.* 2018; Memić 2018, 2020; Dekić *et al.* 2020a, 2024; Kalamujić Stroil *et al.* 2021.

Genus *Alburnoides* Jeitteles, 1861

33. *Alburnoides bipunctatus* (Bloch, 1782)

Vernacular names

Dvoprugasta uklija, pliska. Spirlin.

Previous names used for the species

Alburnus bipunctatus bipunctatus (Bloch, 1782).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

It is possible that *Alburnoides bipunctatus* does not occur in the waters of BiH. The taxonomic status of this species needs revision, as three distinct genetic evolutionary lineages have been revealed in the Danube River Basin, with two recorded in BiH, both possibly representing new species still not described (Stierandová *et al.* 2016). Vukić *et al.* (2019) reports a record of astill unnamed *Alburnoides* sp. from the Neretva River basin. Based on the molecular analyses of cytochrome b gene this population belongs to *Alburnoides* sp. 1 sensu Stierandová *et al.* (2016) inhabiting the Sava River Basin (Danube River Basin).

Literature

Anonymous 1886; Glowacki 1896; Zaplata & Taler 1932, Vuković 1963a, 1977a; Berberović 1967; Vuković & Vuković 1968b, 1974; Žuljević 1970; Aganović & Kapetanović 1973; Martinović 1974; Kosorić *et al.* 1975, 1980; Đurović & Vuković 1975; Đurović 1979; Sofradžija *et al.* 1979, 1980, 1984; Kosorić 1981; Mikavica 1988a; Mikavica *et al.* 1991; Mikavica & Savić 1999; Babić & Mikavica 1994; Radević 2000; Treer *et al.* 2000; Korjenić 2005a; Sofradžija *et al.* 1979, 1980; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Skenderović *et al.* 2006, 2020b; Škrijelj *et al.* 2006; Kottelat & Freyhof 2007; Skenderović & Adrović 2008b; Sofradžija 2009; Golub *et al.* 2012, 2017c, 2018a, 2018b; Čičić-Močić 2014, 2016, 2018, 2019; Delić *et al.* 2014; Simonović *et al.* 2015; Skenderović 2015; Stierandová *et al.* 2016; Dekić *et al.* 2017b, 2020a, 2020b, 2024; Adrović 2018; Adrović *et al.* 2018; Keško 2018; Kovačević *et al.* 2018; Memić 2018, 2020; Lolić *et al.* 2019; Muhamedagić *et al.* 2019; Vukić *et al.* 2019; Cvijić *et al.* 2020.

Genus *Alburnus* Rafinesque, 1820

The taxonomic status of *Alburnus* in the Adriatic Basin has not been well studied and the systematics of many species require clearer resolution (Buj *et al.* 2010).

Alburnus albidus (Costa, 1838)

Previous names used for the species

Alburnus albidus alborella (Costa, 1838).

Remarks

Vuković (1963a) and Sofradžija (2009) reported *Alburnus albidus alborella* for the waters of the Adriatic Basin. According to Kottelat & Freyhof (2007), *A. albidus* is endemic to southern Italian rivers of the Adriatic, Ionian, and Tyrrhenian coast, and therefore is not included in the list of fishes of BiH.

34. *Alburnus alburnus* (Linnaeus, 1758)

Vernacular names

Zela, uklija. Bleak.

Previous names used for the species

Alburnus alburnus alburnus (Linnaeus, 1758); *Alburnus lucidus* Heckel, 1843.

Origin and geographic range in BiH

Native. Danube River Basin.

Remarks

Vuković (1963a), Vuković & Ivanović (1971) and Sofradžija (2009) reported it as *Alburnus alburnus* for the waters of the Danube River Basin; Vuković (1977a) and Bogut *et al.* (2006) reported the subspecies *A. alburnus alburnus* for the waters of the Danube River Basin.

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Medić 1901; RHZZM 1908; Ćurčić 1910; Plančić 1923b; Zaplata & Taler 1932; Aganović & Milošević 1959; Vuković 1963a, 1968a, 1977a; Kosorić & Vuković 1966a; Vuković & Seratlić-Savić 1967; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968, 1969; Vuković & Ivanović 1971; Vuković & Vuković 1974; Kosorić 1978; Orelj 1979; Sofradžija *et al.* 1979, 1980; Berberović 1980; Bianco 1980; Kosorić *et al.* 1980, 1983, 1989; Pejić *et al.* 1980; Sofradžija *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Mikavica 1987b; Economidis & Bănărescu 1991; Mikavica *et al.* 1991; Mrakovčić *et al.* 1995, 2006; Leiner 1998; Korjenić 1999; Mikavica & Savić 1999; Radević 2000; Simonović 2001; Škrijelj 2002; Škrijelj *et al.* 2003, 2005a, 2005b; Korjenić *et al.* 2003, 2005; Kadribašić 2005; Bogut *et al.* 2006; Skenderović *et al.* 2006, 2010, 2011a, 2011b, 2015, 2020a, 2020b, 2021; Adrović 2007, 2012, 2018; Kottelat & Freyhof 2007; Skenderović & Adrović 2008b, 2012, 2019; Vuković *et al.* 2008; Zupančić 2008; Sofradžija 2009; Buj *et al.* 2010; Bećiraj & Šahinović 2012; Golub *et al.* 2012, 2013, 2016a, 2017c, 2018a, 2018b, 2019; Čičić-Močić 2014, 2016, 2019; Delić *et al.* 2014; Nedić *et al.* 2014a, 2014c, 2014d, 2016b, 2018; Riđanović *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Bećiraj *et al.* 2016; Ćulibrk *et al.* 2016; Dekić *et al.* 2017b, 2020a, 2020b; Kovačević *et al.* 2018; Memić 2018; Lolić *et al.* 2019; Zuliani *et al.* 2019; Cvijić *et al.* 2020; Pilić *et al.* 2020.

Alburnus arborella (Bonaparte, 1841)

Vernacular names

Primorska uklija. Italian bleak.

Previous names used for the species

Alburnus alborella (De Filippi, 1844), *Alburnus alburnus alborella* (De Filippi, 1844).

Remarks

According to the literature (Kosorić & Vuković 1966a; Kosorić *et al.* 1989; Mrakovčić *et al.* 1995; Škrijelj 2002; Bogut *et al.* 2006), this species was reported in the middle and lower section of the Neretva River downstream to its mouth, and in the Tihaljina/Trebižat River catchment and Buna River in BiH. However, the population from the Neretva River catchment is now recognized as a different species, *Alburnus neretvae* (Buj *et al.*, 2010). Recent literature (Žujo-Zekić 2021; Žujo-Zekić *et al.* 2018, 2020) mentioned the presence of *A. arborella* in the Buško Lake, however, without providing diagnostic characters. *Alburnus arborella* is distributed in rivers which drain into the northern part of the Adriatic Sea from the Chienti drainage basin in Marche to Ancona Province (Slovenia, Switzerland, Italy) and the Tyrrhenian basin in the Arno drainage (Italy) (Kottelat & Freyhof 2007). The distribution and taxonomic status of the populations of *Alburnus* in the Adriatic Basin have not been fully resolved and require additional research (Geiger *et al.* 2014), and *A. arborella* is therefore not included in the checklist of BiH.

Literature

Kosorić & Vuković 1966a; Čanković 1967; Čanković *et al.* 1968a; Ivanović 1968; Vuković & Ivanović 1971; Kosorić *et al.* 1989; Mrakovčić *et al.* 1995; Škrijelj 2002; Bogut *et al.* 2006; Šanda *et al.* 2008a, 2008b; Buj *et al.* 2010; Perea *et al.* 2010; Žujo-Zekić 2021; Žujo-Zekić *et al.* 2018, 2020.

35. *Alburnus neretvae* Buj, Šanda & Perea, 2010

Vernacular names

Neretvanska uklija. Neretva bleak.

Previous names used for the species

Alburnus arborella (Bonaparte, 1841); *Alburnus alburnus alborella* (De Filippi, 1844).

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

The population from the Neretva River catchment previously identified as *Alburnus arborella* has been described as a different species, *A. neretvae* (Buj *et al.* 2010), endemic to the Neretva River catchment in Croatia and BiH, also recorded in waters of the karstic fields Rastoke and Jezero near Vrgorac, Kuti and Baćinska Lakes, and the lower Neretva River in Croatia. In BiH, it was recorded in the Bregava River, Hutovo Blato wetland, Mušnica River (Gatačko Polje field), Trebišnjica River (Popovo Polje field) (Buj *et al.* 2010), and previously under the name *A. arborella* in the Tihaljina/Trebižat River catchment, Buna River (Kosorić & Vuković 1966a) and in the lower and middle sections of the Neretva River, from the mouth of the Buna River downstream to the mouth of the Neretva River into the sea (Kosorić & Vuković 1966a; Kosorić *et al.* 1989; Mrakovčić *et al.* 1995). It was also recorded in the Salakovac reservoir, approximately 20 km upstream of Mostar (Škrijelj 2002). Pronounced genetic differentiation was revealed between most of the mentioned populations (Charpila *et al.* 2025). As this species is very similar to the congeneric Italian bleak (*A. arborella*) in terms of anatomical characters, additional systematic research is recommended (Geiger *et al.* 2014).

Literature

Kosorić & Vuković 1966a; Žuljević 1967; Muhanović 1984; Burek 1985; Sadović 1986; Kosorić *et al.* 1989; Mrakovčić *et al.* 1995; Škrijelj 2002; Šanda *et al.* 2008a, 2008b; Buj *et al.* 2010; Perea *et al.* 2010; Tutman *et al.* 2012a; Darwall *et al.* 2014; Oikonomou *et al.* 2014; Mrdak & Milošević 2017; Benovics *et al.* 2019; Skenderović *et al.* 2020b; Charmpila *et al.* 2025.

36. *Alburnus sava* Bogutskaya, Zupančič, Jelić, Diripasko & Naseka, 2017

Vernacular names

Bucov. Sava bleak.

Previous names used for the species

Chalburnus chalcoides (Güldenstädt, 1772); *Chalchalburnus chalcoides mento* (Agassiz, 1832); *Alburnus mento* (Heckel, 1837); *Chalburnus chalcoides danubicus* (Antipa, 1909); *Alburnus sarmaticus* Freyhof & Kottelat, 2007.

Origin and geographic range in BiH

Endemic (DBE). Danube River Basin.

Remarks

Vuković (1963a) listed *Chalburnus chalcoides danubicus* in BiH for waters of the Danube River Basin, while Vuković (1977a), Bogut *et al.* (2006) and Sofradžija (2009) listed *C. chalcoides*. Kottelat & Freyhof (2007) considered the same species to be *Alburnus sarmaticus*. However, Bogutskaya *et al.* (2017) describe the population from the upper Sava River (Danube River Basin) as a different species (*A. sava*), but only individuals from the Kupa River (Croatia and Slovenia) were available for examination. Vucić *et al.* (2017) updated the distribution revealing a much more widespread occurrence in the Sava River and its catchment (Bosna River near Zenica), and also in the Drina River in BiH. Additional systematic research from the entire Sava catchment is recommended.

Literature

Anonymous 1886; Glowacki 1896; Ćurčić 1910; Vuković 1963a, 1977a; Čanković *et al.* 1968a, 1968c; Vuković & Ivanović 1971; Obratil 1982; Veljović 1982; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Vuković *et al.* 2008; Sofradžija 2009; Drešković *et al.* 2011; Milanović *et al.* 2015; Mustafić *et al.* 2015; Simonović *et al.* 2015; Bogutskaya *et al.* 2017; Golub *et al.* 2017c; Vucić *et al.* 2017; Skenderović *et al.* 2020b; Dekić *et al.* 2024.

Genus *Ballerus* Heckel, 1843

37. *Ballerus ballerus* (Linnaeus, 1758)

Vernacular names

Kesega. Blue bream.

Previous names used for the species

Abramis ballerus (Linnaeus, 1758).

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Glowacki 1896; Medić 1902; Ćurčić 1910; Plančić 1923b; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Veljović 1982, 1985; Delić 1984; Radević 2000; Simonović 2001; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Simonović *et al.* 2015; Kovačević *et al.* 2018.

38. *Ballerus sapa* (Pallas, 1814)

Vernacular names

Crnooka deverika. White-eye bream.

Previous names used for the species

Abramis sapa (Pallas, 1811).

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Glowacki 1896; Ćurčić 1910; Plančić 1923b; Vuković 1963a, 1977a; Veljović 1982, 1985; Radević 2000, 2002; Simonović 2001; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Ćemer 2010; Čičić-Močić 2014, 2016, 2019; Simonović *et al.* 2015; Adrović 2018; Memić 2018; Skenderović & Adrović 2019; Skenderović *et al.* 2020a, 2020b, 2021; Dekić *et al.* 2024.

Genus *Blicca* Heckel, 1843

39. *Blicca bjoerkna* (Linnaeus, 1758)

Vernacular names

Krupatica. Silver bream.

Previous names used for the species

Abramis bjoerkna (Linnaeus, 1758); *Abramis bjoerkna bjoerkna* (Linnaeus, 1758); *Abramis björkna* (Linnaeus, 1758); *Abramis blicca* (Bloch, 1782); *Blicca argyroleuca* Heckel, 1843.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Ćurčić 1910; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Berberović 1980; Obratil 1982; Veljović 1982, 1985; Delić 1984; Mikavica *et al.* 1991; Leiner 1998; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Vuković *et al.* 2008; Sofradžija 2009; Golub *et al.* 2013; Simonović *et al.* 2015; Nedić *et al.* 2016a, 2016b, 2018; Skenderović & Adrović 2019; Skenderović *et al.* 2020b, 2021; Dekić *et al.* 2024.

Genus *Chondrostoma* Agassiz, 1832

40. *Chondrostoma knerii* Heckel, 1843

Vernacular names

Podustva. Dalmatian nase.

Previous names used for the species

Chondrostoma kneri Heckel, 1843; *Chondrostomus knerii* Heckel, 1843.

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

Geographically restricted to the middle and lower Neretva River catchment in BiH and Croatia (Tutman *et al.* 2019); present up to the Grabovica hydropower plant, Hutovo Blato wetland, Buna and Trebižat Rivers (Hamzić 2024).

Literature

Aganović 1952a; Vuković 1963a, 1964, 1977a; Kosorić & Vuković 1966a; Čanković 1967; Čanković *et al.* 1968a; Berberović *et al.* 1970b, 1970c; Vuković *et al.* 1970b; Guzina *et al.* 1971; Kaluđerčić *et al.* 1971; Vuković & Ivanović 1971; Švob & Kilalić 1972; Vuković & Vuković 1974; Aganović & Kapetanović 1978; Guzina & Vuković 1978; Kosorić 1978; Seratlić *et al.* 1978; Vuković & Kosorić 1978; Kosorić *et al.* 1983, 1989; Elvira 1987; Vuković & Sofradžija 1987; Mikavica & Bošnjak 1989; Mikavica & Dizdarević 1990; Economidis & Bănărescu 1991; Crivelli 1996; Elvira 1997; Mikavica 1998; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Glamuzina *et al.* 2007, 2008a; Kottelat & Freyhof 2007; Robalo *et al.* 2007; Jelić *et al.* 2008; Šanda *et al.* 2008a, 2008b; Tutman *et al.* 2008b, 2012a, 2019; Zupančić 2008; Dulčić *et al.* 2009; Sofradžija 2009; Perea *et al.* 2010; Drešković *et al.* 2011; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Milanović *et al.* 2015; Benovics *et al.* 2019; Freyhof *et al.* 2020; Skenderović *et al.* 2020b; Dekić *et al.* 2024.

41. *Chondrostoma nasus* (Linnaeus, 1758)

Vernacular names

Podust, škobalj. Nase.

Previous names used for the species

Chondrostoma nasus nasus (Linnaeus, 1758); *Cyprinus nasus* Linnaeus, 1758.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Medić 1902; Langhoffer 1904; Čurčić 1910; Plančić 1923b; Zaplata & Taler 1932; Taler 1945b; Kosorić & Kardoš 1955; Aganović & Milošević 1959; Smlatić 1961; Pavlović *et al.* 1960; Vuković 1963a, 1977a; Čanković 1964, 1968a; Čanković *et al.* 1968a, 1968b; Kekić 1968; Šenk & Aganović 1968; Kiškarolj & Čanković 1969; Šenk 1969; Vuletić 1969; Berberović *et al.* 1970c; Vuković & Ivanović 1971; Aganović & Kapetanović 1973; Hadžović 1975; Kosorić *et al.* 1975, 1980; Vuković & Kosorić 1978; Habeković *et al.* 1981; Kačanski *et al.* 1981;

Kosorić 1981; Sofradžija & Hadžiselimović 1981b; Obratil 1982; Veljović 1982, 1985; Mikavica 1987b, 1988a; Bogičević 1988; Mikavica & Bošnjak 1989; Mikavica *et al.* 1989, 1991, 1997; Mikavica & Dizdarević 1990; Babić & Mikavica 1994; Elvira 1997; Mikavica & Savić 1999; Radević 2000, 2002; Mrakovčić *et al.* 2002; Alić *et al.* 2004; Škrijelj *et al.* 2005a, 2005b; Bogut *et al.* 2006; Skenderović *et al.* 2006, 2010, 2011a, 2011b, 2015, 2020a, 2020b; Kottelat & Freyhof 2007; Mehanović 2007; Skenderović & Adrović 2008b, 2019; Vuković *et al.* 2008; Sofradžija 2009; Vujanović *et al.* 2011; Golub *et al.* 2012, 2017a, 2017c, 2018a, 2018b; Muhamedagić & Habibović 2013; Čičić-Močić 2014, 2016, 2018; Delić *et al.* 2014; Simonović *et al.* 2015; Bećiraj *et al.* 2016; Dekić *et al.* 2017b, 2020a, 2020b; 2024; Kovačević *et al.* 2018; Memić 2018; Nedić *et al.* 2018; Lolić *et al.* 2019; Cvijić *et al.* 2020.

42. *Chondrostoma phoxinus* Heckel, 1843

Vernacular names

Podbila. Livno nase.

Previous names used for the species

Chondrostomus phoxinus Heckel, 1843.

Origin and geographic range in BiH

Endemic (CCE). Adriatic Basin.

Remarks

Endemic to the Cetina River catchment in BiH and Croatia. Distributed in Western Bosnia, the catchment of the Cetina River, sinks and sources of Glamočko Polje, Livanjsko Polje, Duvanjsko Polje fields, as well as Buško Blato, Blidinjsko and Mandek lakes (Vuković 1977a, Hamzić 2024).

Literature

Heckel & Kner 1858; Ćurčić 1916; Karaman 1923, 1928; Taler 1951a, 1953a; Vuković 1963a, 1964, 1977a; Aganović 1969; Vuković & Žnidaršić-Krzyk 1969; Berberović *et al.* 1970a, 1970b, 1970c; Lončar 1970; Vuković *et al.* 1970b; Kaluđerčić *et al.* 1971; Veledar 1971; Vuković & Ivanović 1971; Veledar & Kosorić 1972, 1974; Jerković *et al.* 1973; Vuković & Vuković 1974; Kaćanski *et al.* 1978; Berberović 1980; Mikavica & Kosorić 1986; Elvira 1987; Vuković & Sofradžija 1987; Ibrašimović 1988; Mikavica & Bošnjak 1989; Mikavica & Dizdarević 1990; Economidis & Bănărescu 1991; Elvira 1997; Kottelat 1997; Kerovec *et al.* 1998; Mikavica 1998; Žujo 2004; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008b; Zupančić 2008; Čaleta *et al.* 2009b; Sofradžija 2009; Perea *et al.* 2010; Drešković *et al.* 2011; Žujo-Zekić 2012, 2013, 2021; Darwall *et al.* 2014; Geiger *et al.* 2014; Milanović *et al.* 2015; Mustafić & Mrakovčić 2017; Žujo-Zekić *et al.* 2016, 2018; Benovics *et al.* 2019; Freyhof *et al.* 2020; Mušović *et al.* 2020; Dekić *et al.* 2024; Hamzić 2024.

Genus *Delminichthys* Freyhof, Lieckfeldt, Bogutskaya, Pitra & Ludwig, 2006

43. *Delminichthys adpersus* (Heckel, 1843)

Vernacular names

Imotska gaovica. Spotted minnow.

Previous names used for the species

Leucos adpersus Heckel, 1843; *Paraphoxinus adpersus* Heckel, 1843; *Phoxinellus adpersus* (Heckel, 1843).

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

Endemic to the Neretva River catchment in BiH and Croatia.

Literature

Seeley 1886; Kolombatović 1886; Kišpatić 1893; Sabioncello 1967; Berberović & Pavlović 1970; Berberović *et al.* 1970d, 1971b; Karaman 1972; Sofradžija & Berberović 1972; Bozja & Švob 1974; Vuković & Vuković 1974; Ristić 1977; Pavlović & Berberović 1978; Vuković 1982b; Vuković & Sofradžija 1987; Šorić 1992; Zupančič & Bogutskaya 2002; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Freyhof *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008b; Zupančič 2008; Sofradžija 2009; Perea *et al.* 2010; Drešković *et al.* 2011; Palandačić *et al.* 2012; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Milanović *et al.* 2015; Benovics *et al.* 2019; Freyhof *et al.* 2020; Reier *et al.* 2022, Dekić *et al.* 2024; Lukač *et al.* 2024.

44. *Delminichthys ghetaldii* Steindachner, 1882

Vernacular names

Popovska gaovica. Popovo minnow.

Previous names used for the species

Paraphoxinus ghetaldii Steindachner, 1882; *Phoxinellus ghetaldii* (Steindachner, 1882); *Paraphoxinus pstrosii* Steindachner, 1882; *Phoxinellus pstrossi* Steindachner, 1882.

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

In earlier literature, the species was mentioned as *Paraphoxinus ghetaldi* (Vuković 1977a) and *Phoxinellus ghetaldii* in Sofradžija (2009). It is now considered as endemic to the Neretva River catchment in BiH and Croatia. However, due to the unresolved taxonomic status of related species, its occurrence in the Neretva River catchment is uncertain (Zupančič & Bogutskaya 2002). *Phoxinellus pstrossi* was mentioned by Vuković (1963a, 1977a) as occurring in the Trebišnjica River catchment, but according to Zupančič & Bogutskaya (2002), it is tentatively considered as a junior synonym of *Delminichthys ghetaldii*.

Literature

Steindachner 1882a, 1882b, 1882c, 1883; Kišpatić 1893; Ćurčić 1913, 1915a, 1915b; Taler 1951a, 1952, 1953a, 1953b; Vuković 1963a, 1967b, 1977a, 1977b, 1982b; Kosorić & Vuković 1966a; Sket 1967; Berberović *et al.* 1969b, 1970d; Vuković & Miladinović 1969; Aganović & Kapetanović 1970, 1971; Guzina & Vuković 1972; Vuković & Ivanović 1971; Karaman 1972; Sofradžija & Berberović 1972; Sofradžija *et al.* 1972; Vuković & Vuković 1974; Ristić 1977; Pavlović & Berberović 1978; Pocrnjić *et al.* 1979; Kosorić *et al.* 1983, 1989; Leiner 1984; Vuković & Sofradžija 1987; Povž *et al.* 1990; Šorić 1992; Crivelli 1996; Kottelat 1997; Mikavica 1998; Zupančič & Bogutskaya 2000, 2002; Bogut *et al.* 2006; Mrakovčić *et al.*

2006; Freyhof *et al.* 2006, 2020; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Muhamedagić *et al.* 2008; Šanda *et al.* 2008b; Zupančič 2008; Pavlović *et al.* 2009; Sofradžija 2009; Palandačić *et al.* 2010; Dekić *et al.* 2011, 2012a, 2013a, 2014, 2016c, 2017a, 2024; Drešković *et al.* 2011; Darwall *et al.* 2014, Geiger *et al.* 2014; Oikonomou *et al.* 2014; Lolić *et al.* 2015; Milanović *et al.* 2015; Reier *et al.* 2022; Lukač *et al.* 2023, 2024.

Genus *Leucaspius* Heckel & Kner, 1858

45. *Leucaspius delineatus* (Heckel, 1843)

Vernacular names

Bjelica. Sun bleak.

Previous names used for the species

Squalius delineatus Heckel, 1843.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

It was translocated to the Adriatic Basin in Jablaničko Lake accidentally as bait for perch fishing (Pavličević *et al.* 2016; Glamuzina *et al.* 2017; Muhamedagić *et al.* 2019; Hamzić 2024).

Literature

Vuković 1963a, 1977a; Čanković *et al.* 1968a, 1968c; Veljović 1982, 1985; Radević 2000; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Pavličević *et al.* 2016; Glamuzina *et al.* 2017; Muhamedagić *et al.* 2019; Skenderović *et al.* 2020b; Dekić *et al.* 2024, Hamzić 2024.

Genus *Leuciscus* Cuvier, 1816

46. *Leuciscus aspius* (Linnaeus, 1758)

Vernacular names

Bolen. Asp.

Previous names used for the species

Aspius aspius (Linnaeus, 1758); *Aspius aspius aspius* (Linnaeus, 1758); *Cyprinus aspius* Linnaeus, 1758; *Aspius rapax* Agassiz, 1835.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Horvat 1901; Medić 1901; Ćurčić 1910; Plančić 1923b; Knop 1947; Vuković 1963a, 1977a; Žitnan *et al.* 1969; Berberović 1980; Habeković *et al.* 1981; Veljović 1982, 1985; Mikavica *et al.* 1991; Pažur 1993; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002, 2006; Adrović 2007, 2012; Kottelat & Freyhof 2007; Mateš 2008; Suić *et al.* 2008; Sofradžija 2009; Skenderović *et al.* 2010, 2011b, 2020b; Drešković *et al.* 2011; Čičić-Močić 2014; Nedić *et al.* 2014a, 2014c, 2014d, 2018; Skenderović 2015; Riđanović

et al. 2015; Simonović *et al.* 2015; Lukić *et al.* 2016; Kovačević *et al.* 2018; Memić 2018; Dekić *et al.* 2024.

47. *Leuciscus idus* (Linnaeus, 1758)

Vernacular names

Jaz. Ide.

Previous names used for the species

Idus melanotus (Heckel, 1843).

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Ćurčić 1910; Plančić 1923b; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Sofradžija 1977; Berberović 1980; Kosorić *et al.* 1980; Obratil 1982; Veljović 1982; Delić 1984; Veljović 1985; Mikavica *et al.* 1991; Pažur 1993; Mikavica & Savić 1999; Radević 2000, 2002; Mrakovčić *et al.* 2002, 2006; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Suić *et al.* 2008; Sofradžija 2009; Simonović *et al.* 2015; Golub *et al.* 2017c; Kovačević *et al.* 2018; Skenderović *et al.* 2020b; Dekić *et al.* 2020a, 2024.

48. *Leuciscus leuciscus* (Linnaeus, 1758)

Vernacular names

Kljenić. Common Dace.

Previous names used for the species

Cyprinus dobula Linnaeus, 1758; *Cyprinus leuciscus* Linnaeus, 1758; *Leuciscus dobula* (Linnaeus, 1758); *Leuciscus vulgaris* Fleming, 1828; *Squalius lepusculus* Heckel, 1852.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Glowacki 1896; Vuković 1963a, 1977a; Šenk & Aganović 1968; Sofradžija 1977; Veljović 1982, 1985; Radević 2000, 2002; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Vuković *et al.* 2008; Sofradžija 2009; Simonović *et al.* 2015; Golub *et al.* 2017c; Dekić *et al.* 2024.

Genus *Leucos* Heckel, 1843

49. *Leucos basak* Heckel, 1843

Vernacular names

Gera, plotica. Dalmatian roach.

Previous names used for the species

Rutilus rubilio rubilio (Bonaparte, 1758); *Rutilus rubilio* (Bonaparte, 1837); *Sarmarutilus rubilio* (Bonaparte, 1837); *Rutilus basak* (Heckel, 1843).

Origin and geographic range in BiH

Endemic (ABE). Adriatic Basin.

Remarks

Endemic to BiH and Croatia; in BiH, it is recorded in the lower Neretva River catchment (Neretva and its tributaries: Tihaljina, Trebižat, Jasenica, Bregava, Buna and Hutovo Blato wetland), and upstream to the confluence with the Buna River (Tutman *et al.* 2018). In the older literature, it was mentioned as *Rutilus rubilio* (Vuković 1963a, 1977a; Kosorić & Vuković 1966a, 1966b; Vuković & Seratlić-Savić 1967; Vuković & Ivanović 1970; Vuković & Vuković 1970), in Bogut *et al.* (2006) and in Sofradžija (2009) as *R. basak*, for the lower Neretva River. Subsequently, Bianco & Ketmaier (2014) revised the taxonomy of the genus *Rutilus*, creating the genus *Leucos*, and the species in the Neretva River catchment was regarded as *Leucos basak*.

Literature

Heckel & Kner 1858; Karaman 1928; Aganović 1952a, 1967; Vuković 1963a, 1968a, 1970, 1977a; Kosorić & Vuković 1966a, 1966b; Vuković & Seratlić-Savić 1967; Vuković & Vuković 1968a, 1970, 1974; Berberović *et al.* 1970a; Vuković *et al.* 1970a; Vuković & Ivanović 1970, 1971; Seratlić-Savić & Vuković 1969, 1971; Guzina *et al.* 1971; Berberović & Sofradžija 1972; Knežević & Ivanović 1975; Knežević 1977; Aganović & Kapetanović 1978; Guzina & Vuković 1978; Kosorić 1978; Vuković & Kosorić 1978; Berberović 1980; Marić 1981; 1988; Kosorić *et al.* 1983, 1989; Grgurević 1984; Bianco & Taraborelli 1985; Galić 1986; Ljubojević 1989; Economidis & Bănărescu 1991; Crivelli 1996; Mikavica 1998; Glamuzina *et al.* 2001, 2002, 2008a; Tutman *et al.* 2002, 2009a, 2009b, 2012a, 2018; Bianco *et al.* 2004; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Muhamedagić *et al.* 2019; Šanda *et al.* 2008a, 2008b; Zupančič 2008; Dulčić *et al.* 2009; Sofradžija 2009; Marić 2010; Perea *et al.* 2010; Matić-Skoko *et al.* 2011; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Milanović *et al.* 2015; Benovics *et al.* 2019; Dekić *et al.* 2024.

Genus *Pelecus* Agassiz, 1835

50. *Pelecus cultratus* (Linnaeus, 1758)

Vernacular names

Sabljarka. Ziege.

Previous names used for the species

Cyprinus cultratus Linnaeus, 1758.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Medić 1902; Ćurčić 1910; Vuković 1963a, 1977a; Veljović 1982, 1985; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.*

2006; Škrijelj *et al.* 2005a, 2005b; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Milanović *et al.* 2015; Simonović *et al.* 2015; Lukić *et al.* 2016, Dekić *et al.* 2024.

Genus *Phoxinellus* Heckel, 1843

51. *Phoxinellus alepidotus* Heckel, 1843

Vernacular names

Pijurica. Adriatic minnow.

Previous names used for the species

Paraphoxinus alepidotus (Heckel, 1843); *Phoxinellus alepidotus* Heckel, 1843.

Origin and geographic range in BiH

Endemic (CCE). Adriatic Basin.

Remarks

Endemic to the Cetina River catchment in BiH and Croatia.

Literature

Heckel 1843; Heckel & Kner 1858; Gunther 1859; Steindachner 1882c; KaturiĆ 1883; Kolombatović 1886; Seeley 1886; Kišpatić 1893; Trgovčević 1905; Ćurčić 1916; Karaman 1923, 1928; Protić 1924; Taler 1953a, 1953b; Vuković 1963a, 1964, 1977a, 1977b, 1982b; Sabioncello 1967; Sket 1967; Biberdžić 1968; Aganović 1969; Berberović *et al.* 1969a, 1970d, Sofradžija *et al.* 1969; Vuković & Miladinović 1969; Vuković & žnidaršić-Krzyk 1969; Aganović & Kapetanović 1970; Vuković *et al.* 1970b, 1971a, 1971b; Guzina & Vuković 1972; Miladinović *et al.* 1971; Kaluđerčić *et al.* 1971; Trewavas 1971; Vuković & Ivanović 1971; Karaman 1972; Sofradžija & Berberović 1972; Bozja & Švob 1974; Vuković & Vuković 1974; Kačanski *et al.* 1978; Pocrnjić *et al.* 1979; Berberović 1980; Marić 1980, 1983, 1990; Leiner 1984; Howes 1985; Vuković & Sofradžija 1987; Povž *et al.* 1990; Škrijelj & Sofradžija 1990; Šorić 1992; Crivelli 1996; Kottelat 1997; Zupančić & Bogutskaya 2000, 2002; Škrijelj 2002; Bogutskaya & Zupančić 2003; Delić *et al.* 2005; Freyhof *et al.* 2006, 2020; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008b; Zupančić 2008; Sofradžija 2009; Palandačić *et al.* 2010; Perea *et al.* 2010; Dekić *et al.* 2011, 2016a, 2024; Drešković *et al.* 2011; Darwall *et al.* 2014; Milanović *et al.* 2015; Benovics *et al.* 2019; Muhamedagić *et al.* 2019; Žujo-Zekić 2021; Reier *et al.* 2022; Lukač *et al.* 2023; Palandačić *et al.* 2024.

52. *Phoxinellus pseudalepidotus* Bogutskaya & Zupančić, 2003

Vernacular names

Prikanac. Mostar minnow.

Origin and geographic range in BiH

Endemic (NCE, BiH). Adriatic Basin.

Remarks

Endemic to the Neretva River catchment; present data suggest that the species is restricted to the Mostarsko Blato wetland only (Bogutskaya & Zupančić 2003). However, it is considered as more widespread (Mihinjač *et al.* 2014).

Literature

Zupančič & Bogutskaya 2000, 2002; Bogutskaya & Zupančič 2003; Freyhof *et al.* 2006, 2020; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008a, 2008b, 2009; Zupančič 2008; Palandačić *et al.* 2010; Perea *et al.* 2010; Markotić *et al.* 2012, 2013a, 2013b, 2018, 2019; Darwall *et al.* 2014; Geiger *et al.* 2014; Mihinjač *et al.* 2014; Milanović *et al.* 2015; Benovics *et al.* 2019; Reier *et al.* 2022; Palandačić *et al.* 2024.

Genus *Phoxinus* Rafinesque, 1820

53. *Phoxinus karsticus* Bianco & De Bonis, 2015

Vernacular names

Krški pijor. Karstic minnow.

Origin and geographic range in BiH

Endemic (ABE). Adriatic Basin.

Remarks

Based on morphological characters, Bianco & De Bonis (2015) described *P. karsticus* from the endorheic system of the karstic field Popovo Polje within the Trebišnjica River catchment in BiH. It is probably endemic to the same area (Bianco & De Bonis 2015). Palandačić *et al.* (2015, 2017) supported the validity of the species using mitochondrial data, but this was only partially supported by nuclear data. Further studies are necessary to resolve its taxonomy. Distributed also in Montenegro (Palandačić *et al.* 2015, 2017).

Literature

Bianco & De Bonis 2015; Palandačić *et al.* 2015, 2017; Vucić *et al.* 2018; Reier *et al.* 2022; Lukač *et al.* 2023.

Phoxinus ketmaieri Bianco & De Bonis, 2015

Remarks

Based on morphological characters, Bianco & De Bonis (2015) described this species from Krk Island and the Zrmanja River in Croatia and assumed that its distribution likely also includes the Cetina River and other rivers of the Dalmatian ichthyogeographic district, such as the Krka and Neretva. However, according to Palandačić *et al.* (2017), *Phoxinus ketmaieri* is not valid based on molecular species delimitation analyses, and is likely a junior synonym of *P. lumaireul*. Therefore, it is not included in the list of fishes from BiH.

Literature

Bianco & De Bonis 2015; Palandačić *et al.* 2015, 2017; Vucić *et al.* 2018.

54. *Phoxinus lumaireul* (Schinz, 1840)

Vernacular names

Primorski pijor, primorska gaga. Italian minnow.

Previous names used for the species

Cyprinus phoxinus Linnaeus, 1758; *Leuciscus phoxinus* (Linnaeus, 1758); *Phoxinus laevis* Fitzinger, 1832; *Phoxinus phoxinus phoxinus* (Linnaeus, 1758).

Origin and geographic range in BiH

Native. Danube River and the Adriatic Basins.

Remarks

In the earlier literature, it was mentioned as *Phoxinus phoxinus* (Vuković 1963a, 1963c, 1966b, 1967a, 1977a). According to Kottelat (2007), the minnows from Italy, Slovenia, Croatia and BiH were recognised as *P. lumaireul* (Schinz, 1840), based on anatomical data. In BiH, *P. lumaireul* is distributed in both the Danube River and the Adriatic Basin. Results from molecular studies (Vucić *et al.* 2018) confirmed that *P. lumaireul* is a minnow species which prevails in Croatia and is also reported from the westernmost river systems of the Sava River catchment in BiH, with a remarkable genetic variability across the entire known distribution range. However, the taxonomic status and range of *P. lumaireul* in BiH remain uncertain (Palandačić *et al.* 2015, 2017; Neuburg *et al.* 2023). It is also necessary to resolve its taxonomy and the taxonomic status of the genus *Phoxinus* in the Adriatic Basin as well, due to the presence of several genetic lineages, which may represent yet unnamed species (Palandačić *et al.* 2015, 2017, 2020).

Literature

Brusina 1892; Glowacki 1896; Zaplata & Taler 1932; Protić 1933; Šenk 1962, 1969; Vuković 1963a, 1963c, 1966b, 1967a, 1977a; Berberović 1967; Čanković *et al.* 1968a; Vuković & Miladinović 1969; Vuković & Žnidaršić-Krzyk 1969; Vuković & Seratlić-Savić 1970; Kosorić *et al.* 1971; Berberović & Sofradžija 1972, 1974a; Kosorić 1974, 1977a, 1981; Kačanski *et al.* 1977; Kosorić *et al.* 1980, 1983, 1989; Guzina *et al.* 1981; Sofradžija *et al.* 1984; Kosorić & Mikavica 1986; Mikavica *et al.* 1991; Babić & Mikavica 1994; Mikavica & Savić 1999; Radević 2000; Škrijelj *et al.* 2006; Kottelat 2007; Kottelat & Freyhof 2007; Muhamedagić *et al.* 2008; Šanda *et al.* 2008a, 2008b; Sofradžija 2009; Perea *et al.* 2010; Bećiraj & Šahinović 2012; Golub *et al.* 2012, 2016a, 2018b; Delić *et al.* 2014; Palandačić *et al.* 2015, 2017; Simonović *et al.* 2015; Bećiraj *et al.* 2016, Čičić-Močić 2016, 2018, 2019; Keško 2018; Kovačević *et al.* 2018; Memić 2018, 2020; Vucić *et al.* 2018; Benovics *et al.* 2019; Muhamedagić *et al.* 2019; Dekić *et al.* 2020a, 2020b; Skenderović *et al.* 2020a, 2020b; Neuburg *et al.* 2023; Carosi *et al.* 2024.

Genus *Rutilus* Rafinesque, 1820

55. *Rutilus rutilus* (Linnaeus, 1758)**Vernacular names**

Bodorka. Roach.

Previous names used for the species

Leuciscus rutilus (Linnaeus, 1758); *Rutilus rutilus carpathorossicus* Vladykov, 1930.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

According to Hamzić (2024), translocated to the Adriatic Basin; in Hutovo blato wetland – Neretva catchment, and Bilećko lake – Trebišnjica catchment.

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Ćurčić 1910; Plančić 1923b; Aganović 1957; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968, 1969; Žitnan *et al.* 1969; Vuković & Aganović 1971; Vuković & Vuković 1974; Đurović & Vuković 1975; Guzina & Vuković 1978; Vuković & Kosorić 1978; Berberović 1980; Kosorić *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Veljović 1982, 1985; Delić 1984; Kujačić 1987; Ljubojević 1989; Mikavica *et al.* 1991; Leiner 1998; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002; Gajević 2005; Gajević & Korjenić 2005; Skenderović *et al.* 2006, 2011b, 2020a, 2020b; Škrijelj *et al.* 2005a, 2005b, 2006; Bogut *et al.* 2006; Adrović 2007, 2012, 2018; Kottelat & Freyhof 2007; Skenderović & Adrović 2008b; Vuković *et al.* 2008; Adrović *et al.* 2009a, 2009b; Sofradžija 2009; Bećiraj & Šahinović 2012; Golub *et al.* 2013; Čičić-Močić 2014, 2016, 2019; Delić *et al.* 2014; Simonović *et al.* 2015; Skenderović 2015; Ćulibrk *et al.* 2016; Bajrić *et al.* 2016; Bećiraj *et al.* 2016; Mrdak & Milošević 2017; Kovačević *et al.* 2018; Memić 2018; Zuliani *et al.* 2019; Dekić *et al.* 2020a; Kalamujić Stroil *et al.* 2021; Hamzić 2024.

56. *Rutilus virgo* (Heckel, 1852)

Vernacular names

Plotica, platnica. Danube roach.

Previous names used for the species

Leuciscus virgo Linnaeus, 1758; *Rutilus pigus* (Lacepède, 1804); *Rutilus pigus virgo* (Heckel, 1852).

Origin and geographic range in BiH

Endemic (DBE). Danube River Basin.

Remarks

First mentioned in Vuković (1963a) and later in Sofradžija (2009) as *Rutilus pigus virgo*, then in Vuković (1977a) and in Bogut *et al.* (2006) as *R. pigus*. However, *R. pigus* is a Padany/Venetian endemic that is differentiated genetically (Ketmaier *et al.* 2008) and morphologically (Kottelat & Freyhof 2007) from the similar *R. virgo*. Therefore, *R. pigus* is not included in the list of species from BiH. *R. virgo* is endemic to the Danube River Basin (Kottelat & Freyhof 2007).

Literature

Brusina 1892; Glowacki 1896; Medić 1902; RHZZM 1908; Ćurčić 1910; Plančić 1923b; Zaplata 1932; Zaplata & Taler 1932; Aganović & Milošević 1959; Smlatić 1961; Vuković 1963a, 1977a; Čanković 1967; Šenk & Aganović 1968; Aganović & Kapetanović 1973; Kosorić *et al.* 1975, 1980; Kosorić 1981; Obratil 1982; Mikavica 1987b; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000, 2002; Mrakovčić *et al.* 2002, 2006; Bogut *et al.* 2006; Škrijelj *et al.* 2006; Kottelat & Freyhof 2007; Pavlović *et al.* 2009; Sofradžija 2009; Drešković *et al.* 2011; Bećiraj & Šahinović 2012; Muhamedagić & Habibović 2013; Čičić-Močić 2014; Delić *et al.* 2014; Milanović *et al.* 2015; Simonović *et al.* 2015; Bećiraj *et al.* 2016; Golub *et al.* 2017b, 2017c, 2018b; Kovačević *et al.* 2018; Nedić *et al.* 2018; Lolić *et al.* 2019; Skenderović & Adrović 2019; Zuliani *et al.* 2019; Dekić *et al.* 2020a, 2024; Skenderović *et al.* 2020b, 2021.

Genus *Scardinius* Bonaparte, 1837

57. *Scardinius dergle* Heckel & Kner, 1858

Vernacular names

Drlja. Dalmatian rudd.

Previous names used for the species

Scardinius hesperdicus Bonaparte, 1845; *Scardinius erythrophthalmus hesperdicus* Heckel, 1851.

Origin and geographic range in BiH

Endemic (ABE). Adriatic Basin.

Remarks

Endemic to the Adriatic Basin in Croatia and the Cetina River catchment in BiH. In BiH, it inhabits karst streams in Livanjsko Polje field, and Buško Blato and Mandečko lakes near Livno.

Literature

Heckel & Kner 1858; Ćurčić 1916; Karaman 1923, 1928; Taler 1953a, Vuković 1963a, 1964, 1977a; Aganović 1969; Vuković *et al.* 1971a, 1971b; Kottelat 1997; Freyhof *et al.* 2005; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008b; Zupančič 2008; Sofradžija 2009; Tutman *et al.* 2012c; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Žujo-Zekić *et al.* 2018; Mušović *et al.* 2020; Sabolić *et al.* 2021; Žujo-Zekić 2021.

58. *Scardinius erythrophthalmus* (Linnaeus, 1758)

Vernacular names

Crvenperka. Rudd.

Previous names used for the species

Leuciscus erythrophthalmus (Linnaeus, 1758); *Scardinius erythrophthalmus erythrophthalmus* (Linnaeus, 1758).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

A widespread European species; in BiH it is present in the Danube River Basin, though it has also been translocated to the Adriatic Basin in the Trebišnjica River system (Bileća Lake).

Literature

Anonymous 1886; Glowacki 1896; Medić 1898; Ćurčić 1910; Plančić 1923b; Aganović 1957; Vuković 1963a, 1977a; Bralo 1967; Čanković 1967; Čanković *et al.* 1968a, 1968b; Kiškarolj & Čanković 1968, 1969; Vuković & Veledar 1974; Vuković 1977a; Guzina & Vuković 1978; Sofradžija *et al.* 1978b; Seratlić & Vuković 1979; Berberović 1980; Habeković *et al.* 1981; Marić 1981; Obratil 1982; Veljović 1982, 1985; Mikavica *et al.* 1991; Pažur 1993; Mikavica & Savić 1999; Radević 2000; Škrijelj *et al.* 2005a, 2005b; Bogut *et al.* 2006; Skenderović *et al.* 2006, 2010, 2011a,

2011b, 2020a, 2020b; Adrović 2007, 2012; Kottelat & Freyhof 2007; Skenderović & Adrović 2008b; Vuković *et al.* 2008; Sofradžija 2009; Golub *et al.* 2013, 2017c; Čičić-Močić 2014; Nedić *et al.* 2014a, 2014b, 2014c, 2014d; Riđanović *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Kovačević *et al.* 2018; Memić 2018; Dekić *et al.* 2020a; Adrović *et al.* 2021; Žujo-Zekić 2021.

59. *Scardinius plotizza* Heckel & Kner, 1858

Vernacular names

Peškelj. Neretva rudd.

Previous names used for the species

Scardinius erythrophthalmus scardafa (Bonaparte, 1837); *Scardinius scardafa* (Bonaparte, 1837); *Scardinius platizza* Heckel, 1845.

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

First mentioned in BiH as *Scardinius erythrophthalmus scardafa* (Bonaparte, 1837) by Vuković (1977a) and Bogut *et al.* (2006). However, according to Kottelat & Freyhof (2007), *S. scardafa* is endemic to Italy. *Scardinius. plotizza* is endemic to the Neretva River catchment in BiH and Croatia (Kottelat & Freyhof 2007); in BiH, it is distributed in the rivers of the Adriatic Basin/Neretva River catchment (lakes on the Neretva River, Hutovo Blato wetland, Krupa River, Deransko Lake); Trebišnjica catchment (Svitava reservoir and the Trebišnjica channel downstream from Trebinje) (Tutman *et al.* 2012c).

Literature

Heckel & Kner 1858; Karaman 1928; Jedlička 1947; Aganović 1952a; Taler 1953a; Ivanišević 1962; Vuković & Ivanišević 1962; Vuković 1963a, 1977a; Kosorić & Vuković 1966a; Guzina *et al.* 1971; Ivanović 1972; Guzina & Vuković 1978; Kosorić 1978; Seratlić *et al.* 1978; Kosorić *et al.* 1983, 1989; Sadović 1986; Economidis & Bănărescu 1991; Crivelli 1996; Kottelat 1997; Glamuzina *et al.* 2001, 2002, 2008a; Tutman *et al.* 2002, 2009b, 2012a, 2012c; Bianco *et al.* 2004; Bianco & Kottelat 2005; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008a, 2008b; Zupančić 2008; Dulčić *et al.* 2009; Prusina *et al.* 2009; Sofradžija 2009; Perea *et al.* 2010; Marčić *et al.* 2012; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Ivanković *et al.* 2017; Benovics *et al.* 2019; Sabolić *et al.* 2021.

Genus *Squalius* Bonaparte, 1837

Several species, including many synonyms and misidentifications, have been reported from the Adriatic Basin of Croatia and BiH, whose taxonomic status including relationships of several populations and their distributional range, has yet to be resolved in detail (Buj *et al.* 2020a).

Squalius albus (Bonaparte, 1838)

Previous names used for the species

Leuciscus cephalus albus Bonaparte 1938.

Remarks

In Vuković (1963a, 1977a), Bogut *et al.* (2006) and Sofradžija (2009), it was mentioned as *Leuciscus cephalus albus* for the waters of the Neretva River catchment. According to Buj *et al.* (2020a) for these reports, the recommended nomenclature is *Squalius squalus*. On the other hand, *S. albus* is still of questionable taxonomic status (Lucentini *et al.* 2014); according to Kottelat & Freyhof (2007), it is endemic to Italy (Lake Trasimeno). Due to the unresolved taxonomic status, *S. albus* it is not included in the BiH checklist.

Literature

See *Squalius squalus*.

60. *Squalius cephalus* (Linnaeus, 1758)**Vernacular names**

Kljen. Chub.

Previous names used for the species

Leuciscus cephalus (Linnaeus, 1758); *Leuciscus cephalus cephalus* (Linnaeus, 1758).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

In Vuković (1963a, 1977a) and Sofradžija (2009), it was mentioned as *Leuciscus cephalus cephalus*, and in Bogut *et al.* (2006) as *Squalius cephalus cephalus*, all for the Danube River Basin in BiH. According to Hamzić (2024), translocated to the Adriatic Basin.

Literature

Heckel & Kner 1858; Anonymous 1886; Brusina 1892; Glowacki 1896; Langhoffer 1904; Plančić 1923b; Karaman 1928; Zaplata & Taler 1932; Kosorić & Kardoš 1955; Aganović & Milošević 1959; Vuković 1959; 1960, 1963a, 1977a, 1982b; Pavlović *et al.* 1960; Vuković *et al.* 1962; Aganović *et al.* 1966; Kosorić & Vuković 1966a; Berberović 1967, 1980; Čanković 1967; Kekić 1967; Sket 1967; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968, 1969; Vuković & Vuković 1968b, 1974; Šenk 1969; Guzina *et al.* 1971, 1981; Ivanović & Sekulović 1971; Vuković & Ivanović 1971; Berberović & Sofradžija 1972; Marić 1972; Aganović & Kapetanović 1973; Bozja & Švob 1974; Kosorić 1974, 1977a, 1978, 1981; Đurović & Vuković 1975; Kosorić *et al.* 1975, 1980, 1983, 1989; Sofradžija 1977; Kaćanski *et al.* 1977; Aganović & Kapetanović 1978; Guzina & Vuković 1978; Habeković *et al.* 1981; Veljović 1982; Šolaja & Pocrnjić 1983b; Kaplanović 1984; Pocrnjić *et al.* 1984; Ratković & Mikavica 1984; Sofradžija *et al.* 1984; Bianco & Knežević 1987; Mikavica 1987b, 1998; Pocrnjić & Šolaja 1988; Economidis & Bănărescu 1991; Mikavica *et al.* 1991, 1997; Pažur 1993; Babić & Mikavica 1994; Bogutskaya & Zupančič 1999; Mikavica & Savić 1997, 1999; Rončević 1997; Radević 2000, 2002; Mrakovčić *et al.* 2002; Korjenić *et al.* 2003; Škrijelj *et al.* 2005a, 2005b, 2006; Tinjak 2005; Bogut *et al.* 2006; Korjenić 2006a, 2006b; Adrović 2007, 2012, 2018; Kottelat & Freyhof 2007; Kurtović *et al.* 2007, 2009; Đurđević *et al.* 2008; Nišić 2008; Skenderović & Adrović 2008b; Škrijelj & Mašović 2008; Šanda *et al.* 2008a, 2008b; Vuković *et al.* 2008; Mitrašinović & Suljević 2009; Skenderović *et al.* 2006, 2010, 2011a, 2011b 2020a, 2020b; Ivanc *et al.* 2011; Bećiraj & Šahinović 2012; Golub *et al.* 2012, 2013, 2016a, 2017b, 2017c, 2018a, 2018b; Ivanc & Dekić 2012; Tutman *et al.* 2012a; Čičić-Močić 2014, 2016, 2018, 2019; Delić *et al.* 2014; Filipović Marijić *et al.* 2014a, 2014b;

Nedić *et al.* 2014a, 2014c, 2014d, 2018; NHMW 2014; Bajrić *et al.* 2015, 2017, 2020; Riđanović *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Bećiraj *et al.* 2016; Mihailović *et al.* 2016; Dekić *et al.* 2017b, 2020a, 2020b; 2024; Benovics *et al.* 2018; Hajdarević *et al.* 2018; Kovačević *et al.* 2018; Memić 2018, 2020; Vesnić *et al.* 2018; Lolić *et al.* 2019; Muhamedagić *et al.* 2019; Zuliani *et al.* 2019; Cvijić *et al.* 2020; Bakrač *et al.* 2020a; Tanović *et al.* 2022, 2023; Hamzić 2024.

***Squalius illyricus* Heckel & Kner, 1858**

Vernacular names

Ilirski klen. Illyrian chub.

Remarks

Hamzić (2024) mentioned this species for BiH, but it is endemic to the Krka River and Cetina River catchments in Croatia only (Čaleta *et al.* 2019) and is therefore not included in the list of fishes from BiH.

61. *Squalius microlepis* Heckel, 1843

Vernacular names

Makal. Makal dace.

Previous names used for the species

Leuciscus microlepis (Heckel, 1843); *Squalius microlepis* Heckel, 1843.

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

Endemic to the Neretva River catchment in BiH and Croatia. Two phenotypically distinct forms of this species inhabit the upper and middle-lower stretches of the Trebižat River. They are largely separated by the subterranean stretch of the river, and is possible that they might represent distinct species (Bogutskaya *et al.* 2019). Further research is required (Perea *et al.* 2010; Geiger *et al.* 2014; Schönhuth *et al.* 2018).

Literature

Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008b; Zupančić 2008; Sofradžija 2009; Perea *et al.* 2010; Drešković *et al.* 2011; Darwall *et al.* 2014; Geiger *et al.* 2014; Milanović *et al.* 2015; Ivanković *et al.* 2017; Benovics *et al.* 2019; Bogutskaya *et al.* 2019; Freyhof *et al.* 2020; Dekić *et al.* 2024.

62. *Squalius squalus* (Bonaparte, 1837)

Vernacular names

Primorski klen, bijeli klen. Cavedano chub.

Previous names used for the species

Leuciscus cephalus cabeda Risso, 1826; *Leuciscus squalus* Bonaparte, 1837; *Leuciscus cephalus albus* Bonaparte, 1838; *Leuciscus cavedanus* (Bonaparte, 1838); *Squalius albus* (Bonaparte, 1838); *Squalius cavedanus* (Bonaparte, 1838).

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

The taxonomic status and distribution range of *Squalius squalus* is still a matter of an ongoing debate (Buj *et al.* 2020a). According to their results, its distribution range is fragmented and encompasses the Istrian rivers and Visovac Lake on the Krka River in Croatia, and Trebišnjica River and the Ravno field in BiH; it was not found in the main course of the Krka or Neretva rivers. Previous literature mentioned *Leuciscus cephalus albus* Bonaparte, 1838 (Vuković 1977a), and *L. cavedanus* (Bonaparte, 1838) (Karaman 1928) in BiH, according to Buj *et al.* (2020a) for these reports, the recommended taxonomy is *S. squalus*.

Literature

Aganović 1952a; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968, 1969; Vuković 1977a; Kosorić *et al.* 1983; Buntić 1985; Economidis & Bănărescu 1991; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Šorić 2007; Šanda *et al.* 2008a, 2008b; Muhamedagić *et al.* 2008; Pavlović *et al.* 2009; Geiger *et al.* 2014; Benovics *et al.* 2019; Buj *et al.* 2020a; Skenderović *et al.* 2020b; Dekić *et al.* 2024.

63. *Squalius svallize* Heckel & Kner, 1858**Vernacular names**

Strugač, sval. Neretva chub.

Previous names used for the species

Leuciscus svallize (Heckel & Kner, 1858); *Leuciscus svallize svallize* (Heckel & Kner, 1858).

Origin and geographic range in BiH

Endemic (NCE BiH, CRO). Adriatic Basin.

Remarks

Endemic to the Neretva River catchment in BiH and Croatia.

Literature

Heckel & Kner 1858; Karaman 1928; Aganović 1952a, 1952c; Taler 1953a; Vuković 1963a, 1977a; Kosorić & Vuković 1966a; Čanković 1967; Kapetanović & Vuković 1968; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968, 1969; Guzina *et al.* 1971; Berberović & Sofradžija 1972; Marić 1972; Švob & Kilalić 1972; Kosorić 1974, 1977a, 1978, Vuković & Vuković 1974; Kačanski *et al.* 1977; Sofradžija 1977; Kosorić *et al.* 1983, 1989; Ratković & Mikavica 1984; Buntić 1985; Bianco & Knežević 1987; Vuković & Sofradžija 1987; Bogutskaya 1994; Crivelli 1996; Mikavica 1998; Bogutskaya & Zupančič 1999; Korjenić *et al.* 2003; Škrijelj *et al.* 2003; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Glamuzina *et al.* 2008a; Jelić *et al.* 2008; Muhamedagić *et al.* 2008; Šanda *et al.* 2008b; Škrijelj & Mašović 2008; Zupančič 2008; Dulčić *et al.* 2009; Pavlović *et al.*

2009; Sofradžija 2009; Ivanković *et al.* 2010, 2011; Perea *et al.* 2010; Drešković *et al.* 2011; Tutman *et al.* 2012a; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Milanović *et al.* 2015; Bećiraj *et al.* 2016; Mrdak & Milošević 2017; Vesnić *et al.* 2018; Buj *et al.* 2020a; Skenderović *et al.* 2020b; Lukač *et al.* 2023; Dekić *et al.* 2024.

64. *Squalius tenellus* Heckel, 1843

Vernacular names

Sitnoļjuskavi klen. Livno masnica.

Previous names used for the species

Leuciscus turskyi tenellus (Heckel, 1843); *Leuciscus turskyi* (Heckel, 1843); *Squalius turskyi* Heckel, 1843.

Origin and geographic range in BiH

Endemic (CCE). Adriatic Basin.

Remarks

Endemic to the Cetina River catchment in BiH, introduced in Croatia from Buško Blato in BiH (Buj *et al.* 2020a). In BiH, the species is rare in the tributaries and canals connected to Buško Blato, whereas a more abundant population is present in Lake Blidinje (Spasojević, unpublished data). Literature data mentioned *Leuciscus turskyi tenellus* (Heckel, 1843) (Aganović *et al.* 1974, 1976) and *Leuciscus turskyi* (Heckel, 1843) (Gilić 1983) in BiH, but according to Buj *et al.* (2020a) recommended taxonomy is *S. tenellus*.

Literature

Heckel & Kner 1858; Ćurčić 1916; Karaman 1923, 1928; Protić 1924; Taler 1953a; Vuković 1963a, 1964, 1977a; Pipić 1966; Berberović 1967; Čanković *et al.* 1968a; Aganović 1969; Pranjić 1969; Vuković & Žnidaršpé-Krzyk 1969; Kaluđerčić *et al.* 1971; Vuković & Ivanović 1971; Berberović & Sofradžija 1972; Marić 1972; Aganović *et al.* 1974, 1976; Vuković & Veledar 1974; Sofradžija 1977; Kaćanski *et al.* 1978; Seratlić & Vuković 1979; Berberović 1980; Gilić 1983; Volić 1984; Guzina *et al.* 1986; Vuković & Sofradžija 1987; Mikavica 1988b; Economidis & Bănărescu 1991; Kottelat 1997; Bănărescu & Herzig-Strachil 1998; Mikavica 1998; Freyhof *et al.* 2005; Freyhof *et al.* 2006, 2020; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Muhamedagić *et al.* 2008; Pavić 2008; Šanda *et al.* 2008b; Zupančić 2008; Perea *et al.* 2010; Darwall *et al.* 2014; Geiger *et al.* 2014; Mustafić & Mrakovčić 2017; Benovics *et al.* 2019; Bogutskaya *et al.* 2019; Buj *et al.* 2020a; Mušović *et al.* 2020; Skenderović *et al.* 2020b; Žujo-Zekić 2021; Carosi *et al.* 2024; Dekić *et al.* 2024.

Genus *Telestes* Bonaparte, 1837

65. *Telestes dabar* (Bogutskaya, Zupančić, Bogut & Naseka, 2012).

Vernacular names

Dabarska gaovica. Dabar minnow.

Previous names used for the species

Paraphoxinus metohiensis Steindachner, 1901; *Phoxinellus metohiensis* (Steindachner, 1901); *Telestes metohiensis* (Steindachner, 1901).

Origin and geographic range in BiH

Endemic (NCE, BiH). Adriatic Basin.

Remarks

Described based on anatomical comparisons of isolated geographic populations of specimens previously identified as *Telestes metohiensis* (Bogutskaya *et al.* 2012). However, according to molecular data, some specimens identified as *T. dabar* are actually *T. metohiensis* (Buj *et al.* 2017). Furthermore, Reier *et al.* (2022) recognized both species as distinct based on molecular data. Occurs only in the underground waters of Dabarsko Polje (Eastern Herzegovina) (Bogutskaya *et al.* 2012).

Literature

Bogutskaya *et al.* 2012; Dekić *et al.* 2012a; Geiger *et al.* 2014; Lolić *et al.* 2015; Buj *et al.* 2017; Benovics *et al.* 2019; Francuski *et al.* 2019; Freyhof *et al.* 2020; Reier *et al.* 2022; Lukač *et al.* 2023.

66. *Telestes metohiensis* (Steindachner, 1901)

Vernacular names

Gatačka gaovica. Gacko minnow.

Previous names used for the species

Paraphoxinus metohiensis Steindachner, 1901; *Phoxinellus metohiensis* (Steindachner, 1901).

Origin and geographic range in BiH

Endemic (NCE, BiH). Adriatic Basin.

Remarks

In earlier literature, it was mentioned as *Paraphoxinus metohiensis* (Vuković 1977a), *Phoxinellus metohiensis* (Sofradžija 2009), and as *Telestes metohiensis* in Bogutskaya *et al.* (2012); endemic to the Neretva River catchment in BiH.

Literature

Steindachner 1901; Trgovčević 1905; Ćurčić 1915a, 1915b; Taler 1953a, 1953b; Vuković 1963a, 1977a, 1982b; Bašić 1964; Sabioncello 1967; Berberović *et al.* 1969b, 1970d; Vuković & Miladinović 1969; Aganović & Kapetanović 1970, 1971; Guzina & Vuković 1972; Miladinović *et al.* 1971; Vuković & Ivanović 1971; Karaman 1972; Sofradžija & Berberović 1972; Bozja & Švob 1974; Vuković & Vuković 1974; Pavlović & Berberović 1978; Pocrnjić *et al.* 1979; Leiner 1984; Vuković & Sofradžija 1987; Povž *et al.* 1990; Šorić 1992; Crivelli 1996; Kottelat 1997; Zupančič & Bogutskaya 2000, 2002; Bogutskaya & Zupančič 2003; Ketmaier *et al.* 2004; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Jelić *et al.* 2008; Šanda *et al.* 2008b; Zupančič 2008; Pavlović *et al.* 2009; Sofradžija 2009; Palandačić *et al.* 2010; Perea *et al.* 2010; Dekić *et al.* 2011, 2012a, 2012b, 2013a, 2013b, 2024; Drešković *et al.* 2011; Bogutskaya *et al.* 2012; Darwall *et al.* 2014; Geiger *et al.* 2014; Lolić *et al.* 2015; Milanović *et al.* 2015; Mandić *et al.* 2016; Buj *et al.* 2017; Benovics *et al.* 2019; Francuski *et al.* 2019; Freyhof *et al.* 2020; Reier *et al.* 2022; Lukač *et al.* 2023.

Telestes polylepis Steindachner, 1866

Vernacular names

Croatian riffle dace.

Remarks

Hamzić (2024) recorded this species in BiH, but it is endemic only to the karstic fields in the Lika region in Croatia (Čaleta *et al.* 2019) and is therefore not included in the list of fishes of BiH.

67. *Telestes souffia* (Risso, 1827)

Vernacular names

Jelšovka. Vairone, Souffia.

Previous names used for the species

Leuciscus souffia Risso, 1826; *Leuciscus agassizii* Valenciennes, 1844; *Leuciscus souffia agassizii* Valenciennes, 1844.

Origin and geographic range in BiH

Native. Danube River Basin.

Remarks

Vuković (1963a) mentioned the species as *Telestes agassizii*, Vuković (1977a) for BiH, and Sofradžija (2009) as *Leuciscus souffia agassizii*, but all are now considered junior synonyms of *T. souffia* (Risso, 1827).

Literature

Anonymous 1886; Glowacki 1896; Vuković 1963a, 1963b, 1977a, 1979, 1982b, 1986, 1987, 1990; Vuković & Ivanović 1971; Vuković & Vuković 1974; Sofradžija 1977; Guzina & Vuković 1978; Mikavica *et al.* 1991; Korjenić 1998; Mikavica 1998; Mikavica & Savić 1999; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Gilles *et al.* 2009; Sofradžija 2009; Drešković *et al.* 2011; Milanović *et al.* 2015; Simonović *et al.* 2015; Dekić *et al.* 2024.

Telestes turskyi (Heckel, 1843)

Vernacular names

Turski klen. Turski dace.

Previous names used for the species

Leuciscus turskyi tenellus (Heckel, 1843); *Leuciscus turskyi* (Heckel, 1843); *Squalius turskyi* Heckel, 1843.

Remarks

Endemic to the Krka River catchment in Croatia (Buj *et al.* 2022). Sofradžija (2009) stated that the species inhabits BiH freshwaters, but this is likely based on misidentification of specimens. Therefore, it is not included in this checklist.

Literature

Vuković & Ivanović 1971; Mikavica 1988b; Mrakovčić *et al.* 2006; Sofradžija 2009.

Telestes ukliva (Heckel, 1843)

Vernacular names

Ukliva.

Remarks

Hamzić (2024) recorded this species in BiH, but it is endemic only to the Cetina River catchment in Croatia (Čaleta *et al.* 2019). Therefore, the species is not included in our checklist either.

Genus *Vimba* Fitzinger, 1873

68. *Vimba vimba* (Linnaeus, 1758)

Vernacular names

Šljivar, nosara. Vimba.

Previous names used for the species

Abramis melanops (Heckel, 1837); *Abramis vimba* (Linnaeus, 1758); *Vimba vimba carinata* (Pallas, 1811).

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Glowacki 1896; RHZZM 1908; Ćurčić 1910; Plančić 1923b; Zaplata & Taler 1932; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Berberović 1980; Kosorić *et al.* 1980; Kačanski *et al.* 1981; Veljović 1982, 1985; Mikavica 1987b; Mikavica *et al.* 1991; Leiner 1998; Mikavica & Savić 1999; Radević 2000, 2002; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Čičić-Močić 2014, 2016, 2018; Nedić *et al.* 2014a, 2014c, 2014d, 2018; Riđanović *et al.* 2015; Simonović *et al.* 2015; Skenderović & Adrović 2019; Dekić *et al.* 2020a, 2024; Skenderović *et al.* 2020b, 2021.

Order Siluriformes Hay, 1929

Family Siluridae Cuvier, 1816

Genus *Silurus* Linnaeus, 1758

69. *Silurus glanis* Linnaeus, 1758

Vernacular names

Som. European catfish.

Remarks

Translocated to the Adriatic Basin (Glamuzina *et al.* 2017).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Literature

Anonymous 1886, 1931, 1947b; Brusina 1892; Glowacki 1896; Ribić 1901; Ćurčić 1910; Plančić 1923b; Ćurčić 1933; Bek 1935; Jedlička 1935; Taler 1945b; Smlatić 1961; Vuković 1963a, 1977a; Kiškarolj 1965; Čanković *et al.* 1968a, 1968b, 1968c; Munjko 1972; Kosorić *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Sofradžija 1982b, 2009; Veljović 1982, 1985; Delić 1984; Mikavica 1987b; Mikavica *et al.* 1991; Pažur 1993; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002; Mateš 2004; Bogut *et al.* 2006; Adrović 2007; 2012; Kottelat & Freyhof 2007; Suić *et al.* 2008; Šanda *et al.* 2008b; Vuković *et al.* 2008; Sofradžija 2009; Tutman *et al.* 2012a; Golub *et al.* 2013, 2017c, 2018b; Čičić-Močić 2014; Delić *et al.* 2014; Nedić *et al.* 2014a, 2014c, 2014d, 2016b, 2018; Riđanović *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Glamuzina *et al.* 2017; Kovačević *et al.* 2018; Memić 2018; Mušović *et al.* 2018, 2020; Žujo-Zekić *et al.* 2018; Dekić *et al.* 2020a, 2024; Skenderović *et al.* 2020b, 2021; Žujo-Zekić 2021; Hamzić 2024.

Family Ictaluridae

Genus *Ameiurus* Rafinesque, 1820

70. *Ameiurus melas* (Rafinesque, 1820)

Vernacular names

Crni somić. Black bullhead.

Previous names used for the species

Ictalurus melas (Rafinesque, 1820).

Origin and geographic range in BiH

Non-native. Danube River Basin.

Remarks

Non-native species introduced from North America whose distribution in BiH is not yet known because of possible misidentifications of specimens of *Ameiurus nebulosus*. Recently, it was also mentioned as *A. melas* for the artificial lake Modrac (Halilović & Adrović 2015).

Literature

Bogut *et al.* 2006; Šanda *et al.* 2008b, 2009; Sofradžija 2009; Tutman *et al.* 2012a; Halilović & Adrović 2015; Halilović & Adrović 2015; Simonović *et al.* 2015; Skenderović *et al.* 2016, 2020b; Glamuzina *et al.* 2017; Memić 2018; Nedić *et al.* 2018; Piria *et al.* 2018; Dekić *et al.* 2024.

71. *Ameiurus nebulosus* (Lesueur, 1819)

Vernacular names

Američki somić. Brown bullhead.

Previous names used for the species

Ictalurus nebulosus (Lesueur, 1819).

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Non-native species introduced from North America. Reported from both the Adriatic and Danube River Basin in BiH, but its distribution is yet to be confirmed because of possible misidentifications of specimens of *Ameiurus melas*.

Literature

Vuković 1963a, 1977a; Kiškarolj 1965; Čanković *et al.* 1968a, 1968c; Žitnan *et al.* 1969; Obradović 1974; Kosorić 1978; Grbelja *et al.* 1980; Nadaždin *et al.* 1980; Obratil 1982; Veljović 1982; Delić 1984; Leiner 1998; Radević 2000; Bogut *et al.* 2006; Adrović 2007, 2012; Adrović & Skenderović 2007; Šanda *et al.* 2008b, 2009; Glamuzina *et al.* 2008a, 2017; Vuković *et al.* 2008; Sofradžija 2009; Šukalo *et al.* 2012; Tutman *et al.* 2012a; Golub *et al.* 2013; Mikavica *et al.* 2013; Čičić-Močić 2014; Nedić *et al.* 2014a, 2014c, 2014d, 2016b, 2018, 2021; Riđanović *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Kovačević *et al.* 2018; Piria *et al.* 2018; Skenderović & Adrović 2019; Dekić *et al.* 2020a, 2024; Mušović *et al.* 2020; Skenderović *et al.* 2020b; Žujo-Zekić 2021.

Genus *Ictalurus* Rafinesque, 1820

Ictalurus punctatus Rafinesque, 1818

Vernacular names

Kanalski som. Channel catfish.

Origin and geographic range in BiH

Non-native. Danube River Basin.

Remarks

Non-native species introduced to Europe from North America. Literature records in BiH (Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009) are still dubious (Hamzić 2024), and it is not clear whether the species has established a population in the region. Therefore, it is not included in this checklist.

Literature

Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009; Hamzić 2024.

Order Salmoniformes Bleeker, 1859

Family Esocidae Cuvier, 1817

Genus *Esox* Linnaeus, 1758

72. *Esox lucius* Linnaeus, 1758

Vernacular names

Štuka. Northern Pike.

Previous names used for the species

Esox lucioides Agassiz & Girard, 1850; *Lucius lucius* (Linnaeus, 1758).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to the Adriatic Basin in Hutovo Blato wetland (Glamuzina *et al.* 2017).

Literature

Heckel & Kner 1858; Anonymous 1886; Brusina 1892; Glowacki 1896; Anonymous 1909; Ćurčić 1910; Plančić 1923b; Ćurčić 1933; Bek 1935; Jedlička 1935; Taler 1945b; Smlatić 1961; Vuković 1963a, 1977a; Kiškarolj 1965; Čanković 1967; Čanković *et al.* 1968a, 1968b, 1968c; Munjko 1972; Kosorić *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Veljović 1982, 1985; Delić 1984; Mikavica *et al.* 1991; Pažur 1993; Leiner 1998; Mikavica & Savić 1999; Radević 2000; Simonović 2001; Mrakovčić *et al.* 2002; Skenderović *et al.* 2006; Adrović 2007, 2012; Kottelat & Freyhof 2007; Suić *et al.* 2008; Vuković *et al.* 2008; Sofradžija 2009; Skenderović *et al.* 2011a, 2020a, 2020b; Bećiraj & Šahinović 2012; Tutman *et al.* 2012a; Golub *et al.* 2013, 2017c, 2018a, 2018b; Čičić-Močić 2014, 2016; Delić *et al.* 2014; Nedić *et al.* 2014a, 2014c, 2014d, 2016b, 2018; Riđanović *et al.* 2015; Simonović *et al.* 2015; Bećiraj *et al.* 2016; Glamuzina *et al.* 2017; Kovačević *et al.* 2018; Memić 2018; Skenderović & Adrović 2019; Zuliani *et al.* 2019; Dekić *et al.* 2020a, 2024; Adrović *et al.* 2021.

Family Umbridae Bonaparte, 1845
Genus *Umbra* Kramer, 1777

73. *Umbra krameri* Walbaum, 1792

Vernacular names

Crnka. European mudminnow.

Previous names used for the species

Umbra krameri Fitzinger, 1832; *Umbra canina* Károli, 1882.

Origin and geographic range in BiH

Native. Danube River Basin.

Remarks

In BiH, there are few confirmed records in the Gromiželj swamp near Bijeljina (Sekulić *et al.* 2013) and the lower Matura River system (Marić *et al.* 2019).

Literature

Vuković 1963a, 1977a; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Sofradžija 2009; Drešković *et al.* 2011; Sekulić *et al.* 2013; Marić *et al.* 2015, 2017, 2019; Lukić *et al.* 2016; Ćurčić 2017; Čolić 2018; Dekić *et al.* 2024.

Family Salmonidae Cuvier, 1816
Genus *Coregonus* Linnaeus, 1758

73. *Coregonus peled* (Gmelin, 1789)

Vernacular names

Sjeverna ozimica. Peled.

Previous names used for the species

Coregonus oxyrinchus (Linnaeus, 1758).

Origin and geographic range in BiH

Non-native. Adriatic Basin.

Remarks

Non-native species introduced in several water bodies in the past, but the recent distribution is not well known. Hamzić *et al.* (2011) collected several specimens in Boračko Lake near Konjic during August 2011.

Literature

Simonović 2001; Hamzić *et al.* 2011; Piria *et al.* 2018.

Genus *Hucho* Günther, 1866

75. *Hucho hucho* (Linnaeus, 1758)

Vernacular names

Mladica. Huchen.

Previous names used for the species

Salmo hucho Linnaeus, 1758.

Origin and geographic range in BiH

Endemic (DBE). Danube River Basin.

Remarks

Huchen is endemic to the Danube River Basin, where it inhabits streams and rivers with fast water flow. It is one of the largest salmonid fish of the world and one of the most endangered species inhabiting the Danube River Basin of Central Europe (IUCN 2025). Some populations are supported by artificial reproduction and restocking programmes.

Literature

Anonymous 1886, 1903; Kesterčanek 1886; Brusina 1892; Hawlitschek 1895; Glowacki 1896; Mojsisovics 1897; Medić 1901; Horvat 1901; Hirc 1902; Ćurčić 1910; Plančić 1923b; Zaplata & Taler 1932; Protić 1933; Taler 1945a, 1951b; Hafner-Lahorski 1947, 1948; Vuković Teslić 1950; Kosorić & Kardoš 1955; Bogdanović 1957; Rukavina & Delić 1957; Smlatić 1961; Vuković 1963a, 1977a, 1982a; Sabioncello 1967; Čanković 1967; Čanković *et al.* 1968a; Aganović & kapetanović 1973; Kosorić *et al.* 1975; Sofradžija 1979; Kosorić 1981; Sofradžija & Hadžiselimović 1981b; Pažur *et al.* 1982; Imamović 1986; Mikavica 1987a, 1987b; 1988a, Skalin 1990; Mikavica *et al.* 1991; Pažur 1993; Babić & Mikavica 1994; Mikavica & Savić 1999; Radević 1999, 2000; Simonović *et al.* 2000, 2011, 2015; Georgiev 2003; Mateš 2004, 2008; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Škrijelj *et al.* 2006; Kottelat & Freyhof 2007; Pavlović *et al.* 2009; Sofradžija 2009; Ajanović *et al.* 2010; Popović 2010; Drešković *et al.* 2011; Weiss *et al.* 2011; Muhamedagić & Habibović 2013; Witkowski *et al.* 2013; Geiger *et al.* 2014; Freyhof *et al.* 2015, 2020; Bećiraj *et al.* 2016; Čičić-Močić 2016, 2018; Lukić *et al.* 2016; Weiss & Schenekar 2016; Glamuzina *et al.* 2017; Kovačević *et al.* 2018; Piria *et al.* 2018; Lolić *et al.* 2019; Skenderović *et al.* 2020b; Savić 2022; Dekić *et al.* 2020a, 2024.

Genus *Oncorhynchus* Suckley, 1861

76. *Oncorhynchus mykiss* (Walbaum, 1792)

Vernacular names

Kalifornijska pastrva. Rainbow trout.

Previous names used for the species

Parasalmo gairdneri (Richardson, 1836); *Salmo irideus* Gibbons, 1855; *Salmo gairdneri* Richardson, 1836.

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Non-native species introduced from North America in 1902 to the hatchery at Vrelo Bosne (Taler 1954). It can likely be found in all major watercourses in BiH where there are salmonid ponds. Usually introduced for the restocking of rivers and reservoirs in the sport fishery sector (Glamuzina *et al.* 2017). There is no published information on whether this species reproduces naturally in BiH.

Literature

Zaplata & Taler 1932; Mršić 1935; Vuković Teslić 1950; Taler 1953c, 1954; Bogdanović 1954; Aganović 1957; Aganović & Kapetanović 1963; Vuković 1963a, 1977a; Čanković 1967; Čanković *et al.* 1968a, 1970, 1971; Kiškarolj & Čanković 1969; Kaluđerčić *et al.* 1971; Kosorić *et al.* 1971; Guzina *et al.* 1977; Vuković & Kosorić 1978; Vicković 1979; Sofradžija 1982a; Imamović 1986, 1987; Kosorić *et al.* 1989; Mikavica & Grujić 1995a, 1995b; Radević 2000; Korjenić *et al.* 2003; Škrijelj *et al.* 2003; Bogut *et al.* 2006; Adrović & Skenderović 2007; Kottelat & Freyhof 2007; Muhamedagić *et al.* 2008; Šanda *et al.* 2008b, 2009; Sofradžija 2009; Dreca 2009; Hasković *et al.* 2011; Ivanc *et al.* 2011; Bećiraj & Šahinović 2012; Serdarević 2012; Simonović *et al.* 2015; Stanković *et al.* 2015, 2016; Pavličević *et al.* 2016; Bećiraj *et al.* 2016; Čičić-Močić 2016, 2018; Mrdak & Milošević 2017; Glamuzina *et al.* 2017; Adrović 2018; Keško 2018; Kovačević *et al.* 2018; Memić 2018; Piria *et al.* 2018; Savić *et al.* 2018; Mušović *et al.* 2020; Skenderović *et al.* 2020b; Žujo-Zekić 2021; Neuburg *et al.* 2023; Carosi *et al.* 2024; Dekić *et al.* 2024.

Genus *Salmo* Linnaeus, 1758

Salmo dentex (Heckel, 1852)

Vernacular names

Zubatak. Dentex trout.

Previous names used for the species

Salar dentex Heckel, 1851; *Salmo trutta dentex* Heckel, 1851.

Origin and geographic range in BiH

Adriatic Basin.

Remarks

Based on the results of Snoj *et al.* (2010), it is possible that this species represents a form in the life cycle of *Salmo marmoratus* in the Neretva River catchment and of *S. trutta* in the Skadar Lake catchment. Results of Snoj *et al.* (2010) clearly indicate that *S. dentex* does not represent a monophyletic lineage and should not be considered a distinct species. Therefore, it is not included in the list.

Literature

Heckel 1852; Ćurčić 1938; Taler 1953a; Vuković 1963a, 1977a; Kosorić & Vuković 1966a; Kosorić *et al.* 1983, 1989; Vuković & Ivanović 1971; Kačanski *et al.* 1977; Mikavica 1998; Hamzić 2002; Bogut *et al.* 2006; Glamuzina & Bartulović 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Razpet *et al.* 2007; Has-Schön *et al.* 2008; Tutman *et al.* 2008a, 2012a; Zupančič 2008; Dulčić *et al.* 2009; Glamuzina *et al.* 2009; Sofradžija 2009; Snoj *et al.* 2010; Skenderović *et al.* 2020b.

77. *Salmo farioides* Karaman 1938**Vernacular names**

Primorska pastrva. Balkan brook trout.

Previous names used for the species

Salmo trutta farioides.

Origin and geographic range in BiH

Endemic to Adriatic and Ionian Basins.

Remarks

Endemic to the Adriatic Basin in BiH, Croatia, Serbia, Montenegro, Albania and Republic of Northern Macedonia and Ionian Basin in Albania and Greece (Kottelat & Freyhof 2007). The taxonomic status of *Salmo farioides* has always been questionable due to inadequate original descriptions (Kottelat 1997; Razpet *et al.* 2007).

Literature

Kottelat 1997; Hamzić 2002; Korjenić 2005b; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Razpet *et al.* 2007; Šanda *et al.* 2008a, 2008b, 2009; Zupančič 2008; Sofradžija 2009; Suljević 2012; Tutman *et al.* 2012a; Geiger *et al.* 2014; Žujo-Zekić *et al.* 2018, Žujo-Zekić 2021; Carosi *et al.* 2024.

78. *Salmo labrax* Pallas, 1814**Vernacular names**

Crnomorska pastrva. Black Sea trout.

Previous names used for the species

Salmo trutta labrax Pallas, 1814.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Some authors considered that *Salmo labrax* is just the Danube lineage of the *S. trutta* complex (Pustovrh *et al.* 2014a, 2014b), while others consider it as a distinct species (Kottelat 1997; Kottelat & Freyhof 2007). According to Hamzić (2024), translocated to the Adriatic Basin.

Literature

Kottelat & Freyhof 2007; Ajanović *et al.* 2010; Škraba *et al.* 2017; Škraba Jurlina *et al.* 2020; Marić *et al.* 2022; Hamzić 2024.

79. *Salmo letnica* (Karaman 1924)

Vernacular names

Ohridska pastrva. Ohrid trout.

Origin and geographic range in BiH

Non-native. Adriatic Basin.

Remarks

Endemic to Ohrid Lake, with still unclear taxonomic status (Sušnik *et al.* 2007a). Non-native species that was introduced in 1987 to the hatchery in Blagaj and in the same year stocked in Boračko Lake and Plivska Lakes where it spawns successfully (Hamzić *et al.* 2012a; Hamzić 2024).

Literature

Hamzić *et al.* 2012a; Hamzić 2024.

80. *Salmo marmoratus* Cuvier, 1829

Vernacular names

Glavatica. Marble trout.

Previous names used for the species

Salmo trutta marmoratus Cuvier, 1829; *Trutta fario marmorata* Siebold, 1863; *Trutta adriatica* Kolombatović, 1890.

Origin and geographic range in BiH

Endemic (ABE). Adriatic Basin.

Remarks

Endemic to the Adriatic Basin (Vuković 1977a; Bogut *et al.* 2006), mainly known from the Soča River in Slovenia. Native species with still unclear taxonomic status, as some genetic and phenotypic differences between populations from the Soča and Neretva Rivers were recently discovered (Razpet *et al.* 2007). Kottelat & Freyhof (2007) proposed this species as a junior synonym of *Salmo montenigrinus*. Due to the introduction of highly competitive non-native species and highly unregulated stocking, this species, if valid, is extremely endangered in BiH (Hamzić 2024).

Literature

Heckel & Kner 1858; Steindachner 1882b; Heintz 1908; Karaman 1927, 1938; Ćurčić 1938; Taler 1945a, 1950; 1951b, 1953a, 1953b; Aganović 1952a; Vuković 1963a, 1977a, 1982a; Landiges & Vogt 1965; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968; Kosorić 1969, 1972a, 1974, 1977a, 1978; Dorofejeva & Seratlić 1970; Tortonese 1970; Guzina *et al.* 1971; Kaluđerčić *et al.* 1971; Kosorić & Vuković 1971; Dorofejeva & Seratlić-Savić 1972; Vuković & Vuković 1977; Kaćanski *et al.* 1977; Vuković *et al.* 1977; Aganović & Kapetanović 1978; Kosorić *et al.* 1983, 1989; Ratković & Mikavica 1984; Vuković & Sofradžija 1987; Povž *et al.* 1990; Mikavica 1998; Hamzić 2002; Durmić-Pašić *et al.* 2005, 2007a; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Razpet *et al.* 2007; Durmić-Pašić 2008; Muhamedagić *et al.* 2008, 2019; Zupančić 2008; Drešković *et al.* 2011; Pustovrh *et al.* 2011, 2014a, 2014b; Tutman *et al.* 2012a; Skenderović *et al.* 2020b; Dekić *et al.* 2024; Hamzić 2024.

81. *Salmo montenigrinus* (Karaman, 1933)

Vernacular names

Crnogorska pastrva. Montenegro trout.

Origin and geographic range in BiH

Endemic (ABE). Adriatic Basin.

Remarks

Endemic to the Adriatic Basin of Montenegro and BiH. In BiH, it is present in the southern part of the country (Kottelat & Freyhof 2007). In the 1980s and 1990s, this species was stocked in the Trebišnjica River catchment with fry from the “Mareza” and “Morača” fish farms (Hamzić 2024).

Literature

Kottelat & Freyhof 2007; Hamzić 2024.

82. *Salmo obtusirostris* (Heckel, 1851)

Vernacular names

Mekousna pastrva. Softmouth trout, Soft-muzzled trout.

Previous names used for the species

Salmothymus obtusirostris Heckel, 1851; *Thymallus microlepis* Steindachner, 1874; *Salmothymus obtusirostris oxyrhynchus* Steindachner, 1882.

Origin and geographic range in BiH

Endemic (ABE). Adriatic Basin.

Remarks

First reported in BiH as *Salmo obtusirostris oxyrhynchus* (Vuković 1963a, 1977a). Endemic to the rivers of the southeastern Adriatic Basin (Kottelat & Freyhof 2007). It is recorded in the Krka, Jadro, Žrnovnica and Vrljika Rivers in Croatia, the Neretva River in BiH and Zeta and Morača Rivers in Montenegro (Glamuzina *et al.* 2018).

Literature

Zaplata & Taler 1932; Ćurčić 1938; Taler 1950, 1951b; Rukavina & Delić 1959; Janković 1961; Vuković 1963a, 1977a, 1982a; Kosorić & Vuković 1966a; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968; Rukavina & Ćorić 1968; Šenk & Kosorić 1968; Kiškarolj & Čanković 1969; Šenk 1969; Berberović *et al.* 1970e; Čanković *et al.* 1970, 1971; Guzina *et al.* 1971; Kaluđerčić *et al.* 1971; Kosorić *et al.* 1971; Vuković & Ivanović 1971; Korovina & Vuković 1972; Švob & Kilalić 1972; Kosorić 1974, 1977a, 1977b; Kaćanski *et al.* 1977; Vuković & Vuković 1977; Aganović & Kapetanović 1978; Vuković & Kosorić 1978; Kosorić *et al.* 1983, 1989; Vuković & Sofradžija 1987; Economidis & Bănărescu 1991; Babić & Mikavica 1994; Kottelat 1997; Mikavica 1998; Mikavica *et al.* 2001, 2002; Hamzić 2002; Snoj *et al.* 2002a, 2002b, 2010; Durmić-Pašić *et al.* 2005, 2007a, 2007b; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sušnik *et al.* 2007b; Razpet *et al.* 2007; Jelić *et al.* 2008; Muhamedagić *et al.* 2008, 2019; Šanda *et al.* 2008a, 2008b, 2009; Zupančić 2008; Pavlović *et al.* 2009; Drešković *et al.* 2011; Tutman *et al.* 2012a, 2023; Darwall *et al.* 2014; Delić *et al.* 2014; Esteve *et al.* 2014; Geiger *et al.* 2014; Pustovrh *et al.* 2014a, 2014b; Milanović *et al.* 2015; Glamuzina *et al.* 2017, 2018; Mrdak & Milošević 2017; Freyhof *et al.* 2020; Dekić *et al.* 2020b, 2024; Skenderović *et al.* 2020b; Neuburg *et al.* 2023.

Salmo taleri (Karaman, 1933)

Vernacular names

Zetska pastrva.

Remarks

According to Kottelat & Freyhof (2007) this species inhabits only waters of Montenegro. However, Schöffmann & Marić (2024) suggested this species is widespread in the Sava River catchment. Because the occurrence and distribution of this species are unclear, it is not included in the list of fishes from BiH, awaiting further information.

Literature

Vuković 1963a, 1977a; Hamzić 2024.

83. *Salmo trutta* Linnaeus, 1758

Vernacular names

Pastrmka. Brown trout.

Previous names used for the species

Salmo fario Linnaeus, 1758; *Salmo trutta m. fario* Linnaeus, 1758; *Salmo trutta m. lacustris* Linnaeus, 1758; *Trutta fario* (Linnaeus, 1758).

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Native brown trout from the Mediterranean Basin can be assigned to three lineages, Adriatic (AD), Mediterranean (ME), and *marmoratus* (MA), while those from the Danube Basin belong to the Danubian (DA) lineage (Bernatchez *et al.* 1992; Bernatchez 2001). Numerous different sublineages are also

recognized within the AD, ME, and DA lineages (Kottelat & Freyhof 2007; Segherloo *et al.* 2021), and in general, the taxonomic status of many of these species remains unclear. Based on previous reports, *Salmo trutta* is not native to BiH (Kottelat & Freyhof 2007), but it was often used for stocking the rivers of the Danube and the Adriatic Basins in Croatia and BiH (Buj *et al.* 2020b). The local native genetic pool has recently been compromised with the import of fertilized eggs of different origins (i.e., Atlantic lineage), as reared juveniles are widely used for restocking activities. *Salmo trutta* in the Neretva River catchment does not form a genetically unified assemblage. The assemblages are not correlated with phenotype but rather with sampling location. Thus, the most distinctive brown trout associations were observed in relation to either remote (supposedly unstocked) or more urban (supposedly stocked) locations (Razpet *et al.* 2007; Kalamujić 2013). However, according to Neuburg *et al.* (2023), brown trout in the upper Neretva form a genetically homogeneous unit that is clearly distinct from other geographically separated populations such as the brown trout lineages from the Danube and Aegean Basins and the Atlantic lineage (introduced into the Neretva system).

Literature

Brusina 1892; Glowacki 1896; Hirc 1900; Langhoffer 1904; Anonymous 1909; Wisniewski 1932; Zaplata & Taler 1932; Protić 1933; Ćurčić 1938; Taler 1950, 1951b, 1951d; Vuković Teslić 1950; Aganović 1952a, 1952b, 1957, 1967; Kosorić & Kardoš 1955; Šinžar 1955; Šenk 1954, 1956b, 1969; Bogdanović 1957; Aganović & Milošević 1959; Rukavina & Delić 1959; Pavlović *et al.* 1960, 1962; Vuković 1963a, 1977a, 1982a; Smlatić 1961; Cicović 1964; Kosorić & Vuković 1966a; Čanković 1967; Cerović & Ivanović 1968; Čanković *et al.* 1968a; Kiškarolj & Čanković 1968, 1969; Rukavina & Ćorić 1968; Šenk & Aganović 1968; Rojnik 1969; Kačanski & Kosorić 1970; Guzina *et al.* 1971; Kaluđerčić *et al.* 1970, 1971; Kosorić *et al.* 1971, 1980, 1983, 1989; Švob & Kilalić 1972; Aganović & Kapetanović 1973, 1978; Kosorić 1974, 1977a, 1978, 1981; Kačanski *et al.* 1977, 1978; Vuković & Kosorić 1978; Sofradžija 1982a, 2009; Ratković & Mikavica 1984; Imamović 1986, 1987; Kosorić & Mikavica 1986; Mikavica 1988a; Economidis & Bănărescu 1991; Mikavica *et al.* 1991; Babić & Mikavica 1994; Mikavica & Savić 1999; Radević 2000; Hamzić 2002; Trožić-Borovac 2002; Korjenić 2003, 2004b, 2005b, 2007; Korjenić *et al.* 2003, 2009; Škrijelj *et al.* 2003, 2006; Alić *et al.* 2004; Bogunić 2005; Durmić-Pašić *et al.* 2005, 2007a; Kadribegović 2005; Trožić-Borovac *et al.* 2005; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Pojskić *et al.* 2007; Razpet *et al.* 2007; Muhamedagić *et al.* 2008, 2019; Sofradžija 2009; Mišić 2011; Bećiraj & Šahinović 2012; Golub *et al.* 2012; Ivanc & Dekić 2012; Suljević 2012; Tutman *et al.* 2012a; Čorbo 2013; Kalamujić 2013; Muhamedagić & Habibović 2013; Kerkez *et al.* 2014; NHMW 2014; Pustovrh *et al.* 2014a, 2014b; Spasojević 2014; Pojskić & Kalamujić 2015; Simonović *et al.* 2015, 2017; Bećiraj *et al.* 2016; Čičić-Močić 2016, 2018, 2019; Dekić *et al.* 2016b, 2017b, 2020a, 2020b, 2024; Pavličević *et al.* 2014, 2016; Škraba *et al.* 2017; Veladžić *et al.* 2017; Adrović 2018; Keško 2018; Kovačević *et al.* 2018; Memić 2018; Nikolić *et al.* 2018; Piria *et al.* 2018; Lolić *et al.* 2019; Mušović *et al.* 2020; Skenderović *et al.* 2020a, 2020b; Škraba Jurlina *et al.* 2020; Neuburg *et al.* 2023; Carosi *et al.* 2024.

Genus *Salvelinus* Richardson, 1836

84. *Salvelinus alpinus* (Linnaeus, 1758)

Vernacular names

Jezerska zlatovčica. Arctic charr.

Previous names used for the species

Salmo alpinus Linnaeus, 1758.

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Non-native species introduced in BiH for aquaculture in both Adriatic and Danube basins, and later also used for restocking rivers and reservoirs (Hamzić 2024).

Literature

Zaplata & Taler 1932; Vuković Teslić 1950; Aganović 1957, 1967; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Vuković & Kosorić 1978; Imamović 1986; Sofradžija 2009; Glamuzina *et al.* 2017; Piria *et al.* 2018; Skenderović *et al.* 2020b; Dekić *et al.* 2024; Hamzić 2024.

85. *Salvelinus fontinalis* Mitchell, 1814

Vernacular names

Potočna zlatovčica. Brook trout.

Previous names used for the species

Salmo fontinalis Mitchell, 1814.

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Introduced in BiH for aquaculture purposes in both the Adriatic and Danube basins, and later also used for restocking rivers and reservoirs (Hamzić 2024). Recent distribution and the abundance of the species is not known.

Literature

Zaplata & Taler 1932; Taler 1950; Vuković 1963a, 1977a; Čanković 1967; Čanković *et al.* 1968a; Vuković & Kosorić 1978; Kosorić 1981; Radević 2000; Muhamedagić *et al.* 2008, Šanda *et al.* 2008b, 2009; Sofradžija 2009; Stanić 2013; Glamuzina *et al.* 2017; Piria *et al.* 2018; Kalamujić Stroil *et al.* 2020; Skenderović *et al.* 2020b; Neuburg *et al.* 2023; Dekić *et al.* 2024; Hamzić 2024.

Salvelinus salvelinus (Linnaeus, 1758)

Remarks

Vuković (1963a) reported the occurrence of the species in the Plivsko Lake, Danube River Basin, where it was introduced. According to Hamzić (2024), this report is dubious and the species is therefore not included in this list.

Literature

Vuković (1963a).

Genus *Thymallus* Linck, 1790

86. *Thymallus thymallus* (Linnaeus, 1758)

Vernacular names

Lipljen. European grayling.

Previous names used for the species

Salmo thymallus Linnaeus, 1758; *Thymallus vulgaris* Nilsson, 1832.

Origin and geographic range in BiH

Native to the the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to different rivers of the Adriatic Basin, in BiH, like the upper Neretva River and Šuica River (Duvanjsko Polje karst field) (Vuković & Sofradžija 1987; Hamzić 2024).

Literature

Anonymous 1886; Glowacki 1896; Langhoffer 1904; Jedlička 1931, 1932; Zaplata & Taler 1932; Taler 1944, 1945a; Aganović 1952b, 1952c; Stepanek 1953; Šenk 1953, 1956a, 1969; Kosorić & Kardoš 1955; Aganović 1957, 1965a, 1965b; Bogdanović 1957; Janković 1960; Pavlović *et al.* 1960, 1962; Smlatić 1961; Vuković 1963a, 1977a, 1982a; Horvat 1964; Mijatović 1966; Sabioncello 1967; Čanković *et al.* 1968a; Rukavina & Ćorić 1968; Šenk & Aganović 1968; Dedić 1971; Vuković & Ivanović 1971; Aganović & kapetanović 1973; Kačanski *et al.* 1977; Vuković & Kosorić 1978; Kosorić 1981; Kosorić *et al.* 1983, 1989; Imamović 1986; Kosorić & Mikavica 1986; Mikavica 1988a; Mikavica *et al.* 1988, 1991; Babić & Mikavica 1994; Mikavica & Savić 1999; Radević 2000; Mitrašinović 2002; Korjenić 2004a, 2005a, 2005b; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Škrijelj *et al.* 2006; Kalamujić *et al.* 2007; Kottelat & Freyhof 2007; Bakrač-Bećiraj 2008; Bećiraj *et al.* 2008, 2016; Kalamujić 2008; Muhamedagić *et al.* 2008, 2019; Dekić *et al.* 2009b, 2024; Korjenić *et al.* 2009; Sofradžija 2009; Ajanović *et al.* 2010; Ivanc *et al.* 2011; Marić *et al.* 2011, 2012; Mišić 2011; Bećiraj & Šahinović 2012; Golub *et al.* 2012; Muhamedagić & Habibović 2013; Trožić-Borovac *et al.* 2013; Simonović *et al.* 2015; Čičić-Močić 2018; Kovačević *et al.* 2018; Nedić *et al.* 2018; Lolić *et al.* 2019; Bakrač *et al.* 2020b; Skenderović *et al.* 2020b; Neuburg *et al.* 2023; Hamzić 2024.

Order Gadiformes Goodrich, 1909

Family Lotidae Bonaparte, 1832

Genus *Lota* Oken, 1817

87. *Lota lota* (Linnaeus, 1758)

Vernacular names

Manić, menek. Burbot.

Previous names used for the species

Lota fluviatilis Perty, 1832; *Lota lota lota* (Linnaeus, 1758); *Lota vulgaris* Fitzinger, 1832.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Glowacki 1896; Brusina 1892; Medić 1901; Ćurčić 1910, 1933; Plančić 1923b; Vuković 1963a, 1977a; Čanković *et al.* 1968a, 1968c; Veljović 1982, 1985; Sofradžija *et al.* 1984; Bojčić 1987; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002, 2006; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Ivanc & Dekić 2012; Delić *et al.* 2014; Simonović *et al.* 2015; Glamuzina *et al.* 2017; Golub *et al.* 2018a, 2018b; Kovačević *et al.* 2018; Skenderović *et al.* 2020b; Dekić *et al.* 2024; Hamzić 2024.

Order Syngnathiformes
Family Syngnathidae Bonaparte, 1831
Genus *Syngnathus* Linnaeus, 1758

ME3 *Syngnathus abaster* Risso, 1827

Vernacular names

Šilo kratkokljuno. Black-striped pipefish.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Although the black-striped pipefish was previously included in the list of fishes of BiH (Vuković 1963a, 1977a; Bogut *et al.* 2006), Tutman *et al.* (2012b) provided the first reliable record of its presence with the exact location, including morphometric and meristic information of a specimen.

Literature

Vuković 1963a, 1977a; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Tutman *et al.* 2012b.

Order Gobiiformes Bleeker, 1859

A review by Tutman *et al.* (2020b) updated the information on the taxonomic status and distribution of freshwater gobies in BiH (families Oxudercidae and Gobiidae), and four species are excluded from the national checklist given the lack of data supporting their occurrence: *Knipowitschia panizzae*, *Pomatoschistus microps*, *Proterorhinus marmoratus* and *Zosterisessor ophiocephalus*. Therefore, the freshwater gobiid fauna of BiH includes seven species in six genera. The Adriatic Basin (Neretva River catchment) is inhabited by three endemic species: *Knipowitschia radovici*, *Orsinigobius croaticus* and *Ninnigobius canestrinii*, while the Danube River Basin is inhabited by one native species, *Proterorhinus semilunaris*, and four invasive species: *Babka gymnotrachelus*, *Neogobius fluviatilis*, *Neogobius melanostomus* and *Ponticola kessleri*. Distribution of the endemics *K. radovici* and *O. croaticus* is restricted to small areas in the lower Neretva River catchment on both sides of the Croatia–BiH border.

Family Odontobutidae Hoese & Gill, 1993
Genus *Perccottus* Dybowski, 1877

88. *Perccottus glenii* Dybowski, 1877

Vernacular names

Amurski spavač. Amur sleeper.

Origin and geographic range in BiH

Non-native. Danube River Basin.

Remarks

Reported from the Sava River near Slavonski Brod and Bosanski Brod (Ćaleta *et al.* 2010).

Literature

Ćaleta *et al.* 2010.

Family Oxudercidae Günther, 1861

Genus *Knipowitschia* Iljin, 1927

Knipowitschia panizzae (Verga, 1841)

Vernacular names

Glavočić vodenjak. Adriatic dwarf goby.

Previous names used for the species

Gobius pannizae Verga, 1841; *Padogobius panizzai* (Verga, 1841).

Origin and geographic range in BiH

Adriatic Basin.

Remarks

First recorded as *Gobius panizzae* by Vuković (1963a), but without information on its distribution in BiH. Later, Vuković (1977a) reported it as *Padogobius panizzai* in the lower Neretva River catchment, while Sofradžija (2009) listed it as *Knipowitschia panizzae* for the same area; it was not mentioned by Bogut *et al.* (2006). Since its taxonomic status in the Adriatic Sea is doubtful (Kovačić & Pallaoro 2003), and is not recorded in BiH (Šanda & Kovačić 2009), until confirmation *K. panizzae* should not be included in this checklist.

Literature

Vuković 1977a; Mikavica 1998; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Šanda & Kovačić 2009; Drešković *et al.* 2011.

89. *Knipowitschia radovici* Kovačić, 2005

Vernacular names

Norinski glavočić. Norin goby.

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

Stenoendemic species that inhabits only the lower Neretva River catchment in Croatia (tributary Norin River; Kovačić 2005) and BiH (Hutovo Blato wetland; Šanda & Kovačić 2009; Tutman *et al.* 2013). However, molecular results grouped this species with *Knipowitschia panizzae* and *K. mrakovcici* (Geiger

et al. 2014, Thacker *et al.* 2019); therefore, according to Tougard *et al.* (2021), the taxonomic status of this species requires additional clarification.

Literature

Glamuzina *et al.* 2001, 2002, 2008a; Tutman *et al.* 2002, 2009b, 2012a, 2020b; Bogut *et al.* 2006; Dulčić *et al.* 2008; Šanda *et al.* 2008a, 2008b, 2009; Šanda & Kovačić 2009; Darwall *et al.* 2014; Geiger *et al.* 2014; Thacker *et al.* 2019; Freyhof *et al.* 2020; Tougard *et al.* 2021.

Genus *Ninnigobius* Whitley, 1951

90. *Ninnigobius canestrinii* (Ninni, 1883)

Vernacular names

Glavočić crnotrus. Black-spot goby.

Previous names used for the species

Gobius canestrinii Ninni, 1883; *Pomatoschistus canestrinii* (Ninni, 1883).

Origin and geographic range in BiH

Endemic (ABE). Adriatic Basin.

Remarks

Endemic to the Adriatic Basin. It was first reported as *Gobius canestrinii* by Vuković (1963a), but without any distribution data. Later, Vuković (1977a), Bogut *et al.* (2006), Sofradžija (2009) and Drešković *et al.* (2011) listed it as *Pomatoschistus canestrinii* in the lower Neretva River. However, no official records were published (Kosorić 1978; Kosorić *et al.* 1983; Šanda & Kovačić 2009) until Tutman *et al.* (2013) presented the first reliable report of its occurrence in the Hutovo Blato wetland.

Literature

Vuković 1963a, 1977a; Mikavica 1998; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Tutman *et al.* 2012a, 2013, 2020b; Geiger *et al.* 2014.

Genus *Orsinigobius* Gandolfi, Marconato & Torricelli, 1986

91. *Orsinigobius croaticus* (Mrakovčić, Kerovec, Mišetić & Schneider, 1994)

Vernacular names

Vrgoračka gobica. Croatian goby.

Previous names used for the species

Knipowitschia punctatissima croatica Mrakovčić, Kerovec, Mišetić & Schneider, 1994.

Origin and geographic range in BiH

Endemic (NCE, BiH, CRO). Adriatic Basin.

Remarks

Endemic to the Neretva River catchment, originally described as a subspecies, *Knipowitschia punctatissima croatica*, from a restricted area in Croatia (Mrakovčić *et al.* 1996). It was later recognized

as a species, *K. croatica* (Kottelat 1997). Following molecular studies by Geiger *et al.* (2014) and Thacker *et al.* (2019), it was recently transferred to the genus *Orsinigobius*, as *O. croaticus*. It occurs only in the karst watercourses of the Neretva River catchment in BiH and Croatia. Its current distribution in BiH covers the lower Neretva River up to the town of Čapljina, its tributaries Bregava and Trebižat Rivers, and the Hutovo Blato wetland (Šanda & Kovačić 2009; Drešković *et al.* 2011; Glamuzina *et al.* 2013; Tutman *et al.* 2013, 2020b).

Literature

Glamuzina *et al.* 2001, 2002, 2008a; Tutman *et al.* 2002, 2009b, 2012a, 2020b; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Dulčić *et al.* 2008; Jelić *et al.* 2008; Šanda *et al.* 2008a, 2008b, 2009; Šanda & Kovačić 2009; Zupančič 2008; Drešković *et al.* 2011; Darwall *et al.* 2014; Geiger *et al.* 2014; Oikonomou *et al.* 2014; Milanović *et al.* 2015; Horvatić *et al.* 2017; Zanella *et al.* 2017; Freyhof *et al.* 2020.

Genus *Pomatoschistus* Gill, 1863

Pomatoschistus microps (Krøyer, 1838)

Previous names used for the species

Gobius microps Krøyer, 1838; *Pomatoschistus microps* (Krøyer, 1838).

Remarks

First reported as *Gobius microps* (Vuković 1963a), while Vuković (1977a) and Bogut *et al.* (2006) referred to it as *Pomatoschistus microps* for the lower Neretva River catchment. It was not listed by Sofradžija (2009). This species is not present in the Adriatic Sea (Dulčić & Kovačić 2020), therefore all reports from the Neretva River catchment result from misidentification of specimens. Accordingly, it is not included in this checklist.

Literature

Vuković 1963a, 1977a; Bogut *et al.* (2006).

Family Gobiidae Cuvier, 1816

Genus *Babka* Iljin, 1927

92. ***Babka gymnotrachelus*** (Kessler, 1857)

Vernacular names

Glavočić trkač. Racer goby.

Origin and geographic range in BiH

Non-native. Danube River Basin.

Remarks

It was first reported in BiH as *Neogobius gymnotrachelus*, between the cities of Orašje and Tolisa in the middle part of the Sava River (Nedić *et al.* 2018).

Literature

Nedić *et al.* 2018; Skenderović *et al.* 2020b.

Genus *Neogobius* Iljin, 1927

93. *Neogobius fluviatilis* (Pallas, 1814)

Vernacular names

Riječni glavoč. Pontian monkey goby.

Previous names used for the species

Gobius fluviatilis Pallas, 1811.

Origin and geographic range in BiH

Non-native. Danube River Basin.

Remarks

Vuković (1977a) was the first to record the presence of *Neogobius fluviatilis* (as *Gobius fluviatilis*) in the lowlands of the Danube River Basin in northern BiH, with remarks that this species was rare, but without providing any precise distribution data. Similar information was given by Bogut *et al.* (2006) and Sofradžija (2009). Nedić *et al.* (2014a, 2014c) and Delić *et al.* (2014) recorded it in the Sava and Una rivers, respectively.

Literature

Vuković 1977a; Mrakovčić *et al.* 2002; Škrijelj *et al.* 2005a; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Čaleta *et al.* 2010; Piria 2011b; Delić *et al.* 2014; Nedić *et al.* 2014a, 2014c, 2014d; Simonović *et al.* 2015; Golub *et al.* 2017c, 2018a, 2018b; Tutman *et al.* 2020b.

94. *Neogobius melanostomus* (Pallas, 1814)

Vernacular names

Round goby. Glavočić okrugljak.

Origin and geographic range in BiH

Non-native. Danube River Basin.

Remarks

Reported from the Sava River (Piria *et al.* 2011a, 2016a; Jakovlić *et al.* 2015), and Una River near Kozarska Dubica (Čolić *et al.* 2018).

Literature

Piria *et al.* 2011a, 2016a; Jakovlić *et al.* 2015; Simonović *et al.* 2015; Čolić *et al.* 2018; Tutman *et al.* 2020b.

Genus *Padogobius* Berg, 1932

Padogobius bonelli (Bonaparte, 1846)

Remarks

Listed in Hamzić (2024) with the remark that its presence has yet to be confirmed. Therefore, it is not included in this checklist.

Literature

Hamzić (2024).

Genus *Ponticola* Iljin, 1927

95. *Ponticola kessleri* (Günther, 1861)

Vernacular names

Bičkaš. Pontian bighead goby.

Previous names used for the species

Gobius kessleri (Günther, 1861); *Neogobius kessleri* (Günther, 1861).

Origin and geographic range in BiH

Non-native. Danube River Basin.

Remarks

This species was first mentioned in BiH as *Gobius kessleri* by Vuković (1977a), in the lowlands of the Danube River Basin, without precise distribution data, and it was considered rare. This information was repeated also by Bogut *et al.* (2006), as *Neogobius kessleri*, while Sofradžija (2009) considered its occurrence in the Sava River and its lower tributaries. Recently, Jakovlić *et al.* (2015) and Piria *et al.* (2016a) also reported its presence in the Sava River in Croatia.

Literature

Vuković 1977a; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Čaleta *et al.* 2010; Tutman *et al.* 2020b; Skenderović *et al.* 2020b.

Genus *Proterorhinus* Smitt, 1900

Proterorhinus marmoratus (Pallas, 1814)

Previous names used for the species

Gobius marmoratus Pallas, 1814.

Remarks

It was first mentioned in BiH by Vuković (1977a) for the lowland waters of the Danube River Basin. However, its precise location was not provided, though it was considered rare. This information was later repeated by Bogut *et al.* (2006), while Sofradžija (2009) listed it for the waters of Sava River. However, molecular studies of the genus in the Danube River Basin (Neilson & Stepien 2009a, 2009b) confirmed that *Proterorhinus marmoratus* is present only in marine and brackish waters at the mouth of

the Danube Basin, while inland, freshwater representatives were confirmed as *P. semilunaris*. Therefore, *P. marmoratus* is not included in this checklist.

Literature

See *Proterorhinus semilunaris*.

96. *Proterorhinus semilunaris* (Heckel, 1837)

Vernacular names

Mramorasti glavoč. Western tubenose goby.

Previous names used for the species

Gobius marmoratus Pallas, 1814; *Proterorhinus marmoratus* (Pallas, 1814).

Origin and geographic range in BiH

Native. Danube River Basin.

Remarks

Literature records mention *Proterorhinus marmoratus*, even though this was the result of misidentification of specimens (Hamzić, 2024). In recent years, *P. semilunaris* was recorded in the Sava River at the confluence with the Ukrina River (Čaleta & Marčić, unpublished data).

Literature

Vuković 1963a, 1977a; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Sofradžija 2009; Tutman *et al.* 2020b; Dekić *et al.* 2024.

Genus *Zoosterisessor* Whitley, 1935

Zoosterisessor ophiocephalus (Pallas, 1811)

Vernacular names

Glavoč travaš. Grass goby.

Previous names used for the species

Gobius ophiocephalus Pallas, 1811.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

This species was originally recorded in BiH as *Gobius ophiocephalus* by Vuković (1963a), but no distribution data was provided. Sofradžija (2009) did not include the species in his study, but Vuković (1977a) and Bogut *et al.* (2006) reported it in the lower Neretva, again without providing distributional data. It seems it does not occur in BiH, but is present in the lower reaches of the Neretva River in Croatia, in brackish waters and lagoons (Dulčić *et al.* 2007). Although this species is regularly found in transitional waters (Kottelat & Freyhof 2007), there is no reliable information of its occurrence in BiH, and therefore it is not included in this checklist.

Literature

Vuković 1977a; Bogut *et al.* 2006; Tutman *et al.* 2020b.

Order Carangiformes Jordan, 1923
Family Pleuronectidae Rafinesque, 1815
Genus *Platichthys* Girard, 1854

ME4 *Platichthys flesus* (Linnaeus, 1758)

Vernacular names

Iverak, kalkan. Flounder.

Previous names used for the species

Pleuronectes flesus Linnaeus, 1758; *Pleuronectes flesus luscus* Pallas, 1814.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Distributed in the lower reaches of the Neretva River and the Hutovo blato wetland, where it enters from the coastal areas of the Adriatic Sea.

Literature

Vuković 1963a, 1977a; Kosorić 1978; Kosorić *et al.* 1983, 1989; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Glamuzina *et al.* 2008a; Sofradžija 2009; Tutman *et al.* 2012a.

Order Atheriniformes Rosen, 1966
Family Atherinidae Risso, 1827
Genus *Atherina* Linnaeus, 1758

ME5 *Atherina boyeri* Risso, 1810

Vernacular names

Oliga, gavun. Sand smelt.

Previous names used for the species

Hepsetia boyeri Jordan & Hubbs, 1919.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Occurs in the lower reaches of the Neretva River and the Hutovo blato wetland, where it enters from coastal areas of the Adriatic Sea.

Literature

Vuković 1963a, 1977a; Mikavica 1998; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Tutman *et al.* 2012a.

Atherina hepsetus Linnaeus, 1758

Vernacular names

Gavun. Mediterranean sand smelt.

Previous names used for the species

Atherina athaerina Nardo, 1827; *Atherina mochon* Cuvier, 1829; *Hepsetia mochon* Schultz, 1948.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Listed in Vuković (1963a) as *Atherina mocho* and in Vuković (1977a) as *A. mochon*. In Bogut *et al.* (2006) and Hamzić (2024), it is mentioned as *A. hepsetus*. It was not mentioned by Sofradžija (2009). Glamuzina & Dobrosravić (2020) did not report the species either from the lower Neretva River stretches in Croatia. Environmental conditions of the lower Neretva River are indeed not suitable for the species; therefore, its occurrence in the BiH is unlikely and the species is not included in this checklist.

Literature

Vuković 1963a, 1977a; Bogut *et al.* 2006; Hamzić 2024.

Order Cyprinodontiformes Berg, 1940

Family Cyprinodontidae Gill, 1865

Genus *Aphanius* Nardo, 1827

Aphanius fasciatus (Valenciennes, 1821)

Vernacular names

Obrvan. Mediterranean toothcarp.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Recorded by Sofradžija (2009) and Hamzić (2024) in the lower Neretva River, without distributional data provided. However, since environmental conditions of the lower Neretva River are not suitable for the species, its occurrence in the BiH is unlikely and the species is not included in this checklist.

Literature

Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Hamzić 2024.

Family Poeciliidae Bonaparte, 1831

Genus *Gambusia* Poey, 1854

Gambusia affinis Girard, 1859

Previous names used for the species

Gambusia affinis (Baird & Girard, 1853).

Remarks

The record provided by Sofradžija (2009) is most likely based on misidentification of specimens. Landeka *et al.* (2015) confirmed the presence of only *Gambusia holbrooki* across the entire Mediterranean biogeographical region in Croatia and BiH. Therefore, it is not included in this checklist.

Literature

See in *Gambusia holbrooki*.

97. *Gambusia holbrooki* Girard, 1859

Vernacular names

Gambuzija. Eastern mosquitofish.

Previous names used for the species

Gambusia affinis Girard, 1859; *Gambusia affinis holbrooki* (Girard, 1859).

Origin and geographic range in BiH

Non-native. Adriatic Basin.

Remarks

Older literature mentioned *Gambusia affinis* Girard, 1859 (Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009) and *G. affinis holbrooki* (Girard, 1859) (Landeka *et al.* 2015) in BiH, but those records are likely misidentification of specimens, since according to Landeka *et al.* (2015), only *G. holbrooki* occurs in the Mediterranean biogeographic region in Croatia and BiH.

Literature

Vuković 1963a, 1977a; Kosorić 1978; Kosorić *et al.* 1983, 1989; Bogut *et al.* 2006; Glamuzina *et al.* 2008a, 2017; Šanda *et al.* 2008a, 2008b; Sofradžija 2009; Tutman *et al.* 2012a; Landeka *et al.* 2015; Piria *et al.* 2018.

Order Mugiliformes Günther, 1880

Family Mugilidae Jarocki, 1822

Genus *Chelon* Artedi, 1793

ME6 *Chelon auratus* (Risso, 1810)

Vernacular names

Cipol zlatac. Golden mullet.

Previous names used for the species

Mugil auratus Risso, 1810; *Liza aurata* (Risso, 1810).

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Reported in the literature (Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009) as occurring in the lower Neretva River, where it enters from coastal areas of the Adriatic Sea. However, data on its distribution are based on very general information and no actual specimens with detailed description have ever been reported (Kosorić 1978; Kosorić *et al.* 1983; Mikavica 1998). This has led to doubts about its occurrence in BiH. Reliable data about its occurrence in the Neretva River in Croatia were presented by Glamuzina & Dobroslavić (2020). Therefore, more information confirming its occurrence is necessary, with previous reports considered unreliable.

Literature

Aganović 1952a; Vuković 1963a, 1977a; Kosorić 1978; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009.

ME7 *Chelon labrosus* (Risso, 1827)

Vernacular names

Cipol putnik. Thicklip mullet.

Previous names used for the species

Mugil labrosus Risso, 1827; *Mugil chelo* Cuvier, 1829; *Liza chelo* Popov, 1929; *Liza labrosus* Buen, 1935.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Reported in the literature as occurring in the lower Neretva River, where it enters from coastal areas of the Adriatic Sea (Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009). However, data on its occurrence are based on very general information and no actual specimens with detailed descriptions have ever been reported (Kosorić 1978; Kosorić *et al.* 1983; Mikavica 1998). More reliable data about its occurrence for the Neretva River in Croatia were presented by Glamuzina & Dobroslavić (2020). Therefore, more information confirming its occurrence is necessary, with previous reports being unreliable.

Literature

Vuković 1963a, 1977a; Kosorić 1978; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009.

ME8 *Chelon ramada* (Risso, 1827)

Vernacular names

Cipol balavac. Thinlip mullet.

Previous names used for the species

Mugil ramada Risso, 1826; *Liza ramada* Risso, 1827; *Mugil capito* Cuvier, 1829.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Distributed in the lower reaches of the Neretva River and the Hutovo blato wetland, where it enters from coastal areas of the Adriatic Sea.

Literature

Vuković 1963a, 1977a; Kosorić 1978; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Tutman *et al.* 2012a.

ME9 *Chelon saliens* (Risso, 1810)

Vernacular names

Cipol dugaš. Sharpnose mullet.

Previous names used for the species

Mugil saliens Risso, 1810 ; *Liza saliens* (Risso, 1810).

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Reported in the literature as occurring in the lower Neretva River, where it enters from coastal areas of the Adriatic Sea (Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009). However, data on its distribution are based on very general information and no actual specimens with detailed description have ever been reported (Kosorić 1978; Kosorić *et al.* 1983; Mikavica 1998). This has led to doubts about its occurrence in BiH. Reliable data about its occurrence in the Neretva River in Croatia were presented by Glamuzina & Dobroslavić (2020). Therefore, more information confirming its occurrence is necessary, with previous reports unreliable.

Literature

Vuković 1963a, 1977a; Kosorić 1978; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Tutman *et al.* 2012a.

Genus *Mugil* Linnaeus, 1758

ME10 *Mugil cephalus* Linnaeus, 1758

Vernacular names

Cipol bataš. Striped mullet.

Previous names used for the species

Mugil vulpinus Nardo, 1847.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Distributed in the lower reaches of the Neretva River and the Hutovo Blato wetland, where it enters from coastal areas of the Adriatic Sea.

Literature

Aganović 1952a; Vuković 1963a, 1977a; Kosorić 1978; Kosorić *et al.* 1983, 1989; Bogut *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Tutman *et al.* 2012a; Skenderović *et al.* 2020b.

Genus *Oedalechilus* Fowler, 1903

Oedalechilus labeo (Cuvier, 1829)

Vernacular names

Cipol plutaš. Boxlip mullet.

Previous names used for the species

Mugil labeo Cuvier, 1829.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

According to Vuković (1963a, 1977a), it is distributed in the lower reaches of the Neretva River, where it enters from the coastal areas of the Adriatic Sea. However, no actual specimens with detailed descriptions have ever been reported (Kosorić 1978; Kosorić *et al.* 1983; Mikavica 1998). The report of the occurrence of the species in BiH by Vuković (1963a, 1977a) is therefore unreliable. Furthermore, environmental conditions of the lower Neretva River are not suitable for the species; therefore, its occurrence in the BiH is unlikely and the species is not included in this checklist.

Literature

Vuković 1963a, 1977a; Bogut *et al.* 2006.

Order Blenniiformes Bleeker, 1860

Family Blenniidae Rafinesque, 1810

Genus *Salaria* Forsskål, 1775

98. *Salariopsis fluviatilis* (Asso, 1801)

Vernacular names

Riječna babica. Freshwater blenny.

Previous names used for the species

Blennius fluviatilis Asso, 1801; *Blennius vulgaris* Pollini, 1816; *Lipophrys fluviatilis* (Asso, 1801).

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Although previously included in lists of freshwater fishes of BiH (Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009), its occurrence in the lower Neretva River near the border with Croatia was confirmed for the first time by Tutman & Glamuzina (2021).

Literature

Vuković 1963a, 1977a; Mikavica 1998; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Tutman & Glamuzina 2021.

Order Perciformes Bleeker, 1863
Family Percidae Rafinesque, 1815
Genus *Gymnocephalus* Bloch, 1793

99. *Gymnocephalus baloni* Holčik & Hensel, 1974

Vernacular names

Balonov balavac. Danube ruffe.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Kottelat & Freyhof 2007; Pavlović *et al.* 2009; Drešković *et al.* 2011; Lukić *et al.* 2016; Simonović *et al.* 2015; Dekić *et al.* 2024.

100. *Gymnocephalus cernua* (Linnaeus, 1758)

Vernacular names

Balavac. Ruffe.

Previous names used for the species

Acerina cernua (Linnaeus, 1758).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Distributed in the Danube River Basin of BiH (Vuković 1977a), reported as translocated in the Hutovo Blato wetland in the lower portion of the Neretva River (Dulčić *et al.* 2005).

Literature

Anonymous 1886; Glowacki 1896; Ćurčić 1910; Vuković 1963a, 1977a; Čanković *et al.* 1968a; Kosorić *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Veljović 1982, 1985; Mikavica 1987b; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002; Dulčić *et al.* 2005; Bogut *et al.* 2006; Skenderović *et al.* 2006, 2011a, 2020a, 2020b; Adrović 2007, 2012; Kottelat & Freyhof 2007; Vuković *et al.* 2008; Sofradžija 2009; Tutman *et al.* 2012a, 2019; Čičić-Močić 2014; Milanović *et al.*

2015; Simonović *et al.* 2015; Skenderović 2015; Glamuzina *et al.* 2017; Mrdak & Milošević 2017; Adrović 2018; Memić 2018; Lolić *et al.* 2019; Dekić *et al.* 2020a.

101. *Gymnocephalus schraetser* (Linnaeus, 1758)

Vernacular names

Prugasti balavac. Stripped ruffe.

Previous names used for the species

Acerina schraetser (Linnaeus, 1758); *Perca schraetser* Linnaeus, 1758.

Origin and geographic range in BiH

Endemic (DBE). Danube River Basin.

Remarks

Endemic to the Danube River Basin.

Literature

Glowacki 1896; Ćurčić 1910; Vuković 1963a, 1977a; Mikavica 1987b; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Pavlović *et al.* 2009; Sofradžija 2009; Popović 2010; Drešković *et al.* 2011; Simonović *et al.* 2015; Lukić *et al.* 2016; Dekić *et al.* 2024.

Genus *Perca* Linnaeus, 1758

102. *Perca fluviatilis* Linnaeus, 1758

Vernacular names

Grgeč. Perch.

Previous names used for the species

Perca fluviatilis fluviatilis Linnaeus, 1758; *Perca vulgaris* Fitzinger, 1832.

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to the Hutovo Blato wetland and reservoirs in the lower Neretva River (Glamuzina *et al.* 2017).

Literature

Anonymous 1886; Brusina 1892; Glowacki 1896; Ćurčić 1910; Jedlička 1935; Vuković 1963a, 1977a; Kiškarolj 1965; Čanković *et al.* 1968a, 1968b, 1968c; Žitnan *et al.* 1969; Kosorić *et al.* 1980; Habeković *et al.* 1981; Obratil 1982; Veljović 1982, 1985; Delić 1984; Mikavica *et al.* 1991; Leiner 1998; Mikavica & Savić 1999; Radević 2000; Mrakovčić *et al.* 2002; Bogut *et al.* 2006; Skenderović *et al.* 2006, 2011a, 2020a, 2020b; Škrijelj *et al.* 2006; Adrović 2007, 2012, 2018; Kottelat & Freyhof 2007; Vuković *et al.* 2008; Sofradžija 2009; Ivanc *et al.* 2011; Tutman *et al.* 2012a; Golub *et al.* 2013, 2016a, 2017c, 2018a, 2018b; Čičić-Močić 2014; Nedić *et al.* 2014a, 2014c, 2014d, 2018; Ridanović *et al.* 2015;

Simonović *et al.* 2015; Skenderović 2015; Glamuzina *et al.* 2017; Kovačević *et al.* 2018; Memić 2018; Lolić *et al.* 2019; Cvijić *et al.* 2020; Dekić *et al.* 2020a, 2024.

Genus *Sander* Oken, 1817

103. *Sander lucioperca* (Linnaeus, 1758)

Vernacular names

Smuđ. Pikeperch.

Previous names used for the species

Lucioperca lucioperca (Linnaeus, 1758); *Stizostedion lucioperca* (Linnaeus, 1758).

Origin and geographic range in BiH

Native to the Danube River Basin, translocated to the Adriatic Basin.

Remarks

Translocated to the reservoirs in the lower Neretva River (Pavličević *et al.* 2016).

Literature

Anonymous 1886, 1947b; Jurinac 1887; Brusina 1892; Glowacki 1896; Medić 1901; Langhoffer 1904; Ćurčić 1910; Plančić 1923b; Taler 1932; Bek 1935; Vuković 1963a, 1977a; Čanković 1964; Kiškarolj 1965; Čanković *et al.* 1968a, 1968c; Žitnan *et al.* 1969; Munjko 1972; Kosorić *et al.* 1980; Habeković *et al.* 1981; Veljović 1982, 1985; Delić 1984; Mikavica *et al.* 1991; Pažur 1993; Leiner 1998; Mikavica & Savić 1999; Radević 2000; Škrijelj & Mašović 2001; Korjenić *et al.* 2003; Škrijelj *et al.* 2003, 2011; Bogut *et al.* 2006; Skenderović *et al.* 2006, 2011a, 2020a, 2020b, 2021; Adrović 2007, 2012; Adrović & Skenderović 2007; Kottelat & Freyhof 2007; Trožić-Borovac & Škrijelj 2007; Muhamedagić *et al.* 2008; Suić *et al.* 2008; Šanda *et al.* 2008b; Vuković *et al.* 2008; Sofradžija 2009; Jamaković 2011; Tutman *et al.* 2012a, 2019; Čičić-Močić 2014; Nedić *et al.* 2014a, 2014c, 2014d, 2018; Riđanović *et al.* 2015; Simonović *et al.* 2015; Skenderović 2015; Mitrašinović-Brulić *et al.* 2016; Pavličević *et al.* 2016; Šljuka 2011, 2016; Glamuzina *et al.* 2017; Kovačević *et al.* 2018; Memić 2018; Mušović *et al.* 2018, 2020, 2025; Žujo-Zekić *et al.* 2018; Dekić *et al.* 2020a, 2024; Žujo-Zekić 2021.

104. *Sander volgensis* (Gmelin, 1789)

Vernacular names

Smuđ kamenjak. Volga pikeperch.

Previous names used for the species

Perca volgensis Gmelin, 1788; *Stizostedion volgensis* (Gmelin, 1788).

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Glowacki 1896; Medić 1896, 1901; Ćurčić 1910; Anonymous 1947b; Vuković 1963a, 1977a; Radević 2000; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Dekić *et al.* 2024.

Genus *Zingel* Cloquet, 1817

105. *Zingel streber* (Siebold, 1863)

Vernacular names

Mali vretenac. Streber.

Previous names used for the species

Aspro streber Siebold, 1863; *Aspro streber streber* Siebold, 1863; *Aspro vulgaris* Cuvier, 1828.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Anonymous 1886; Glowacki 1896; Medić 1901; Ćurčić 1910; Zaplata & Taler 1932; Vuković 1963a, 1977a; Veljović 1982; Mikavica 1987b; Mikavica *et al.* 1991; Mikavica & Savić 1999; Radević 2000; Korjenić 2004b; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Pavlović *et al.* 2009; Sofradžija 2009; Popović 2010; Drešković *et al.* 2011; Milanović *et al.* 2015; Simonović *et al.* 2015; Kovačević *et al.* 2018; Lukić *et al.* 2016; Dekić *et al.* 2024.

106. *Zingel zingel* (Linnaeus, 1766)

Vernacular names

Veliki vretenac. Zingel.

Previous names used for the species

Perca zingel Linnaeus, 1766; *Aspro zingel* Linnaeus, 1766.

Origin and geographic range in BiH

Native. Danube River Basin.

Literature

Glowacki 1896; Ćurčić 1910; Vuković 1963a, 1977a; Kosorić *et al.* 1980; Veljović 1982; Mikavica *et al.* 1991; Mikavica & Savić 1999; Bogut *et al.* 2006; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Sofradžija 2009; Drešković *et al.* 2011; Milanović *et al.* 2015; Simonović *et al.* 2015; Kovačević *et al.* 2018; Nedić *et al.* 2018; Skenderović *et al.* 2020b; Dekić *et al.* 2024.

Family Gasterosteidae Bonaparte, 1831

Genus *Gasterosteus* Linnaeus, 1758

107. *Gasterosteus aculeatus* Linnaeus, 1758

Vernacular names

Bodonja. Threespine stickleback.

Previous names used for the species

Gasterosteus aculeatus gymnurus Cuvier, 1829.

Origin and geographic range in BiH

Native. Adriatic Basin.

Literature

Karaman 1928; Taler 1953a; Vuković 1963a, 1977a; Vuković & Prolić 1966; Vuković & Kosorić 1967; Čemalović 1975; Kosorić 1978; Kosorić *et al.* 1983, 1989; Hećo 2000; Cano *et al.* 2006, 2008; Mäkinen *et al.* 2006, 2008a, 2008b; Mrakovčić *et al.* 2006; Kottelat & Freyhof 2007; Glamuzina *et al.* 2008a; Mäkinen & Merilä 2008; Šanda *et al.* 2008a, 2008b; DeFaveri *et al.* 2012; Hamzić *et al.* 2012b; Tutman *et al.* 2012a; Zanella *et al.* 2015; Coll-Costa *et al.* 2024; Hamzić 2024.

Genus *Pungitius* Coste, 1848

Pungitius platygaster (Kessler, 1859)

Vernacular names

Gregorac. Ukrainian stickleback.

Remarks

Reported from the Danube up to Belgrade (Kottelat & Freyhof 2007). Mentioned in Sofradžija (2009) for the Danube River Basin, but without a more precise distribution area. There is no confirmation of its presence in recent literature, and additional research is needed.

Literature

Sofradžija 2009.

Family Cottidae Bonaparte, 1831

Genus *Cottus* Linnaeus, 1758

108. *Cottus gobio* Linnaeus, 1758

Vernacular names

Peš. Sculpin.

Previous names used for the species

Cottus gobio gobio Linnaeus, 1758; *Cottus ferrugineus* Bonaparte, 1846; *Cottus gobio ferugineus* Heckel & Kner, 1858.

Origin and geographic range in BiH

Native. Danube River and Adriatic Basin.

Remarks

Distributed in both the Danube and Adriatic basins. Although genetic analyses conducted by Šlechtová *et al.* (2004) support the existence of a single species in the region, several divergent genetic clades have subsequently been identified (Bravničar 2012; Bravničar *et al.* 2015, 2021; Neuburg *et al.* 2023). Specimens sampled in the upper Neretva River seem to be more related to populations from Danube tributaries in the northern Balkan Peninsula, despite the fact that the Neretva River drains towards the Adriatic Basin (Bravničar *et al.* 2015). Furthermore, Heckel & Kner (1858) described *Cottus ferrugineus* as a species endemic to the Adriatic Basin. Mentioned in Vuković (1977a) and Bogut *et al.* (2006)

as *C. cottus ferrugineus* for the lower Neretva River. However, according to Šlechtová *et al.* (2004), *C. ferrugineus* is not a valid species and therefore is not included in this checklist. Until this situation is resolved, only *C. gobio* is included in the checklist of fishes of BiH.

Literature

Brusina 1892; Glowacki 1896; Langhoffer 1904; Zaplata & Taler 1932; Aganović & Milošević 1959; Smlatić 1961; Vuković 1963a, 1977a; Kiškarolj 1965; Okuka 1966; Čanković 1967; Čanković *et al.* 1968a; Vuković & Ivanović 1971; Aganović & kapetanović 1973; Berberović & Sofradžija 1974b; Kosorić 1974, 1977a, 1981; Sofradžija & Berberović 1975; Kačanski *et al.* 1977; Kosorić *et al.* 1980, 1983, 1989; Sofradžija *et al.* 1984; Kosorić & Mikavica 1986; Mikavica 1988a; Mikavica *et al.* 1991; Babić & Mikavica 1994; Mikavica & Savić 1999; Radević 2000; Korjenić 2004b, 2005a; Šlechtová *et al.* 2004; Bogut *et al.* 2006; Škrijelj *et al.* 2006; Kottelat & Freyhof 2007; Muhamedagić *et al.* 2008, 2019; Korjenić *et al.* 2009; Sofradžija 2009; Drešković *et al.* 2011; Bećiraj & Šahinović 2012; Golub *et al.* 2012, 2017c; NHMW 2014; Simonović *et al.* 2015; Bećiraj *et al.* 2016; Čičić-Močić 2016, 2018, 2019; Lukić *et al.* 2016; Adrović 2018; Keško 2018; Kovačević *et al.* 2018; Memić 2018; Lolić *et al.* 2019; Dekić *et al.* 2020a, 2020b; Skenderović *et al.* 2020a, 2020b; Neuburg *et al.* 2023.

Cottus poecilopus Heckel, 1836

Vernacular names

Šareni peš. Siberian bullhead.

Remarks

Mentioned in Vuković (1977a) and Bogut *et al.* (2006) but without any data on its distribution. According to Kottelat & Freyhof (2007), this species is distributed in northern Europe; therefore, previous reports are inaccurate and the species is not included in this list.

Order Centrarchiformes Bleeker, 1859

Family Centrarchidae Bleeker, 1859

Genus *Lepomis* (Rafinesque, 1819)

109. *Lepomis gibbosus* (Linnaeus, 1758)

Vernacular names

Sunčanica. Pumpkinseed.

Previous names used for the species

Eupomotis gibbosus (Linnaeus, 1758).

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Non-native species from North America. According to EU regulation it is considered an invasive alien species of EU concern.

Literature

Žitnan *et al.* 1969; Vuković 1977a; Kosorić 1978; Obratil 1982; Veljović 1982, 1985; Leiner 1998; Radević 2000; Mrakovčić *et al.* 2002; Trožić-Borovac *et al.* 2003; Škrijelj *et al.* 2005a, 2005b; Bogut *et al.* 2006; Adrović 2007, 2012, 2018; Adrović & Skenderović 2007; Glamuzina *et al.* 2008a, 2017; Has-Schön *et al.* 2008; Muhamedagić *et al.* 2008; Šanda *et al.* 2008a, 2008b; Vuković *et al.* 2008; Sofradžija 2009; Tutman *et al.* 2012a, 2019; Golub *et al.* 2013, 2018a, 2018b; Simonović *et al.* 2015; Skenderović 2015; Čulibrk *et al.* 2016; Pavličević *et al.* 2016; Mrdak & Milošević 2017; Kovačević *et al.* 2018; Memić 2018; Piria *et al.* 2018; Vesnić *et al.* 2018; Žujo-Zekić *et al.* 2018; Dekić *et al.* 2020a, 2024; Mušović *et al.* 2020; Skenderović *et al.* 2020b.

Genus *Micropterus* Lacepède, 1802

110. *Micropterus salmoides* (Lacepède, 1802)

Vernacular names

Pastrmski grgeč. Largemouth bass.

Origin and geographic range in BiH

Non-native. Danube River and Adriatic Basin.

Remarks

Non-native species from North America. The first confirmed record of the species was based on a single specimen captured in the Sava River near the small town of Kaoci (northern BiH) in December 2014 (Tutman *et al.* 2017c). The origin of this introduction is unknown, but it was most likely due to dispersal from neighbouring freshwater habitats from other parts of the Sava River catchment in Croatia and the Danube River in Serbia.

Literature

Radević 2001; Bogut *et al.* 2006; Tutman *et al.* 2017c; Piria *et al.* 2018; Lelo & Hamzić 2020; Hamzić 2024.

Order Acanthuriformes Jordan, 1923
Family Moronidae Jordan & Evermann, 1896
Genus *Dicentrarchus* (Gill, 1860)

ME11 *Dicentrarchus labrax* (Linnaeus, 1758)

Vernacular names

Lubin. Sea bass.

Previous names used for the species

Morone labrax (Linnaeus, 1758).

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Occurs in the lower reaches of the Neretva River and the Hutovo Blato wetland, where it enters from the coastal areas of the Adriatic Sea.

Literature

Aganović 1952a; Vuković 1963a, 1977a; Bogut *et al.* 2006; Sofradžija 2009; Tutman *et al.* 2012a.

Family Sparidae Rafinesque, 1810

Genus *Sparus* Linnaeus, 1758

ME12 *Sparus aurata* Linnaeus, 1758

Vernacular names

Komarča. Gilthead bream.

Previous names used for the species

Chrysophrys aurata Valenciennes, 1830.

Origin and geographic range in BiH

Native. Adriatic Basin.

Remarks

Present in the lower reaches of the Neretva River and the Hutovo Blato wetland, where it enters from coastal areas of the Adriatic Sea.

Literature

Vuković 1963a, 1977a; Mikavica 1998; Bogut *et al.* 2006; Sofradžija 2009; Tutman *et al.* 2012a.

Discussion

General overview of the taxonomic status of freshwater fishes of BiH

The ichthyofauna of BiH can be clearly divided into a northern Danubian and a southern Adriatic Sea parts (Supp. file 1; Table 1). Geographically, these parts are divided by the central massif of the Dinaric Arc mountains range. The northern part shares many species with Central Europe, while the southern part shares several species with the Mediterranean Basin (Kottelat & Freyhof 2007). The Adriatic Sea Basin features a high rate of endemism and highly specialised species in the relatively short karst watercourses flowing southwards into the Adriatic Sea. *Anguilla anguilla*, *Phoxinus phoxinus* and *Cottus gobio* are the only native species present in both the Danube and Adriatic Sea drainages. Consequently, the freshwater fishes of BiH, including lampreys, are of particular importance as part of the national, regional and European natural heritage, especially due to their diversity and high degree of endemism. This ichthyofaunal diversity is mainly the result of the complex geological and climatic history of the Balkan Peninsula, which has allowed multiple colonisations events from outside the area and long periods of hydrographic isolation of populations, leading to speciation (Hewitt 1999; Šlechtová *et al.* 2004). In addition, BiH stands at the crossroads of the main biogeographical regions of southeast Europe and the Mediterranean Danube River Basin and Adriatic Basin.

According to historical and recent scientific literature, taking into account all taxonomic changes, 123 species of fish have been recorded and confirmed in the freshwaters of BiH. However, when we consider only strictly freshwater species, excluding euryhaline ones, the number is reduced to 110 species. In relation to previous checklists, significant changes are proposed here in terms of composition and number of reported species. The first monograph by Vuković (1963a) reported 103 species, Vuković (1977a) reported 106 species, Bogut *et al.* (2006) reported 116 species, Sofradžija (2009) reported 118 species, whereas the most recent report on the freshwater species of the BiH

(Hamzić 2024) reported 137 species. Such differences can be the consequence of several factors, e.g., introduction of non-native species, taxonomic re-evaluations of existing taxa based on new information, and description of new species. Indeed, with the development of modern biomolecular research, many taxonomic aspects of the BiH freshwater fishes, including lampreys, have changed since the monographs of Sofradžija (2009) and Hamzić (2024), both still in use by ichthyologists.

Among the species reported as occurring in BiH according to our results, 87 are native, of which 38 are endemic, while 23 are non-native. Furthermore, there are 12 euryhaline species and one purely marine species. Amongst the endemic species, 15 were restricted only to the Neretva River catchment in BiH and Croatia, 10 were restricted to the Danube River Basin area, 10 to the broader Adriatic Basin, and three to the Cetina River catchment. A total of 18 species are new records for BiH, and two species were found to have expanded ranges. Thirty-four families and 75 genera are presented in total. The richest in species families are the Leuciscidae with 37 species, and the Salmonidae with 13 species, whereas there are 17 families with only one species represented in BiH (Supp. file 1).

The total number of species known from the Adriatic Basin (83) is slightly higher than that from the Danube River Basin (77) (Fig. 3). This is mostly due to the number of translocated Danube natives (22) and the introduction of non-native species (17) (Glamuzina *et al.* 2017; Vukić *et al.* 2019). Considering only the numbers of native species, the Danube River Basin has more species (57 vs 34 in the Adriatic Basin). However, from the biogeographic point of view, the fish diversity is indeed more unique in the Adriatic Basin where about 80% of the native species are endemic. The Neretva River has the highest species diversity and the Cetina River the lowest, and both are located in the Adriatic Basin (Supp. file 1; Table 1). Although the distinctiveness of fish populations in BiH rivers flowing into the Adriatic Sea has long been recognized (Vuković 1963a, 1977a; Vuković & Ivanović 1971; Vuković & Sofradžija 1987), it is rarely considered in current fisheries policy studies in BiH (Drešković *et al.* 2011). Nevertheless, this high proportion of endemism requires additional confirmation because the taxonomy of several species remains uncertain, such as in the genus of *Salmo* (e.g., Georgiev 2011; Razpet *et al.* 2007).




Fig. 3. Number of species presented in both the Danube River Basin and the Adriatic Basin, in Bosnia and Herzegovina.

Native and translocated species

Generally, the ichthyofauna of BiH shows significant differences in biodiversity by catchment basins. Species assemblage inhabiting the Danube River Basin in BiH are similar to the one of the Danube River Basin in Central Europe (Kottelat & Freyhof 2007). Among the 57 native species inhabiting the Danube River Basin, some are particularly interesting, like *Cobitis elongata*, and *Hucho hucho*, mostly recorded in the broader area of the Sava River catchment. The native ichthyofauna of the Adriatic Basin, in turn, is mostly endemic to small parts of the freshwater catchments.

Translocated species recorded in the Adriatic Basin are native species transferred from the Danube River Basin (e.g., *Sander lucioperca*, *Silurus glanis*) (Glamuzina *et al.* 2017). Their occurrence is a result of human intervention, usually the result of legally regulated or unconscionable stocking of target species or accidental introduction during stocking. Some of these species have not established self-sustainable populations, like *Gymnocephalus cernuus*, and *Oncorhynchus mykiss* (Dulčić *et al.* 2005; Glamuzina *et al.* 2017). However, where introduced, translocated species often turned out to be invasive and detrimental to native fish communities (Glamuzina *et al.* 2017).

Endemic species

The freshwater fishes, including lampreys, of BiH is characterised by a considerable species richness and diversity, with the presence of several endemic species with a restricted distribution area, making them more vulnerable to extinction. In total, 38 endemic species have been recorded in the freshwaters of BiH to date (Supp. file 1). At present, four species appear to be endemic only to BiH: *Cobitis herzegoviniensis*, *Phoxinellus pseudalepidotus*, *Telestes dabar* and *T. metohiensis*, all found in the Neretva River catchment (Adriatic Basin). The number of regionally endemic species in BiH is comparable with those in neighbouring countries (Slovenia 12, Montenegro 24, Albania, 43, Croatia 44) (Crivelli 1996). Such an abundance of endemic species in BiH can be primarily attributed to its geographic location and isolation from other European river systems, as well as the complex geologic history and climatic characteristics of the region (Perea *et al.* 2010). The Neretva River catchment, as part of the Adriatic Basin, has a prominent place in the BiH ichthyofauna (Vuković & Sofradžija 1987). This system is comprised of two main catchments, drained by the Neretva and Trebišnjica rivers. The Neretva River catchment has one of the highest endemism of freshwater fishes in this part of the Balkan Peninsula (Economidis & Bănărescu 1991), housing nearly all the country's endemic species. The discovery of a new mitochondrial CR haplotype of *Cottus gobio* and confirmation of native *Salmo trutta*, *Barbatula barbatula*, and *Phoxinus* sp., by Neuburg *et al.* (2023) highlight the importance of the upper Neretva as a refuge and source of genetic diversity, emphasizing the importance of this stretch of the river as a biodiversity hotspot. Unfortunately, the entire endemic fish assemblage there faces significant conservation threats due to major anthropogenic alterations such as damming processes, habitat degradation, pollution, and the introduction of non-native species (Tutman *et al.* 2012a; Glamuzina *et al.* 2017). Nearly 90% of the endemic species have been categorized as threatened according to the IUCN (2025) (Table 2). According to IUCN (2025), at least ten species are Critically Endangered (CR): *Acipenser naccarii*, *Lampetra soljani*, *Salmo marmoratus*, *S. farioides*, *S. obtusirostris*, *Orsinigobius croaticus*, *Chondrostomus knerii*, *T. metohiensis*, *T. dabar* and *Delminichtys ghetaldii* (Freyhof & Brooks 2011; Bogutskaya *et al.* 2012). *Telestes metohiensis*, *T. dabar* and *D. ghetaldii* migrate underground during the low waters in the summer. Even though these three minnows are adapted to stagnant waters (e.g., Lake Bileća, Hamzić, personal observation), in other parts of their distributional areas there have been major declines in population density, like in the Popovo Polje field. For all these 10 Critically Endangered species little is known about their life history, migration pathways, biological traits and spawning activities. Therefore, more comprehensive ichthyological surveys of these species are necessary, providing a baseline for effective protection measures such as establishing ichthyological reserves, and ensuring adequate habitat for the recovery of their populations. Unfortunately, to date, BiH has not yet compiled a list of threatened species, nor is there a Red

Book of freshwater fishes at the national level, and no strategies have been implemented for conservation or protection of the lampreys and other fish fauna (Tutman *et al.* 2017a, 2017b, 2020a, 2020b).

Recently described species and new records within hydrographic catchments

Recently described lampreys and fish species for BiH freshwaters are *Phoxinellus pseudalepidotus* (Bogutskaya & Zupančič 2003), *Knipowitschia radovici* (Kovačić 2005), *Alburnus neretvae* (Buj *et al.* 2010), *Telestes dabar* (Bogutskaya *et al.* 2012), *Cobitis herzegoviniensis* (Buj *et al.* 2014), *Phoxinus karsticus* (Bianco & De Bonis 2015) and *Lampetra soljani* (Tutman *et al.* 2017a). These recent discoveries indicate how diverse and remarkable the freshwater ichthyofauna is in BiH, and suggest that there is still more to be discovered in the lakes, rivers and wetlands of the country. Those species were described by 17 scientists from eight European countries, without the participation of scientists or scientific institutions from BiH. Although BiH has not yet ratified the Nagoya Protocol, procedures regarding the permissions for field work, sampling, and transboundary export of BiH natural resources (as whole individuals, tissues, or extracted DNA) are prescribed by law. Even though international collaborations are beneficial as they decrease the knowledge gap concerning biodiversity in BiH, it is important that all research groups strictly follow national legal requirements and do not take advantage of the country's porous borders. Foreign scientists should adhere to the principles of responsible research and respect BiH's right to fair access and benefit sharing resulting from its resources.

For some species, like gobiids, it seems that the lack of adequate sampling techniques has resulted in this group being previously overlooked in the freshwaters of BiH (Šanda & Kovačić 2009; Tutman *et al.* 2009a, 2012b, 2013, 2021). Here, we provided new knowledge about the distributional range of some species of the family, again reinforcing that more studies on the group are necessary in BiH.

Non-native species

The introduction of non-native freshwater fish species induced by humans potentially represents one of the most serious threats to biodiversity and ecosystem integrity in different parts of Europe (Ribeiro & Leunda 2012; Costa *et al.* 2021) and the world (Clavero & García-Berthou 2005; Helfman 2007). Non-native species have been implicated in the decline and extinction of native freshwater fauna worldwide and are considered a major threat to biodiversity (Taylor *et al.* 1984; Lever 1996). These species usually have a high potential for dispersion and occupy niches of native species due to their competitive advantages and the absence of predators (Kolar & Lodge 2001). They are often blamed for adverse ecological impacts on native fishes such as competition for food and space, predation, import and spread of pathogens and parasites, habitat alteration, and even their extinction (Bernery *et al.* 2022).

Currently, 23 non-native fish species are recorded in BiH freshwaters (Supp. file 1). The fact that the number of introduced species in the Neretva River catchment (17 non-native and 18 translocated) is the same as the number of native freshwater species (35) ever recorded in the Adriatic Basin and higher than the number of endemic species (28) is particularly alarming (Vukić *et al.* 2019). As with most aspects of BiH freshwater fish biogeography, the ecology and conservation impacts of non-native species are largely unknown. Most non-natives were introduced to benefit the commercial value of waters in freshwater fisheries (e.g., *Hypophthalmichthys molitrix*, *H. nobilis*, *Micropterus salmoides*) or for aquaculture purposes (e.g., *Onchorhynchus mykiss*, *Salvelinus alpinus*, *S. fontinalis*). A recent example of such an introduction is *Coregonus peled* (Hamzić *et al.* 2011). In addition to intentionally stocked species, several other species were likely introduced accidentally due to poor regulation and control. This includes *Lepomis gibbosus*, *Carassius gibelio* and *Pseudorasbora parva*, which have spread widely and potentially pose the greatest threat to native species, especially in the Adriatic Basin (Glamuzina *et al.* 2017). Non-native species have experienced different fates in the region, but some have established self-sustaining populations, such as *L. gibbosus*, *P. parva* and *Leucaspis delineatus* (Glamuzina *et al.* 2017). Introductions continue to this day,

and *Carassius langsdorfi* (Kalous *et al.* 2012), for instance, is now present in the Adriatic Basin, whereas *M. salmoides* (Tutman *et al.* 2017c) is recorded in both the Adriatic and Danube Basins.

Taxonomic considerations

Many unresolved challenges concerning the phylogenetic placement and taxonomic validity of several fish species still remain, highlighting how still insufficiently is our knowledge on the BiH freshwater fish fauna. This is specifically true for the species of the Adriatic Basin, which is rich in endemics (Palandačić *et al.* 2015, 2017; Reier *et al.* 2022; Lukač *et al.* 2023, 2024). Advancing the phylogeny and taxonomy of the genera *Alburnus*, *Delminichthys*, *Phoxinus*, *Rutilus*, *Telestes*, *Squalius* and *Salmo*, which are widespread in the karst area of eastern Herzegovina (Ketmaier *et al.* 2004; Buj *et al.* 2017, 2020a, 2020b, 2022; Charmpila *et al.* 2025), has been a long-standing task. The taxonomic status of the genera *Gobio* and *Romanogobio* (Takács *et al.* 2014; Zangl *et al.* 2020), as well as the *Barbatula* (Calegari *et al.* 2025) from the Danube Basin also requires a thorough revision.

The taxonomic status of the populations of *Alburnus* in BiH has also not yet been resolved (Buj *et al.* 2010; Vucić *et al.* 2017). The population from the Neretva River catchment is now recognized as a distinct species *A. neretvae*; therefore, it seems that *A. arborella* is likely absent from BiH (Buj *et al.* 2010). Additional systematic research is also necessary for the populations in the Sava catchment, Danube Basin (Vucić *et al.* 2017). Furthermore, the taxonomy of *Rutilus* and species of the Leuciscinae of northern Italy, the Western Balkans, and southern Greece has not been comprehensively studied (Ketmaier *et al.* 2008). A significant intraspecific difference within southern European roaches was noticed (Bianco & Ketmaier 2014; Geiger *et al.* 2014), and the taxonomy of the genera *Rutilus* and *Leucos* is debated with no agreement on the validity and distribution of their species (Petrosino *et al.* 2022). There is a lack of consensus regarding the validity of several species inhabiting the freshwaters of the Western Balkans and BiH (Geiger *et al.* 2014). Subpopulations occurring in this region are either treated as separate, locally endemic species such as in *Telestes*, *Rutilus*, i.e., *R. basak* from the Neretva River system (Geiger *et al.* 2014; Buj *et al.* 2017, 2020a, 2022; Petrosino *et al.* 2022), or considered as being merely subpopulations of more widespread species, such as *Leucos basak* (Bianco & Ketmaier 2014; Pietrock *et al.* 2022). Further research is indeed needed because at present molecular analyses do not fully support their separation into distinct species (Perea *et al.* 2010; Geiger *et al.* 2014; Schönhuth *et al.* 2018).

A similar situation has been observed with the species previously belonging to the genera *Phoxinellus* and *Paraphoxinus*, which were later divided into the genera *Delminichthys* and *Telestes* (Palandačić *et al.* 2015, 2017, 2024; Reier *et al.* 2022; Lukač *et al.* 2023, 2024). However, according to molecular data some specimens identified as *T. dabar* actually represent *T. metohiensis* (Buj *et al.* 2017), while Reier *et al.* (2022) recognized both species as distinct based on molecular data. Therefore, this taxonomic question needs to be more fully clarified and the validity of the *T. dabar* population in the Dabarsko Polje should be properly tested.

Another taxonomic puzzle includes the populations of *Phoxinus phoxinus* sensu lato. According to Kottelat & Freyhof (2007), two species of European minnow occur in the western Balkan Peninsula, with *P. phoxinus* in the Danubian and *P. lumaireul* in the Adriatic part of the peninsula. However, more recent evidence indicates high levels of molecular and anatomical diversity in both species (Palandačić *et al.* 2015). This finding indicates that a multispecies complex of *Phoxinus* is present (Palandačić *et al.* 2015, 2017, 2020; Vucić *et al.* 2018). Neuburg *et al.* (2023) confirmed that the minnows from the upper Neretva River belong to clade 2 of Palandačić *et al.* (2015), comprising populations from the lower Sava catchment (Danube Basin) and rivers flowing directly into the Adriatic Sea (Adriatic Basin) (Palandačić *et al.* 2015, 2017, 2020). The occurrence in the same Basin of two other species of this genus, *P. ketmaieri* and *P. karsticus*, was proposed based only on anatomy. However, Palandačić *et al.* (2017) concluded that *P. ketmaieri* is not valid based on molecular species delimitation techniques, and *P. ketmaieri* was synonymised with *P. lumaireul*. They also concluded that *P. karsticus* forms a clearly distinct group in both mtDNA analysis and RAG1 networks. The same

authors further indicate that, due to taxonomic complexities of the genus *Phoxinus*, species descriptions and synonymisation in the genus should ideally include morphological and molecular data. In sum, more research is needed to confirm the presence of *P. carsticus* in BiH.

A similar complex taxonomic situation is present in the case of the species previously identified as *Salmo dentex* (Snoj *et al.* 2010). The validity of *S. fariooides* has always been questionable due to inadequate original descriptions, rare records and the absence of type specimens (Kottelat 1997). Analysis indicates that the trouts of the Neretva River are genetically heterogeneous, with the single exception of members of *S. obtusirostris*, which have its own distinct set of mtDNA and nuclear DNA markers (Razpet *et al.* 2007).

Furthermore, *Knipowitschia radovici* is a relatively recently described stenoendemic species that inhabits only the lower Neretva River catchment in Croatia (tributary Norin River; Kovačić 2005) and BiH (Hutovo Blato wetland; Šanda & Kovačić 2009; Tutman *et al.* 2013). However, in a recent study that applied molecular tools, Tougard *et al.* (2021) provided new evidence that among the endemic Adriatic species of *Knipowitschia*, *K. panizzae*, *K. radovici* and *K. mrakovcici* were genetically highly similar, although they are regarded as distinct species based on morphology. Based on the high genetic similarity and species delimitation tests, they proposed that *K. radovici* and *K. mrakovcici* are junior synonyms of *K. panizzae*.

Threats, legislation and protection

Regarding nature protection in BiH, the competence lies in both the Federation of Bosnia and Herzegovina and Republic of Srpska, in addition to the Brčko District as three political entities of BiH. Implementing international laws is a competence of the Ministry of Foreign Trade and Economic Relations of BiH at the state level. The Red Lists of Fauna are adopted at the entity level. The Red List of the fauna of the Federation of BiH (Official Gazette of the Federation of Bosnia and Herzegovina 7/14) contains 115 freshwater and marine fishes. Considering exclusively freshwater fish species, seven were classified as Critically Endangered (CR), 16 as Endangered (EN), seven as Vulnerable (VU), and seven are Data Deficient (DD) (Škrijelj *et al.* 2013; Dekić *et al.* 2024). The Red List of Protected Species of Flora and Fauna of the Republic of Srpska lists two species of lampreys and 46 species of actinopterygians without clear threat categories (Dekić *et al.* 2024). Unfortunately, the national assessment of the threat risk of several species of fish in BiH has not yet been thoroughly carried out, and no conservation strategies or protection measures have been implemented. More detailed information can be found in Dekić *et al.* (2024). However, a global assessment following IUCN criteria has been conducted for the majority of BiH fishes (Table 2). *Anguilla anguilla* is classified as Critically Endangered (CR), while 15, mostly endemic species, in BiH are considered Endangered (EN) according to the IUCN, with a significant anthropological impacts threatening their conservation. All of them are regionally endemic with a restricted range, distributed in a single river, river catchment or in several localities in a very small area (Buj *et al.* 2014, 2015a, 2015b), implying the necessity of immediate effective protection. Furthermore, according to the *Regulation on strictly protected and protected wild species* (Official Gazette of the Republic of Srpska 65/20), there are 31 species of fishes that are in the category of Strictly Protected, and 20 species that are in the category of Protected in the Republic of Srpska. In comparison, nine fish species are Strictly Protected, and 13 species of fish are Protected in the Federation of Bosnia and Herzegovina following the *Rules on protection measures for protected and strictly protected species and subspecies* (Official Gazette of the Federation of Bosnia and Herzegovina 21/20). The conservation of protected and strictly protected wild species, except for species protected by fishing regulations, is implemented by prohibiting the use, destruction, and undertaking of all activities that may endanger these species and their habitats, as well as by undertaking measures and activities to manage populations (Dekić *et al.* 2024).

Water pollution and habitat loss through modification of river substrates are among the major threats to freshwater fishes, including lampreys, in BiH (Hamzić, 2002; Tutman *et al.* 2017b, 2020a, 2020b, Palandačić *et al.* 2024). The introduction of non-native species, especially predatory ones, can be added to these threats

(Kosorić & Kardoš 1955; Glamuzina *et al.* 2017; Dekić *et al.* 2024; Hamzić 2024; Palandačić *et al.* 2024). The construction of hydroelectric dams along the Neretva, Drina and Vrbas rivers since the 1950s has altered the natural flow and sedimentation regimes and potentially blocked migration routes, fragmented subpopulations, and markedly reduced the extent of habitat suitable for all life stages (Glamuzina *et al.* 2017). Diffuse and point source agricultural, domestic and industrial pollution have resulted in eutrophication or discharge of toxic substances at some locations (Has-Schön *et al.* 2008). The industrial extraction of riverine gravel and sand for urban development has taken place since the 1970s, causing additional impacts on river bed and habitat degradation in areas such as the Mostarsko blato that affected the life cycle of several species (Glamuzina *et al.* 2017). Additional threat to the lower Neretva River is the intense saltwater intrusion as a result of hydrological regime changes of freshwater inflows, which can cause increased intake of brackish and marine organisms (Tutman *et al.* 2021). The Hutovo Blato wetland is potentially the most endangered habitat for a number of endemic species of fish. Although no specific conservation actions are in place, part of the range of some species lies within the boundaries of protected areas like Hutovo Blato wetland in the lower Neretva River catchment (Tutman *et al.* 2002, 2006, 2008a, 2009b, 2012a, 2012c, 2018, 2019) and the Blidinje Nature Park (Carosi *et al.* 2024), which have been included in the EU Natura 2000 network. Despite numerous ichthyological surveys conducted in the region of the Danube River and the Adriatic Basin, there are substantial gaps in the knowledge of the current taxonomic status and geographical distribution of freshwater fishes, including lampreys, in BiH. Except for Dekić *et al.* (2024) and Hamzić (2024), no information has been presented to address the vulnerability of threatened endemic species in BiH.

Despite the new information presented here, the knowledge of the taxonomy and distribution of several freshwater fishes in BiH remains insufficient. In particular, data on the biology and ecology of most endemic species and their distribution in BiH is very scarce and based on general notes and historical data. Given these gaps, this study can in no way be considered a definitive checklist of the freshwater fishes, including lampreys, of BiH, though hopes are that the data presented herein will serve as the foundation for further analysis and conservation actions. This highlights the importance of establishing conservation measures, considering the high faunistic and scientific value of the species inhabiting the BiH freshwaters (Tutman *et al.* 2017b, 2020a, 2020b).

Acknowledgements

This study is dedicated to Prof. Dr Tonko Šoljan and Prof. Dr. Tihomir Vuković, members of the Academy of Sciences of Bosnia and Herzegovina, honorable scientists and lecturers from the Faculty of Science and Mathematics, University of Sarajevo, Bosnia and Herzegovina. The first author also expresses his gratitude to all the people, professors and students, and the time spent in the city of Sarajevo during studies at the Department of Biology, Faculty of Science and Mathematics, University of Sarajevo. We are thankful to Linda Zanella for improving the English in the manuscript. Radek Šanda was supported by the Ministry of Culture of the Czech Republic (DKRVO 2024-2028/6.I.c, National Museum, 00023272).

References

- Adrović A. 2007. *Biodiverzitet i Ekološke Osobnosti Ihtiopopulacija Hidroakumulacije Modrac*. PhD thesis, University of Tuzla, Sarajevo.
- Adrović A. 2012. *Ribe Modraca. Ihtiološka Monografija*. NAM, Tuzla.
- Adrović A. 2018. *Biodiverzitet Ihtiopopulacija Rijeke Gostelje*. OFF-SET, Tuzla.
- Adrović A. & Skenderović I. 2007. Allochthonous Ichthyofauna of Certain Water Bodies in North–Eastern Bosnia. *Acta Agriculturae Serbica* 12 (23): 37–46.
- Adrović A., Škrijelj R. & Đug S. 2008. Ekološke i biosistematske karakteristike vrste *Cobitis elongatoides* Bacescu & Maier, 1969. iz tekućica Brke, Gostelje i Tinje u slivu rijeke Save. *In*: Petrović M., Petrović M.

& Marković D. (eds) *XIV Savjetovanje o Biotehnologiji*: 359–366. University of Kragujevac, Faculty of Agronomy Čačak, Čačak.

Adrović A., Škrijelj R. & Skenderović I. 2009a. The first record and morphological characteristics of hybrid of *Rutilus rutilus* × *Abramis brama* from Modrac reservoir, Bosnia and Herzegovina. *Croatian Journal of Fisheries* 67 (1): 11–23.

Adrović A., Škrijelj R. & Skenderović I. 2009b. Ecological characteristics of the roach (*Rutilus rutilus* L.) and bream (*Abramis brama* L.) in the reservoir of the Lake Modrac. *Acta Agriculturae Serbica* 14 (27): 23–33.

Adrović A., Skenderović I., Salihović J. & Stjepić Š. 2012. Biodiversity of fish fauna of the Gostelja River and reservoir Modrac. Structure and dynamics of ecosystems Dinarides – status, possibilities and prospects. In: Redžić S (ed.) *Proceedings of the International Conference Structure and Dynamics of Ecosystems Dinarides – Status, Possibilities and Prospects, Special Edition CXLIX* 23: 277–290. Academy of Sciences and Arts of Bosnia and Herzegovina, Department of Natural Sciences and Mathematics, Sarajevo. <https://doi.org/10.5644/proc.eco-03.21>

Adrović A., Golub D. & Dekić R. 2017. Distribution of an invasive fish species *Pseudorasbora parva* (Telostei: Cyprinidae) in Bosnia and Herzegovina. In: Pešić V. & Hadžiablahović S. (eds) *Book of Abstracts of 7th International Symposium of (ISEM7)*: 94. Institute for Biodiversity and Ecology, Sutomore.

Adrović A., Bajrić A., Hajdarević E. & Skenderović I. 2018. Length-weight relationships and condition of some cyprinid fish species from waters of north-east Bosnia. In: Miletić N. & Marković D. (eds) *XXIII Savjetovanje o Biotehnologiji*: 543–548. University of Kragujevac, Faculty of Agriculture Čačak, Čačak.

Adrović A., Kamberović J., Nešković R., Lukić Z. & Jusufović A. 2021. *Biodiverzitet Jezera Kop Šićki Brod*. Centar za ekologiju i energiju, Tuzla.

Aganović M. 1952a. Hutovo blato. *Ribarstvo Jugoslavije* 3: 28–30.

Aganović M. 1952b. Zaštita lipljena i njegov odnos prema pastrmki. *Ribarski list BiH* 6: 7–14.

Aganović M. 1952c. Proučavanje pitanja minimalnih mjera za lipljena u rijekama Drini, Bosni i Vrbasu. *Ribarski List BiH* 6: 15–18.

Aganović M. 1957. Veliko Plivsko Jezero. *Ribarstvo Jugoslavije* 1: 4–9.

Aganović M. 1965a. Komparativna istraživanja režima ishrane, rasta, plodnosti i strukture populacija lipljena u rijekama Bosni i Plivi. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 18: 3–109.

Aganović M. 1965b. O lipljenu rijeke Plive. *Ribarstvo Jugoslavije* 4: 76–77.

Aganović M. 1967. Uzrasna struktura populacija riba Velikog i Malog Plivskog jezera. *Ribarstvo Jugoslavije* 1: 1–6.

Aganović M. 1969. Sastav ribljih populacija rijeke Jaruge. *Ichthyologia* 1: 3–10.

Aganović M. 1970. Plodnost oštrulja (*Aulopyge hüegeli* Heck) iz voda Livanjskog polja. *Ribarstvo Jugoslavije* 1: 15–18.

Aganović M. & Kapetanović N. 1963. Prilog diskusiji o udomaćivanju dužičaste pastrmke u vodama Jugoslavije. *Poljoprivredni Pregled* 12: 7–8.

Aganović M. & Kapetanović N. 1970. Odnos dužine crijevnog trakta i dužine tijela kod tri vrste roda *Paraphoxinus*. *Ichthyologia* 2: 17–24.

Aganović M. & Kapetanović N. 1971. Dužinsko i težinsko rastezanje kod dviju endemskih ribljih vrsta roda *Paraphoxinus*. *Ribarstvo Jugoslavije* 26 (4): 69–71.

- Aganović M. & Kapetanović N. 1973. Prilog upoznavanju uticaja otpadnih voda kožarskih industrija na mješvitu riblju populaciju u nekim vodama BiH. *Ribarstvo Jugoslavije* 3: 51–55.
- Aganović M. & Kapetanović N. 1978. Uzrasna struktura u populacijama nekih vrsta riba iz rijeke Neretve. *Ichthyologia* 10: 1–7.
- Aganović M. & Milošević B. 1959. Prilog ribarsko-biološkom poznavanju slivnog područja rijeke Gostović. *Ribarstvo Jugoslavije* 1: 12–17.
- Aganović M. & Vuković T. 1966. Odnos dužine crijevnog trakta i dužine tijela kod tri lokalne populacije oštrulja (*Aulopyge hügelii* Heck.). *Ribarstvo Jugoslavije* 21: 8–11.
- Aganović M. & Vuković T. 1967. Neki morfometrijski i meristički karakteri oštrulja iz vodotoka na području Livanjskog polja. In: Ržničanin B. (ed.) *I Simpozijum JID, Rezime „Informativni Bilten JID“* 2: 14–15. Jugoslovensko ihtiološko društvo, Sarajevo.
- Aganović M. & Vuković T. 1968. Uzrasno variranje nekih morfometrijskih karaktera oštrulja iz vodotoka sa područja Livanjskog polja. *Ribarstvo Jugoslavije* 23: 30–33.
- Aganović M. & Vuković T. 1971. Promjenjivost morfološko-taksonomskih karaktera u lokalnim populacijama oštrulja *Aulopyge hügelii* Heck. (Pisces, Cyprinidae). In: *I Jugoslavenski Simpozijum iz Sistematike, Posebni Otisak*. Sarajevo.
- Aganović M., Vuković T. & Kapetanović N. 1966. Ihtiofauna jablaničke vodene akumulacije. *Ribarstvo Jugoslavije* 5: 92–95.
- Aganović M., Kapetanović N. & Veleđar I. 1974. Plodnost *Leuciscus turskyi turskyi* (Heckel, 1843). *Ichthyologia* 6: 1–6.
- Aganović M., Kapetanović N. & Veleđar I. 1976. Odnos dužine crijevnog trakta i dužine tijela bez „C“ kod *Leuciscus turskyi turskyi* Heckel iz vodene akumulacije Buško blato. *Ichthyologia* 8: 1–7.
- Ajanović N., Moth-Poulsen T., Peteri A., Wurzbacher S. & Boellerup L. 2010. Fish hatchery in the municipality of Bosanska Krupa in northwestern Bosnia and Herzegovina: a sustainable development pilot project. *Ribarstvo* 68 (2): 79–92.
- Alić B., Milanović A., Čaklović F. & Saračević L. 2004. Content of the copper, zinc, cadmium, lead and mercury in muscle tissue of the Trout (*Salmo trutta* m. *fario* L.) and Nases (*Chondrostoma nasus* L.) caught in rivers Una, Vrbas and Drina. *Meso* 4 (3): 40–46.
- Anonymous 1886. Ribarstvo i rakolov. In: Krešić M. (ed.) *Izvjješće Trgovačko-obrtničke komore u Zagrebu 1886*: 256–259. Trgovačka i obrtnička komora, Zagreb.
- Anonymous 1898. Uzgoj jesetri iz Hrvatske u Rhoni. *Lovačko-Ribarski Viestnik* 7: 87.
- Anonymous 1903. Glavatica i „Lachsforelle“. *Lovačko-Ribarski Viestnik* 12: 71.
- Anonymous 1909. Kako da se podigne ribarstvo? *Lovačko-Ribarski Viestnik* 18: 99–101.
- Anonymous 1914. Losos na ribljem trgu u Zagrebu. *Lovačko-Ribarski Vjesnik* 23: 34–36.
- Anonymous 1931. Gorostasni somovi. *Priroda* 2–3: 91–92.
- Anonymous 1947a. Neobično krupan šaran. *Ribarstvo Jugoslavije* 4: 46.
- Anonymous 1947b. Ribarske prilike u Brčkom i okolini. *Ribarstvo Jugoslavije* 4: 46.
- Anonymous 1948. Šaran od 18 kg. *Ribarstvo Jugoslavije* 7–8: 78.
- Anonymous 1959. Sitne vijesti. *Ribarski list* 1: 23–24.
- Anonymous 1965. Jegulje u Savi, Dunavu i Tisi. *Ribarski List* 5: 107.

- Avdić S., Bakrač A., Adrović A., Hajdarević E., Dekić R. & Delić M. 2023. Seasonal variations of morphometric and hematological parameters in Dalmatian barbelgudgeon *Aulopyge huegelii* Heckel 1843 from Šatorsko Lake. *Croatian Journal of Fisheries* 81: 159–166. <https://doi.org/10.2478/cjf-2023-0017>
- Babić S. & Mikavica D.E. 1994. Ecological characteristics of rivers in the region of Sarajevo (the Fojnica, the Željeznica and the Zujevina). *Zemljište i biljka* 43 (1): 57–66.
- Bajrić A., Adrović A., Hajdarević E. & Skenderović I. 2015. Ishrana i faktor kondicije klijena (*Squalius cephalus*) iz rijeke Spreče. In: Nikolić M., Marković D., Ilić R. & Đurović I. (eds) *XX Savetovanje u Biotehnologiji, Zbornik Radova* 20 (22): 411–415. University of Kragujevac faculty of agronomy, Čačak.
- Bajrić A., Adrović A., Skenderović I. & Hajdarević E. 2016. Comparative review of morphometric and meristic characteristics of Bream (*Abramis brama*, Linnaeus) and Roach (*Rutilus rutilus*, Linnaeus) and their hybrids from the reservoir Modrac. In: Golub D. (ed.) *Zbornik Radova III Simpozijuma Biologa i Ekologa Republike Srpske (SBERS 2015)* 7 (2): 107–113. Prirodno-matematički fakultet Univerziteta u Banjoj Luci, Banja Luka.
- Bajrić A., Adrović A., Hajdarević E. & Skenderović I. 2017. Taksonomske karakteristike klijena (*Squalius cephalus*) iz voda sjeveroistočne Bosne. *Educa* 10 (10): 3–7.
- Bajrić A., Adrović A., Hajdarević E., Skenderović I. & Tanović E. 2018a. Body pigmentation and meristic characteristics of Balkan golden loach (*Sabanejewia balcanica*) from the water catchment area of the river Sava. *Croatian Journal of Fisheries* 76 (2): 72–79. <https://doi.org/10.2478/cjf-2018-0009>
- Bajrić A., Adrović A., Hajdarević E., Skenderović I., Tanović E. & Marković G. 2018b. Karakteristike krvnih ćelija *Sabanejewia balcanica* (Cobitidae) iz nekih tekućica slivnog područja rijeke Save. In: Miletić N. & Marković D. (eds) *XXIII Savjetovanje o Biotehnologiji sa Međunarodnim Učešćem, Zbornik Radova*: 215–221. Univerzitet u Kragujevcu, Agronomski fakultet, Čačak.
- Bajrić A., Hajdarević E., Adrović A. & Skenderović I. 2020. Age and Growth of the Chub (*Squalius cephalus*) from Watercourses of Northeast Bosnia. *Educa* 13: 9–12.
- Bajrić A., Buj I., Adrović A. & Hajdarević E. 2021a. Morphological and genetic diversity of *Sabanejewia balcanica* (Cobitidae, Actinopterygii) in Bosnia and Herzegovina. *Journal of Applied Ichthyology* 37 (1): 89–98. <https://doi.org/10.1111/jai.14130>
- Bajrić A., Hajdarević E. & Adrović A. 2021b. Systematic position and vulnerability of *Sabanejewia balcanica* in the Balkan area. *Works of the Faculty of Forestry University of Sarajevo* 50 (1): 19–26. <https://doi.org/10.54652/rsf.2020.v50.i1.38>
- Bakrač A., Avdić S., Dekić R. & Ičanović I. 2020a. Organosomatic indices and leukocyte formula of chub (*Squalius cephalus* L) from the Suturlija River. *Agro-knowledge Journal* 21 (4): 151–160. <https://doi.org/10.7251/10.7251/AGREN2004151B>
- Bakrač A., Džafić S., Dekić R. & Ičanović I. 2020b. The presence of heavy metals in water and the muscle tissue of grayling (*Thymallus thymallus*) of the Una River. *Agro-knowledge Journal* 21 (3): 107–115. <https://doi.org/10.7251/AGREN2003107B>
- Bakrač-Bećiraj A. 2008. *Fiziologija Ishrane i Prirasta Vrste Thymallus thymallus (Linnaeus, 1758) u Prirodnim i Eksperimentalnim Uslovima*. PhD thesis, University of Banja Luka, Banja Luka.
- Bănăduc D. & Curtean-Bănăduc A. 2014. *Rhodeus sericeus amarus* Bloch, 1782; Monitoring elements in the new NATURA 2000 context in Croatia. *Transylvanian Review of Systematical and Ecological Research* 16 (1): 185–204. <https://doi.org/10.1515/trser-2015-0013>
- Bănărescu P. 1989. Zoogeography and history of the freshwater fish fauna of Europe. In: Holčík J. (ed.) *The Freshwater Fishes of Europe* 1. Aula-Verlag, Wiesbaden.

- Bănărescu P. 1992. *Zoogeography of Freshwaters. Vol. 2: Distribution and Dispersal of Freshwater Animals in North America and Eurasia*. Aula Verlag, Wiesbaden.
- Bănărescu P. & Herzig-Straschil B. 1998. Beitrag zur Kenntnis der *Leuciscus*-Untergattung *Telestes* Bonaparte (Pisces, Cyprinidae). *Annalen des naturhistorischen Museums in Wien, Serie B, Botanik und Zoologie* 100B: 405–424.
- Bănărescu P. & Nalbant T.T. 1957. Eidonomische und taxonomische Untersuchungen an *Cobitis elongata* (Pisces, Cobitidae). *Senckenbergiana Biologica* 38: 283–294.
- Bănărescu P. & Nalbant T.T. 1998. *Cobitis elongata* Heckel & Kner, 1858 (Pisces, Ostariophysi, Cobitidae), distribution, relationships, geographical variation and protection status. *Revue Roumaine de Biologie, Biologie Animale* 43: 57–65.
- Barbieri R., Zogaris S., Kalogianni E., Stoumboudi M.Th., Chatzinikolaou Y., Giakoumi S., Kapakos Y., Kommatas D., Koutsikos N., Tachos V., Vardakas L. & Economou. A.N. 2015. *Freshwater Fishes and Lampreys of Greece: An Annotated Checklist*. Monographs on Marine Sciences No. 8, Hellenic Centre for Marine Research, Athens.
- Bartulović V., Dulčić J., Bogut I., Pavličević J., Hasković E. & Glamuzina B. 2011. First record of the freshwater bream, *Abramis brama* in the river Mala Neretva, Adriatic Sea basin system of Croatia. *Cybium* 35 (2): 165–166.
- Bašić M. 1964. *Paraphoxinus metohiensis* Steind. *Gatačkog i Nevesinjskog Polja*. Diploma thesis, University of Sarajevo, Sarajevo.
- Bećiraj A. & Šahinović R. 2012. Biodiversity of the fish of the Klokot River. In: Redžić S. (ed.) *Proceedings of the Second International Colloquium “Biodiversity – Theoretical And Practical Aspects”, Special Editions CXLVIII*: 143–157. Department of Natural Sciences and Mathematics, Academy of Sciences and Arts of Bosnia and Herzegovina, Sarajevo.
- Bećiraj A., Ivanc A., Piria M. & Dekić R. 2008. Food composition of Grayling *Thymallus thymallus* L., from the river Krušnica. *Croatian Journal of Fisheries* 66 (3): 105–118.
- Bećiraj A., Tahrić E., Šarić-Kadić I. & Ičanović I. 2016. Fish population from Klokot and Krušnica rivers. In: Poleksić V., Rašković B. & Marković Z. (eds) *Proceedings of VII International conference “Water & Fish”*: 191–197. University of Belgrade, Faculty of Agriculture, Serbia.
- Bek B. 1935. Sa ostvama na Uni. *Lovačko-Ribarski Vjesnik* 44: 415–417.
- Benovics M., Kičinjaová M.L. & Šimková A. 2017. The phylogenetic position of the enigmatic Balkan *Aulopyge huegelii* (Teleostei: Cyprinidae) from the perspective of hostspecific *Dactylogyrus* parasites (Monogenea), with a description of *Dactylogyrus omenti* n. sp. *Parasites & Vectors* 10: 547. <https://doi.org/10.1186/s13071-017-2491-z>
- Benovics M., Desdevises Y., Vukić J., Šanda R. & Šimková A. 2018. The phylogenetic relationships and species richness of host-specific *Dactylogyrus* parasites shaped by the biogeography of Balkan cyprinids. *Scientific Reports* 8: 13006.
- Benovics M., Desdevises Y., Šanda R., Vukić J. & Šimková A. 2019. Cophylogenetic relationships between *Dactylogyrus* (Monogenea) ectoparasites and endemic cyprinoids of the north-eastern European peri-Mediterranean region. *Journal of Zoological Systematics and Evolutionary Research* 58 (1): 1–21. <https://doi.org/10.1111/jzs.12341>
- Berberović Lj. 1967. Podaci o kariotipovima slatkovodnih riba iz voda Bosne i Hercegovine. *Godišnjak Biološkog instituta Univerziteta u Sarajevu* 12 (1–2): 5–15.

- Berberović Lj. 1980. Kariološke osobine nekih ciprinidnih vrsta i njihova hibridizacija. *Ichthyologia* 12 (1): 1–8.
- Berberović Lj. & Pavlović B. 1970. Basic data about the chromosome set of the species *Paraphoxinus adspersus* (Heckel, 1843). *Bulletin Scientifique Section A*, 15 (11–12): 391–392.
- Berberović Lj. & Sofradžija A. 1972. Pregled podataka o hromosomskim garniturama slatkovodnih riba Jugoslavije. *Ichthyologia* 4 (3): 1–21.
- Berberović Lj. & Sofradžija A. 1974a. Basic data about diploid chromosome complements of *Phoxinus* (Cyprinidae, Pisces). *Bulletin Scientifique Section A*, 19 (5–6): 142–143.
- Berberović Lj. & Sofradžija A. 1974b. Hromosomi vrste *Cottus gobio* L. (Cottidae, Pisces). *Godišnjak Biološkog instituta Univerziteta u Sarajevu* 27: 5–17.
- Berberović Lj., Hadžiselimović R. & Sofradžija A. 1969a. Chromosome number of the species *Paraphoxinus alepidotus* (Heckel). *Bulletin Scientifique Section A*, 14 (11–12): 386.
- Berberović Lj., Hadžiselimović R. & Sofradžija A. 1969b. Prvi rezultati istraživanja hromozomskih garnitura trebinjske i gatačke gaovice (*Paraphoxinus pstrossi* Steind. i *Paraphoxinus metohiensis* Steind.). *Ichthyologia* 1 (1): 11–18.
- Berberović Lj., Hadžiselimović R. & Sofradžija A. 1969c. Chromosome set of the *Aulopyge hügelii* Heckel 1841. *Bulletin Scientifique Section A*, 18 (1–3): 10–12.
- Berberović Lj., Hadžiselimović R. & Sofradžija A. 1970a. Hromozomske garniture podbile (*Chondrostoma phoxinus* Heck.) i plotice (*Rutilus rubilio* Bon.). *Ribarstvo Jugoslavije* 25 (2–3): 40–42.
- Berberović Lj., Hadžiselimović R. & Sofradžija A. 1970b. Usporedni pregled osnovnih podataka o hromosomskim garniturama vrsta *Chondrostoma phoxinus* Heckel i *Chondrostoma kneri* Heckel. *Ichthyologia* 2 (1): 25–30.
- Berberović Lj., Hadžiselimović R. & Sofradžija A. 1970c. Komparativna analiza hromosomskih garnitura podbile (*Chondrostoma phoxinus* Heckel) i podustve (*Chondrostoma kneri* Heckel). In: *Abstrakti 3. Simpozija "Savremena ihtiološka istraživanja u Jugoslaviji"*: 8. Yugoslav Ichthyological Society, Kotor.
- Berberović Lj., Čurić M., Hadžiselimović R. & Sofradžija A. 1970d. Neka zapažanja o citotaksonomskim osobinama vrsta roda *Paraphoxinus*. In: *Abstrakti 3. Simpozija „Savremena Ihtiološka Istraživanja“*: 7. Yugoslav Ichthyological Society, Kotor.
- Berberović Lj., Hadžiselimović R., Pavlović B. & Sofradžija A. 1970e. Hromozomska garnitura neretvanske mekousne (*Salmothymus obtusirostris oxyrhynchus* (Steindachner). *Genetika* 2 (1): 55–63.
- Berberović Lj., Hadžiselimović R., Pavlović B. & Sofradžija A. 1971a. Sistematski položaj vrste *Aulopyge hügelii* Heckel 1841. u svjetlu osnovnih podataka o njenoj hromozomskoj garnituri. In: *I Jugoslavenski Simpozijum iz Sistematike*: 31. Sarajevo.
- Berberović Lj., Hadžiselimović R., Pavlović B. & Sofradžija A. 1971b. Osnovni usporedni podaci o hromosomskim garniturama vrsta *Paraphoxinus adspersus* (Heckel 1843) i *Paraphoxinus croaticus* Steindachner 1865. *Ichthyologia* 3 (1): 3–12.
- Berberović Lj., Hadžiselimović R. & Sofradžija A. 1973. Chromosome set of the species *Aulopyge hügelii* Heckel, 1841. *Bulletin Scientifique Section A* 1–3: 10–11.
- Berberović Lj., Sofradžija A. & Vagner Lj. 1978. Hromosomi linjaka (*Tinca tinca*). *Ichthyologia* 10 (1): 9–15.
- Bernatchez L. 2001. The evolutionary history of brown trout (*Salmo trutta* L.) inferred from phylogeographic, nested clade, and mismatch analyses of mitochondrial DNA variation. *Evolution* 55: 351–379. <https://doi.org/10.1111/j.0014-3820.2001.tb01300.x>

- Bernatchez L., Guyomard R. & Bonhomme F. 1992. DNA sequence variation of the mitochondrial control region among geographically and morphologically remote European brown trout *Salmo trutta* populations. *Molecular Ecology* 1: 161–173. <https://doi.org/10.1111/j.1365-294X.1992.tb00172.x>
- Bern Convention 1979. Convention on the Conservation of European Wildlife and Natural Habitats. Available from <https://www.coe.int/en/web/bern-convention>
- Bernery C., Bellard C., Courchamp F., Brosse S., Gozlan R., Jarić I., Teletchea F. & Leroy B. 2022. Freshwater fish invasions: a comprehensive review. *Annual Review of Ecology, Evolution, and Systematics* 53: 427–456. <https://doi.org/10.1146/annurev-ecolsys-032522-015551>
- Bianco P.G. 1980. Remarks on the genera *Alburnus* and *Alburnoides* and description of a neotype for *Alburnus albidus* (Costa, 1838), senior synonymous of *Alburnus vulturinus* (Costa, 1838) (Pisces, Cyprinidae). *Cybium* 9: 31–44.
- Bianco P.G. 1990. Potential role of the paleohistory of the Mediterranean and Paratethys basins on the early dispersal of Euro-Mediterranean freshwater fishes. *Ichthyological Explorations of Freshwaters* 1: 167–184.
- Bianco P.G. 1995. Mediterranean endemic freshwater fishes of Italy. *Biological Conservation* 72 (2): 159–170. [https://doi.org/10.1016/0006-3207\(94\)00078-5](https://doi.org/10.1016/0006-3207(94)00078-5)
- Bianco P.G. 2014. An update on the status of native and exotic freshwater fishes of Italy. *Journal of Applied Ichthyology* 30: 62–77. <https://doi.org/10.1111/jai.12291>
- Bianco P.G. & De Bonis S. 2015. A taxonomic study of the genus *Phoxinus* (Actinopterygii, Cyprinidae) from Italy and western Balkans with description of four new species: *P. ketmaieri*, *P. karsticus*, *P. apollonicus* and *P. likai*. In: De Filippo G. & Bianco P.G. (eds) *Researches on Wildlife Conservation* 4: 1–17. IGF Publishing, USA.
- Bianco P.G. & Ketmaier V. 2014. A revision of the *Rutilus* complex from Mediterranean Europe with description of a new genus, *Sarmarutilus*, and a new species, *Rutilus stoumboudae* (Teleostei: Cyprinidae). *Zootaxa* 3481 (3): 379–402. <https://doi.org/10.11646/zootaxa.3841.3.4>
- Bianco P.G. & Knežević B. 1987. The *Leuciscus cephalus* complex (Pisces, Cyprinidae) in the western Balkanic area. In: Kullander S.O. & Fernholm B. (eds) *Proceedings of V Congress of European Ichthyologists*: 49–55. Department of Vertebrate Zoology, Swedish Museum of Natural History, Stockholm.
- Bianco P.G. & Kottelat M. 2005. *Scardinius knezevici*, a new species of rudd from Lake Skadar, Montenegro (Teleostei, Cyprinidae). *Ichthyological Exploration of freshwaters* 16 (3): 231–238.
- Bianco P.G. & Taraborelli T. 1985. Contributo alla conoscenza del genere *Rutilus* Rafinescue in Italia e nei Balcani occidentali (Pisces, Cyprinidae). *Bolletino del Museo Regionale di Scienze Naturale di Torino* 3 (1): 131–172.
- Bianco P.G., Aprea G., Balletto E., Capriglione T., Fulgione D. & Odierna G. 2004. The karyology of the cyprinid genera *Scardinius* and *Rutilus* in southern Europe. *Ichthyological Research* 51: 274–278. <https://doi.org/10.1007/s10228-004-0221-y>
- Biberdžić J. 1968. *Hromosomski Brojevi Karakteristični za Vrste Paraphoxinus alepidotus, Paraphoxinus metohiensis i Paraphoxinus pstrossi*. Diploma thesis, University of Sarajevo, Sarajevo.
- Bobori D.C. & Economidis P.S. 2006. Freshwater fish of Greece: Their biodiversity, fisheries and habitats. *Aquatic Ecosystem Health & Management* 9: 407–418. <https://doi.org/10.1080/14634980601027855>
- Bogdanović J. 1954. Ribarski riječnik II. *Ribarstvo Jugoslavije* 2: 34–35.
- Bogdanović J. 1957. Pliva – jedna od naših najljepših i najbogatijih rijeka. *Ribarstvo Jugoslavije* 6: 143–144.

- Bogićević D. 1988. *Morfometrijski i Meristički Karakteri Vrste Chondrostoma nasus (Linnaeus, 1758) iz rijeke Jadar*. Diploma thesis, University of Sarajevo, Sarajevo.
- Bogunić A. 2005. *Ishrana Potočne Pastrmke – Salmo trutta m. fario (Linnaeus 1758) iz Rijeke Vrbas*. Diploma thesis, University of Sarajevo, Sarajevo.
- Bogut I., Novoselić D. & Pavličević I. 2006. *Biologija riba*. Sveučilište J.J. Strossmayer, Fakultet za poljoprivredu, Osijek, Sveučilište u Mostaru, Mostar.
- Bogutskaya N.G. 1994. A description of *Leuciscus lepidus* (Heckel, 1843) with comments on *Leuciscus* and leuciscine-aspinine relationships (Pisces, Cyprinidae). *Annalen des naturhistorischen Museums in Wien, Serie B, Botanik und Zoologie* 96B: 599–620.
- Bogutskaya N.G. & Zupančić P. 1999. A re-description of *Leuciscus zrmanjae* (Karaman 1928) and new data on the taxonomy of *Leuciscus illyricus*, *L. svallize* and *L. cephalus* (Pisces, Cyprinidae) in the West Balkans. *Annalen des naturhistorischen Museums in Wien, Serie B, Botanik und Zoologie* 101B: 509–529.
- Bogutskaya N.G. & Zupančić P. 2003. *Phoxinellus pseudalepidotus*, a new species from the Neretva basin (Teleostei, Cyprinidae) with an overview of the morphology of *Phoxinellus* species of Croatia and Bosnia-Herzegovina. *Ichthyological Exploration of Freshwaters* 14 (4): 369–383.
- Bogutskaya N.G., Zupančić P., Bogut I. & Naseka A. 2012. Two new freshwater fish species of the genus *Telestes* (Actinopterygii, Cyprinidae) from karst poljes in Eastern Herzegovina and Dubrovnik littoral (Bosnia and Herzegovina and Croatia). *ZooKeys* 180: 53–80. <https://doi.org/10.3897/zookeys.180.2127>
- Bogutskaya N.G., Zupančić P., Jelić D., Diripasko O.A. & Naseka A.M. 2017. Description of a new species of *Alburnus* Rafinesque, 1820 (Actinopterygii, Cyprinidae, Leuciscinae) from the Kolpa River in the Sava River system (upper Danube River drainage), with remarks on the geographical distribution of shemayas in the Danube. *ZooKeys* 688: 81–110. <https://doi.org/10.3897/zookeys.688.11261>
- Bogutskaya N.G., Diripasko O.A., Zupančić P., Jelić D. & Naseka A.M. 2019. Phenotypic diversity in an endangered freshwater fish *Squalius microlepis* (Actinopterygii, Leuciscidae). *ZooKeys* 897: 115–147. <https://doi.org/10.3897/zookeys.897.38768>
- Bohlen J. & Ráb P. 2001. Species and hybrid richness in spined loaches of the genus *Cobitis* L. (Teleostei, Cobitidae), with a checklist of European forms and suggestions for their conservation. *Journal of Fish Biology* 59A: 75–89. <https://doi.org/10.1111/j.1095-8649.2001.tb01380.x>
- Bojčić C. 1987. Strategija razvoja šaranskog ribnjačarstva na osnovi većeg udjela sporednih riba u ukupnoj proizvodnji. *Ribarstvo Jugoslavije* 42: 130–140.
- Bojčić C., Debeljak Lj., Vuković T., Jovanović-Kršljanin B., Apostolski K., Ržanićanin B., Turk M., Volk S., Drecun Đ., Habeković D., Hristić Đ., Fijan N., Pažur K., Bunjevac I. & Marošević Đ. 1982. *Slatkovodno ribarstvo*. Poslovna zajednica slatkovodnog ribarstva Jugoslavije, Zagreb
- Bozja A. & Švob M. 1974. Histološka ispitivanja probavnog trakta nekih vrsta ciprinida. *Ichthyologia* 6 (1): 7–13.
- Bralo Z. 1967. *Taksonomski Karakteri Scardinus erythrophthalmus (Linné) iz Livanjskog Polja*. Diploma thesis. University of Sarajevo, Sarajevo.
- Bravničar J. 2012. *Taxonomic Analysis of Bullhead (Cottus sp.) in Slovenia with Application of Molecular Methods*. Bachelor's thesis, Univerza u Ljubljani, Slovenia.
- Bravničar J., Palandačić A., Marić S., Šanda R. & Snoj A. 2015. Phylogeny of European bullhead (*Cottus* sp.) in Western Balkan. In: Correia A. (ed.) *Abstract book of the XV European Congress of Ichthyology*: 35–36. CIIMAR, Porto.

- Bravničar J., Palandačić A., Jelić D., Podgornik S. & Snoj A. 2021. Molecular data reveal distinct phylogenetic position of *Cottus metae*, establish its distribution range, and invalidate the species status of *C. scaturigo*. *Journal of Zoological Systematics and Evolutionary Research* 59 (2): 428441. <https://doi.org/10.1111/jzs.12434>
- Brownstein C.D. & Thomas J.N. 2025. Toward a phylogenetic taxonomy of sturgeons (Acipenseriformes: Acipenseridae). *Bulletin of the Peabody Museum of Natural History* 66 (1): 3–23. <https://doi.org/10.3374/014.066.0101>
- Brusina S. 1892. Pabirci za hrvatsku ihtiologiju i za ribarstvo. *Glasnik Hrvatskoga Naravoslovnoga Društva* 1–6: 221–288.
- Brusina S. 1902. Naše jesetre. *Rad Jugoslavenske akademije znanosti i umjetnosti* 149 (31): 1–69.
- Buj I., Podnar M., Mrakovčić M., Hamzić A., Čaleta M., Marčić Z., Mustafić P., Brigić A., Zanella D. & Skaramuca B. 2007. Raznolikost i taksonomski položaj vijuna roda *Cobitis* u sljevovima Neretve i Trebišnjice. In: Skaramuca B. & Dulčić J. (eds) *Ugrožene i Endemske Vrste Riba u Slivovima Rijeka Neretve, Trebišnjice i Morače. Zbornik Radova Znanstveno-Stručnog Skupa*: 47. East West Institute, Dubrovnik.
- Buj I., Podnar M., Mrakovčić M., Choleva L., Šlechtova V., Tvrtković N., Čaleta M., Mustafić P., Marčić Z., Zanella D. & Brigić A. 2008. Genetic diversity and phylogenetic relationships of spined loaches (genus *Cobitis*) in Croatia based on mtDNA and allozyme analyses. *Folia Zoologica* 57 (1–2): 71–82.
- Buj I., Vukić J., Šanda R., Perea S., Čaleta M., Marčić Z., Bogut I., Povž M. & Mrakovčić M. 2010. Morphological comparison of bleaks (*Alburnus*, Cyprinidae) from the Adriatic Basin with the description of a new species. *Folia Zoologica* 59 (2): 129–141. <https://doi.org/10.25225/fozo.v59.i2.a8.2010>
- Buj I., Šanda R., Marčić Z., Čaleta M. & Mrakovčić M. 2014. Combining morphology and genetics in resolving taxonomy – a systematic revision of spined loaches (genus *Cobitis*; Cypriniformes, Actinopterygii) in the Adriatic watershed. *PLoS ONE* 9 (6): e99833. <https://doi.org/10.1371/journal.pone.0099833>
- Buj I., Čaleta M., Marčić Z., Šanda R., Vukić J. & Mrakovčić M. 2015a. Different histories, different destinies – impact of evolutionary history and population genetic structure on extinction risk of the Adriatic spined loaches (genus *Cobitis*; Cypriniformes, Actinopterygii). *PLoS ONE* 10 (7): e0131580. <https://doi.org/10.1371/journal.pone.0131580>
- Buj I., Šanda R., Marčić Z., Čaleta M. & Mrakovčić M. 2015b. Sexual dimorphism of five *Cobitis* species (Cypriniformes, Actinopterygii) in the Adriatic watershed. *Folia Zoologica* 64: 97–103. <https://doi.org/10.25225/fozo.v64.i2.a2.2015>
- Buj I., Marčić Z., Čaleta M., Šanda R., Geiger M.F., Freyhof J., Machordom A. & Vukić J. 2017. Ancient connections among the European rivers and watersheds revealed from the evolutionary history of the genus *Telestes* (Actinopterygii; Cypriniformes). *PLoS ONE* 12 (12): e0187366. <https://doi.org/10.1371/journal.pone.0187366>
- Buj I., Marčić Z., Čavlović K., Čaleta M., Tutman P., Zanella D., Duplić A., Raguž L., Ivić L., Horvatić S. & Mustafić P. 2020a. Multilocus phylogenetic analysis helps to untangle the taxonomic puzzle of chubs (genus *Squalius*; Cypriniformes, Actinopteri) in the Adriatic basin of Croatia and Bosnia and Herzegovina. *Zoological Journal of the Linnean Society* 189: 953–974. <https://doi.org/10.1093/zoolinnean/zlz133>
- Buj I., Raguž L., Marčić Z., Čaleta M., Duplić A., Zanella D., Mustafić P., Ivić L., Horvatić S. & Karlović R. 2020b. Plitvice Lakes National park harbors ancient, yet endangered diversity of trout (genus *Salmo*). *Journal of Applied Ichthyology* 37: 20–37. <https://doi.org/10.1111/jai.14120>
- Buj I., Knjaz M., Čaleta M., Marčić Z., Ivić L., Onorato L., Šanda R., Vukić J., Horvatić S., Zanella D. & Mustafić P. 2021. Phylogeographic pattern, genetic diversity, and evolutionary history of the enigmatic

- freshwater fish species *Aulopyge huegelii* (Actinopterygii: Cyprinidae). *Journal of Zoological Systematics and Evolutionary Research* 59: 2086–2102. <https://doi.org/10.1111/jzs.12574>
- Buj I., Marčić Z., Flauder E., Šanda R. & Vukić J. 2022. Population genetic structure of endemic fish species facilitating their survival in changing environments – A case study on the genus *Telestes* in Croatia. *Diversity* 14 (7): 529. <https://doi.org/10.3390/d14070529>
- Buntić I. 1985. *Morfometrijski i Meristički Karakteri Vrste Leuciscus svallize (Heckel et Kner, 1858.) i Podvrste Leuciscus ephalus albus iz Bilećkog Jezera*. Diploma thesis. University of Sarajevo, Sarajevo.
- Burek O. 1985. *Dužina Digestivnog Trakta Vrste Alburnus alborella (de Filippi, 1844) iz Hutovog Blata*. Diploma thesis. University of Sarajevo, Sarajevo.
- Calegari B.B., Freyhof J., Waldock C., Wegscheider B., Josi D., Rüber L. & Seehausen O. 2025. Two new species of stone loaches of the genus *Barbatula* (Cypriniformes: Nemacheilidae) from Europe with a neotype designation of *B. barbatula* (Teleostei: Nemacheilidae). *Journal of Fish Biology* 107 (45): 1364–1397. <https://doi.org/10.1111/jfb.70108>
- Cano J.M., Matsuba C., Mäkinen H.S. & Merilä J. 2006. The utility of QTL-linked markers to detect selective sweeps in natural populations – a case study of the EDA gene and a linked marker in threespine stickleback. *Molecular Ecology* 15: 4613–4621. <https://doi.org/10.1111/j.1365-294X.2006.03099.x>
- Cano J.M., Mäkinen H.S., Leinonen T., Freyhof J. & Merilä J. 2008. Extreme neutral genetic and morphological divergence supports classification of Adriatic threespine stickleback (*Gasterosteus aculeatus*) populations as distinct conservation units. *Biological Conservation* 141: 1055–1066. <https://doi.org/10.1016/j.biocon.2008.01.015>
- Carosi A., Lorenzoni F., Oneto F., Capurro M., Ovčina J., Rezzoagli D., Petroselli C., Selvaggi R., Cappelletti D., Sanz N. & Lorenzoni M. 2024. The role of protected areas in the Balkan freshwater biodiversity conservation: The case of Blidinje Nature Park (Southwestern Bosnia–Herzegovina). *Journal for Nature Conservation* 82: 126739. <https://doi.org/10.1016/j.jnc.2024.126739>
- Cerović D. & Ivanović B. 1968. Some morphological characteristics of *Salmo trutta m. fario* from River Čehotina. *Poljoprivreda i Šumarstvo* 3: 17–24.
- Changeux T. & Pont D. 1995. Current status of the riverine fishes of the French Mediterranean basin. *Biological conservation* 72 (2): 137–158.
- Charnpila E.A., Šanda R., Doadrio I., Perea S., Marić D., Radošević D. & Vukić J. 2025. Endemic Neretva bleak, *Alburnus neretvae* (Cypriniformes: Leuciscidae) from fragile karstic environment: First assessment of the population genetic diversity. *Biologia* 80: 2761–2770. <https://doi.org/10.1007/s11756-025-02002-6>
- Cicović V. 1964. *Prilog Ispitivanju Rasta Pastrmki (Salmo trutta m. fario) iz Potoka Oteše*. Diploma thesis. University of Sarajevo, Sarajevo.
- Clavero M. & García-Berthou E. 2005. Invasive species area leading cause of animal extinctions. *Trends in Ecology and Evolution* 20: 110. <https://doi.org/10.1016/j.tree.2005.01.003>
- Coll-Costa C., Dahms C., Kempainen P., Alexandre C.M., Ribeiro F., Zanella D., Zanella L., Merilä J. & Momigliano P. 2024. Parallel evolution despite low genetic diversity in three-spined sticklebacks. *Proceedings of the Royal Society B* 291 (2020): e20232617. <https://doi.org/10.1098/rspb.2023.2617>
- Costa M., Duarte G., Segurado P. & Branco P. 2021. Major threats to European freshwater fish species. *Science of The Total Environment* 797: e149105. <https://doi.org/10.1016/j.scitotenv.2021.149105>
- Crivelli A.J. 1996. *The Freshwater Fish Endemic to the Northern Mediterranean Region. An Action Plan for their Conservation*. Tour du Valat Publication, Arles.

- Cvijić S., Golub D., Šukalo G., Dekić R., Kostić D., Miljanović B. & Kanlić V. 2020. Ihtiofauna donjeg dijela toka rijeke Vrbanje (Republika Srpska, BiH). *Acta Scientifica Balcanica* 1 (1): 49–60. <https://doi.org/10.7251/skp201101049c>
- Čanković M. 1964. Parazitofauna riba u Bosni i Hercegovini (digeni trematodi II). *Veterinaria* 13 (3): 307–312.
- Čanković M. 1967. Digeni trematodi salmonidnih i nekih drugih riba u Bosni i Hercegovini. *Veterinaria* 16 (3): 381–391.
- Čanković M., Delić S., Kiškarolj M. & Rukavina J. 1968a. *Parazitofauna Slatkovodnih Riba Bosne i Hercegovine (Trematoda, Cestoidea, Nematoda, Acanthocephala)*. Akademija nauka i umjetnosti Bosne i Hercegovine, Sarajevo.
- Čanković M., Kiškarolj M. & Ramadanović R. 1968b. Helmintofauna riba jednog dijela srednjeg toka rijeke Save. I. Digeni trematodi i cestodi. *Veterinaria* 17 (1): 17–28.
- Čanković M., Kiškarolj M. & Ramadanović R. 1968c. Helmintofauna riba jednog dijela srednjeg toka rijeke Save. II. Nematodi i akantocefali. *Veterinaria* 17 (2): 151–158.
- Čanković M., Kiškarolj M. & Kosorić Đ. 1970. Ispitivanje sezonske dinamike nekih vrsta endohelminata neretvanske mekousne (*Salmothymus obtusirostris oxhyrhynchus* Steind.) iz rijeke Bune. *Ichthyologia* 2 (1): 31–36.
- Čanković M., Kiškarolj M. & Kosorić Đ. 1971. Ispitivanja invadiranosti nekih podvrsta *Salmothymus obtusirostris* Heck. endohelmintima u nekim rijekama jadranskog sliva. *Ichthyologia* 3 (1): 13–18.
- Čanković M., Žitnan R. & Građanin S. 1976. Paraziti iz klase Monogenoidea i njihova epizootiološka važnost kod šarana na jednom ribnjačarstvu u Bosni i Hercegovini. *Veterinaria* 25 (1–2): 45–54.
- Čičić-Močić A. 2014. Hidromorfološke promjene vodotoka i njihov uticaj na ihtiofaunu. *Voda i Mi* 85: 27–34.
- Čičić-Močić A. 2016. Ihtiološka istraživanja sliva rijeke Save u Federaciji Bosne i Hercegovine u 2015. godini. *Voda i Mi* 92: 10–22.
- Čičić-Močić A. 2018. Ihtiološka istraživanja sliva rijeke Save u Federaciji Bosne i Hercegovine u 2017. godini. *Voda i Mi* 98: 34–43.
- Čičić-Močić A. 2019. Ihtiološka istraživanja sliva rijeke Save u Federaciji Bosne i Hercegovine u 2018. godini. *Voda i Mi* 101: 18–34.
- Čolić S. 2018. *Genetic Characterization of the European Mudminnow (Umbra krameri Walbaum, 1792) Population from the Matura River Drainage, Bosnia and Herzegovina*. Master thesis, University of Belgrade, Serbia.
- Čolić S., Šukalo G., Čolić V. & Kerkez V. 2018. First record of round goby *Neogobius melanostomus* Pallas, 1814 (Pisces: Gobiidae) in Bosnia and Herzegovina. *Ecologica Montenegrina* 16: 108–110. <https://doi.org/10.37828/em.2018.16.8>
- Čorbo A. 2013. *Ishrana Potočne Pastrmke Salmo trutta L. u Slivu Gornjeg Toka Rijeke Neretve*. Diploma thesis. University of Sarajevo, Sarajevo.
- Čaleta M., Mrakovčić M., Buj I., Mustafić P., Zanella D. & Marčić Z. 2009a. Threatened fishes of the world: *Aulopyge huegelli* Heckel, 1842 (Cyprinidae). *Environmental Biology of Fishes* 85: 229–230. <https://doi.org/10.1007/s10641-009-9445-z>

- Ćaleta M., Mrakovčić M., Buj I., Mustafić P., Zanella D. & Marčić Z. 2009b. Threatened fishes of the world, *Chondrostoma phoxinus* Heckel, 1843 (Cyprinidae). *Environmental Biology of Fishes* 84: 229–230. <https://doi.org/10.1007/s10641-008-9413-z>
- Ćaleta M., Jelić D., Buj I., Zanella D., Marčić Z., Mustafić P. & Mrakovčić M. 2010. First record of the non-native invasive species rotan (*Percottus glenii* Dybowski, 1877) in Croatia. *Journal of Applied Ichthyology* 27: 146–147. <https://doi.org/10.1111/j.1439-0426.2010.01612.x>
- Ćaleta M., Buj I., Mrakovčić M., Mustafić P., Zanella D., Marčić Z., Duplić A., Mihinjač T. & Katavić I. 2015. *Hrvatske Endemske Ribe*. Agencija za zaštitu okoliša, Zagreb.
- Ćaleta M., Marčić Z., Buj I., Zanella D., Mustafić P., Duplić A. & Horvatić S. 2019. A review of extant Croatian freshwater fish and lampreys. *Croatian Journal of Fisheries* 77 (3): 137–234. <https://doi.org/10.2478/cjf-2019-0016>
- Ćemalović Z. 1975. *Morfometrijski i Meristički Karakteri Vrste Gasterosteus aculeatus Linne Rijeke Bune*. Master's thesis, University of Sarajevo, Sarajevo.
- Ćemer A. 2010. *Morfoanatomske Karakteristike Digestivnog Trakta Deverike Abramis brama (Linnaeus, 1758) i Crnooke Deverike Abramis sapa (pallas, 1811) iz Rijeke Save*. Diploma thesis, University of Sarajevo, Sarajevo.
- Ćulibrk B., Bilbija B., Nedić Z. & Nikolić V. 2016. Ectoparasites of fish from Lake Gradina, Bosnia and Herzegovina. *Veterinarski Žurnal Republike Srpske* 16 (1): 11–19. <https://doi.org/10.7251/VETJ1601011C>
- Ćurčić M. 2017. Minders, Brunette (*Umbra krameri* Walbaum, 1792) endangered species in the swamp Gromiželj. *Nova Škola* 12 (1): 14–18. <https://doi.org/10.7251/NSK1701014C>
- Ćurčić V. 1910. Narodno ribarstvo Bosni i Hercegovini. *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 2–3: 379–488.
- Ćurčić V. 1913. Narodno ribarstvo u Bosni i Hercegovini. Glasnik II. Hercegovina. *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 25: 421–514.
- Ćurčić V. 1915a. Narodno ribarstvo u Bosni i Hercegovini. Glasnik II. Hercegovina. Nastavak. *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 27: 37–107.
- Ćurčić V. 1915b. Narodno ribarstvo u Bosni i Hercegovini. Glasnik II. Hercegovina. Svršetak. *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 27: 313–350.
- Ćurčić V. 1916. Narodno ribarstvo u Bosni i Hercegovini. Glasnik III. Zapadnobosanski krš, i to ispostava Kupres i kotarevi Duvno, Glamoč, Livno i Blidinjsko jezero. *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 28: 397–426.
- Ćurčić V. 1933. Neprijatelji riba. *Ribarski List* 5–6: 62–64.
- Ćurčić V. 1938. *Neretva i Njezine Pastrve (Salmonidae)*. Štamparija Risto V. Savić, Sarajevo.
- Darwall W., Carrizo S., Numa C., Barrios V., Freyhof J. & Smith K. 2014. *Freshwater Key Biodiversity Areas in the Mediterranean Basin Hotspot: Informing Species Conservation and Development Planning in Freshwater Ecosystems*. IUCN, Cambridge, United Kingdom and Malaga, Spain. <https://doi.org/10.2305/IUCN.CH.2014.SSC-OP.52.en>
- Dedić S. 1971. *Neka Ekološka Posmatranja na Lipljenu Rijeke Une*. Diploma thesis. University of Sarajevo, Sarajevo.
- DeFaveri J., Zanella L.N., Zanella D., Mrakovčić M. & Merilä J. 2012. Phylogeography of isolated freshwater three-spined stickleback (*Gasterosteus aculeatus*) populations in the Adriatic basin. *Journal of Fish Biology* 80: 62–85. <https://doi.org/10.1111/j.1095-8649.2011.03147.x>

- Dekić R., Ivanc A., Bakrač-Bećiraj A., Bošković J. & Lolić S. 2009a. Hematological parameters of fishes as the indicators of environment. *In: Marković Z., Poleksić V., Živić I., Stanković M. & Dulić Z. (eds) 4th International Conference Fishery. Conference Proceedings: 204–210.* Faculty of Agriculture, Belgrade-Zemun.
- Dekić R., Ivanc A., Bakrač-Bećiraj A., Bošković J., Lolić S. & Vuković D. 2009b. Normal Hematological Values of Cultured Grayling. *In: Marković Z., Poleksić V., Živić I., Stanković M. & Dulić Z. (eds) 4th International Conference Fishery. Conference Proceedings: 358–364.* Faculty of Agriculture, Belgrade-Zemun.
- Dekić R., Ivanc A., Lolić S., Bošković J., Obradović S. & Četković D. 2011. The recent state of distribution of endemic fish species in eastern Herzegovina. *In: Marković Z. (ed.) 5th International Conference Aquaculture and Fishery. Conference Proceedings: 195–199.* University of Banja Luka, Belgrade-Zemun.
- Dekić R., Ivanc A., Lolić S., Manojlović M., Janjić N., Erić Ž. & Četković D. 2012a. Red blood cell parameters of *Telestes metohiensis* from Pribitul watercours. *Skup* 4 (2): 153–160.
- Dekić R., Ivanc A., Lukač M. & Krnić J. 2012b. Morphometric characteristics of red blood cells of *Telestes metohiensis* (Steindacher, 1901). *Veterinaria* 61 (3–4): 115–127.
- Dekić R., Ivanc A. & Erić Ž. 2013a. Distribution of endemic fish species in Eastern Herzegovina. *In: Đorđević D. (ed.) 3rd International Simposium of Natural Resources Management, Proceedings: 127–130.* Faculty of management, Zaječar.
- Dekić R., Ivanc I., Erić Ž., Lolić S., Manojlović M. & Janjić N. 2013b. Morphometric characteristics of *Telestes metohiensis* from different watercourses of the Dabar Field. *Agro-knowledge Journal* 14 (4): 567–573. <https://doi.org/10.7251/agrsr1304567d>
- Dekić R., Ivanc A., Gnjata R., Trbić G., Četković D. & Lolić S. 2013c. Effect of thermal stress of short duration on the red blood cell parameters of *Barbus balcanicus* Kotlik, Tsigenopulos, Rab & Berrebi, 2002. *African Journal of Biotechnology* 12 (18): 2484–2491.
- Dekić R., Ivanc A., Mandić M., Lolić S. & Bakrač-Bećiraj A. 2013d. The effect of temperature change on erythrocyte profile of *Carassius gibelio*. *Croatian Journal of Fisheries* 71 (1): 32–36.
- Dekić R., Ivanc A., Erić Ž., Gnjata R., Trbić G., Lolić S., Manojlović M. & Janjić N. 2014. Hematological characteristics of *Delminichthys ghetaldii* (Steindachner 1882) inhabiting the karst region of eastern Herzegovina. *Archives of Biological Sciences* 66 (4): 1423–1430. <https://doi.org/10.2298/ABS1404423D>
- Dekić R., Bilbija B., Lukač M., Mandić M., Friščić J., Ivanc A. & Bećiraj A. 2016a. Morphometric characteristics of Dalmatian barbel-gudgeon (*Aulopyge huegelii*) and Adriatic minnow (*Phoxinellus alepidotus*) from the Šator Lake. *Skup* 7 (2): 123–128. <https://doi.org/10.7251/SKP1607123D>
- Dekić R., Manojlović M., Friščić J., Lolić S. & Golub D. 2016b. Morphometric and Physiological Characteristics of Brown Trout (*Salmo trutta*) from the Ponor River. *Agro-knowledge Journal* 17 (3): 233–242. <https://doi.org/10.7251/AGREN1603233D>
- Dekić R., Friščić J., Ivanc A. & Kukavica B. 2016c. Characterization of proteins from Popovo Minnow (*Delminichthys ghetaldii* Steindachner, 1882) muscle. *Turkish Journal of Fisheries and Aquatic Sciences* 16: 637–642. https://doi.org/10.4194/1303-2712-v16_3_17
- Dekić R., Friščić J., Lolić S., Lubarda B. & Manojlović M. 2017a. Morphometric characteristics of *Delminichthys ghetaldii* (Steindachner, 1882) from different habitats. *In: Pešić V. & Hadžiablahović S. (eds) 7th International Symposium of Ecologists (ISEM7). Proceedings: 82–85.* Institute of Biodiversity and Ecology, Sutomore.

- Dekić R., Friščić J., Manojlović M., Lolić S. & Golub D. 2017b. Ichthyofaunal diversity of the Drinjača catchment area. *In: Pešić V. & Hadžiablahović S. (eds) 7th International Symposium of (ISEM7). Proceedings: 77–81.* Institute of Biodiversity and Ecology, Sutomore.
- Dekić R., Golub D., Lolić S., Manojlović M. & Paspalj J. 2020a. Physical, chemical and biological parameters as water quality of River Sana (Republic of Srpska, B&H). *In: Đukić A. (ed.) 50th Annual Conference of the Serbian Water Pollution Control Society Water 2020. Conference Proceedings: 195–201.* Serbian Water Pollution Control Society, Trebinje.
- Dekić R., Ivanc A., Četković D. & Lolić S. 2020b. Anthropogenic impact and environmental quality of different tributaries of the River Vrbas (Bosnia and Herzegovina). *In: Bănăduc D., Curtean-Bănăduc A., Pedrotti F., Cianfaglione K. & Akeroyd J.R. (eds) Human Impact on Danube Watershed Biodiversity in the XXI Century. Geobotany Studies: 169–213.* Springer International Publishing, Cham. <https://doi.org/10.1007/978-3-030-37242-2>
- Dekić R., Golub D., Kalamujić Stroil B., Bakrač A., Avdić S., Kanlić V., Škrijelj R. & Adrović A. 2024. Fish Resources and Fisheries in Bosnia and Herzegovina: Overview, Sustainability, and Conservation. *In: Simić V., Simić S. & Pešić V. (eds) Ecological Sustainability of Fish Resources of Inland Waters of the Western Balkans. Freshwater Fish Stocks, Sustainable Use and Conservation: 73–111.* Fish & Fisheries Series 43, Springer Nature, Switzerland. <https://doi.org/10.1007/978-3-031-36926-1>
- Delić A. 1984. Zaštita faune riba u otvorenim vodama novogradiške općine. *Ribarstvo Jugoslavije* 3–4: 52–58.
- Delić A. 1985. Dužinski i težinski rast riba u otvorenim vodama novogradiške općine. *Ribarstvo Jugoslavije* 4–6: 80–85.
- Delić A., Bučar M., Kučinić M. & Mrakovčić M. 2003. New data about the distribution of *Sabanejewia balcanica* (Karaman, 1922). (Cobitidae) in Croatia. *Folia Biologica (Kraków)* 51 (Suppl.): 39–42.
- Delić A., Kučinić M., Marić D. & Bučar M. 2005. New data about the distribution of *Phoxinellus alepidotus* (Heckel, 1843) and *Aulopyge huegelii*. *Natura Croatica* 14 (4): 351–355.
- Delić A., Šanda R., Bučar M., Mihoci I., Vilenica M., Vukić J., Lelo S. & Kučinić M. 2014. New data on distribution of the Monkey goby, *Neogobius fluviatilis* (Pallas, 1814) in Bosnia and Herzegovina and Croatia with notes on ecology and associated fish fauna. *Natura Croatica* 23 (2): 298–302.
- Dorofejeva E.A.D. & Seratlić D. 1970. Osteološke karakteristike *Salmo marmoratus* Cuvier. *In: III Simpozijum Jugoslavenskog Ihtiološkog Društva, Rezime Saopštenja: 14–15.* Yugoslav Ichthyological Society, Kotor.
- Dorofejeva E.A.D. & Seratlić-Savić D. 1972. The systematic status of *Salmo marmoratus* Cuv. with respect to its osteological characteristics. *Zoologicheskaya Zhurnal* 51 (5): 759–763. [In Russian.]
- Dreca A. 2009. *Tempo Rasta Predkonzumne Kalifornijske Pastrmke Oncorhynchus Mykiss (Walbaum, 1792) na Ribogojilištu Žunovnica Hadžići.* Diploma thesis. University of Sarajevo, Sarajevo.
- Drešković N., Đug S., Stupar V., Hamzić A., Lelo S., Muratović E., Lukić-Bilela L., Brujić J., Milanović Đ. & Kotrošan D. 2011. *NATURA 2000 u Bosni i Hercegovini*. 1. izd. Centar za okolišno održivi razvoj, Sarajevo.
- Dulčić J. & Kovačić M. 2020. *Ichthyofauna of the Adriatic Sea.* Institute for Oceanography and Fisheries, Split.
- Dulčić J., Glamuzina B. & Tutman P. 2005. The first record of the ruffe, *Gymnocephalus cernuus* (Percidae), in the Hutovo Blato wetland, Adriatic drainage system of Bosnia and Herzegovina. *Cybium* 29 (2): 205–206.
- Dulčić J., Tutman P., Matic-Skoko S., Kraljević M., Jug-Dujaković J., Glavić N., Kožul V., Glamuzina B., Bartulović V. & Skaramuca B. 2007. A list of Y-O-Y fish species found in the littoral shallows of the Neretva and Mala Neretva estuaries (Eastern Adriatic, Croatian coast). *Acta Adriatica* 48 (1): 89–94.

- Dulčić J., Tutman P., Glamuzina B., Skaramuca B. & Ahnelt H. 2008. Endemic gobies (Gobiidae) of the Hutovo blato wetland (Neretva River basin, Bosnia and Herzegovina) and their conservation status. In: Skaramuca B. & Dulčić J. (eds) *Zbornik Radova Znanstveno-stručnog Skupa Ugrožene i Endemske Vrste Riba u Slivovima Rijeka Neretve, Trebišnjice i Morače*: 61–66. Sveučilište u Dubrovniku, The EastWest Institute, Dubrovnik.
- Dulčić J., Tutman P., Prusina I., Tomšić S., Dragičević B., Hasković E. & Glamuzina B. 2009. Length-weight relationships for six endemic freshwater fishes from Hutovo blato wetland (Bosnia and Herzegovina). *Journal of Applied Ichthyology* 25 (4): 499–500. <https://doi.org/10.1111/j.1439-0426.2010.01593.x>
- Dumbović V., Posavec Vukelić V., Duplić A., Katušić L., Jelić D., Boršić I. & Partl A. 2009. *Study of the inventarisation of the flora and fauna of the Una River and its riparian belt*. Sisačko-moslavačka županija, Sisak.
- Duran S. 1975. *Morfološko-taksonomske Odlike Gobio gobio L. 1758 iz Vodene Akumulacije Peručac*. Diploma thesis. University of Sarajevo, Sarajevo.
- Durmić-Pašić A. 2008. *Filogenetske i Interpopulacijske Relacije Osobnosti Mitohondrijalne DNK Bh. salmonida*. PhD thesis, Prirodno-matematički fakultet Univerziteta u Sarajevu, Bosna i Hercegovina.
- Durmić-Pašić A., Pojskić N., Kalamujić B. & Hadžiselimović R. 2005. Genetic characterization of fish populations and its use in landscape conservation in Bosnia and Herzegovina. In: Ruane J. & Sonnino A. (eds) *International Workshop The Role of Biotechnology for the Characterization and Conservation of Crop, Forestry, Animal and Fishery Genetic Resources*: 171–172. FAO, Turin.
- Durmić-Pašić A., Kalamujić B., Pojskić N. & Hadžiselimović R. 2007a. Simultaneous amplification of six microsatellite loci in three salmonid species. In: Buj I., Zanella L., & Mrakovčić M. (eds) *12th European Congress of Ichthyology ECI XII. Book of Abstracts*: 26. Croatian Ichthyological Society, Cavtat (Dubrovnik).
- Durmić-Pašić A., Pojskić N., Kalamujić B. & Hadžiselimović R. 2007b. Genetic subdivision of endemic species *Salmothymus obtusirostris oxyrhynchus* Heckel population in Neretva river. In: Buj I., Zanella L., & Mrakovčić M. (eds) *12th European Congress of Ichthyology ECI XII. Book of Abstracts*: 9–14. Croatian Ichthyological Society, Cavtat (Dubrovnik).
- Đonko V. 2005. *Morfološke Osobnosti Digestivnog Trakta Deverike Abramis brama (Linnaeus, 1758) iz rijeke Save*. Diploma thesis. University of Sarajevo, Sarajevo.
- Đurđević S., Dekić R., Vuković D. & Ivanc A. 2008. Kvalitet vode i morfološke, merističke i hematološke karakteristike klena (*Leuciscus cephalus*). *Skup 2* : 231–244.
- Đurović E. 1979. Variation of the relative length of the intestinal tract in some populations of *Alburnoides bipunctatus* (Bloch, 1782) in Yugoslavia. *Glasnik Zemaljskog Muzeja Bosne i Hercegovine u Sarajevu* 24: 243–250.
- Đurović E. & Vuković T. 1975. Experimental hybridization of *Aulopyge hügelii* Heckel, 1843 with the species of *Leuciscus cephalus* (Linnaeus, 1758) *Rutilus rutilus* (Linnaeus, 1758) and *Alburnoides bipunctatus* (Bloch, 1782) (Pisces Cyprinidae). *Godišnjak Biološkog nstituta Univerziteta Sarajevo* 28: 83–92.
- Džafić S., Bakrač A., Suljević D. & Dekić R. 2018. Effects of hyperthermia on erythrocyte parameters of carp *Cyprinus carpio* (Linnaeus, 1758) from Bardača swamp, Bosnia and Herzegovina. *Asian Journal of Fisheries and Aquatic Research* 2 (4): 1–7. <https://doi.org/10.9734/ajfar/2018/v2i430015>
- Economidis P.S. & Bănărescu P.M. 1991. The distribution and origins of freshwater fishes in the Balkan peninsula, especially in Greece. *International Revue of Hydrobiology* 76 (2): 257–283.

- Elvira B. 1987. Taxonomic revision of the genus *Chondrostoma* Agassiz, 1835 (Pisces, Cyprinidae). *Cybiurn* 1: 111–140.
- Elvira B. 1997. Taxonomy of the genus *Chondrostoma* (Osteichthyes, Cyprinidae), an updated review. *Folia Zoologica* 46 (Suppl. 1): 1–14.
- Esteve M., McLennan D.A., Zablocki J.A., Pustovrh G. & Doadrio I. 2014. Spawning behaviour and the softmouth trout dilemma. *Archives of Polish Fisheries* 22: 159–165. <https://doi.org/10.2478/aopf-2014-0016>
- Fedorčák J., Šanda R., Stefanov T., Mendel J., Nowak M., Križek P., Perdices A., Vukić J. & Koščo J. 2023. The „True Colours” of Golden Loaches (Teleostei: Cobitidae). *Fishes* 8 (2): 119. <https://doi.org/10.3390/fishes8020119>
- Filipović Marijić V. & Raspor B. 2014a. Relevance of biotic parameters in the assessment of the spatial distribution of gastrointestinal metal and protein levels during spawning period of European chub (*Squalius cephalus* L.). *Environmental Science and Pollution Research* 21 (12): 7596–7606. <https://doi.org/10.1007/s11356-014-2666-5>
- Filipović Marijić V., Vardić Smrzlić I. & Raspor B. 2014b. Does fish reproduction and metabolic activity influence metal levels in fish intestinal parasites, acanthocephalans, during fish spawning and post-spawning period? *Chemosphere* 112: 449–455. <https://doi.org/10.1016/j.chemosphere.2014.04.086>
- Francuski Lj., Ludoški J., Lukač M., Dekić R. & Milankov V. 2019. Integrative study of population structure of *Telestes dabar*, the strictly endemic cyprinid species from the Dinaric karst on the Balkan Peninsula. *European Journal of Wildlife Research* 65: 66. <https://doi.org/10.1007/s10344-019-1302-6>
- Freyhof J. 1997. Zur Fortpflanzungsbiologie des Barbengründlings. *DATZ Aquarien Terrarien* 50: 513–516.
- Freyhof J. 2012. *Threatened Freshwater Fishes and Molluscs of the Balkan, Potential Impacts of Hydropower Projects*. Unpublished report, ECA Watch Austria & EuroNatur.
- Freyhof J. 2024. *Romanogobio carpathorossicus*. The IUCN Red List of Threatened Species 2024: e.T146233861A146234301. <https://doi.org/10.2305/IUCN.UK.2024-2.RLTS.T146233861A146234301.en>. [accessed on 04 Jul 2025]
- Freyhof J. & Brooks E. 2011. *European Red List of Freshwater Fishes*. Publications Office of the European Communities, Luxembourg. <https://doi.org/10.2779/85903>
- Freyhof J. & Stelbrink B. 2007. *Cobitis illyrica*, a new species of loach from Croatia (Teleostei: Cobitidae). *Ichthyological Exploration of Freshwaters* 18: 269–275.
- Freyhof J., Lieckfeldt D., Pitra C. & Ludwig A. 2005. Molecules and morphology, Evidence for introgression of mitochondrial DNA in Dalmatian cyprinids. *Molecular Phylogenetics and Evolution* 37 (2): 347–354. <https://doi.org/10.1016/j.ympev.2005.07.018>
- Freyhof J., Lieckfeldt D., Bogutskaya N.G., Pitra C. & Ludwig A. 2006. Phylogenetic position of the Dalmatian genus *Phoxinellus* and description of the newly proposed genus *Delminichthys* (Teleostei, Cyprinidae). *Molecular Phylogenetics and Evolution* 38: 416–425. <https://doi.org/10.1016/j.ympev.2005.07.024>
- Freyhof J., Weiss S., Adrović A., Čaleta M., Duplić A., Hrašovec B., Kalamujić B., Marčić Z., Milošević D., Mrakovčić M., Mrdak D., Piria M., Schwarz U., Simonović P., Šljuka S., Tomljanović, T. & Zabric D. 2015. *The Huchen Hucho hucho in the Balkan region: Distribution and Future Impacts by Hydropower Development*. RiverWatch & EuroNatur, Vienna.

- Freyhof J., Bergner L. & Ford M. 2020. *Threatened Freshwater Fishes of the Mediterranean Basin Biodiversity Hotspot: Distribution, Extinction Risk and the Impact of Hydropower*. EuroNatur and RiverWatch, Vienna.
- Fricke R., Eschmeyer W.N. & van der Laan R. (eds) 2024. *Eschmeyer's Catalog of Fishes: Genera, Species, References*. Available from <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> [accessed 3 Jul. 2024].
- Friedrich T., Wiesner C., Zangl L., Daill D., Freyhof J. & Koblmüller S. 2018. *Romanogobio skywalkeri*, a new gudgeon (Teleostei: Gobionidae) from the upper Mur River, Austria. *Zootaxa* 4403 (2): 336–350. <https://doi.org/10.11646/zootaxa.4403.2.6>
- Gajević M. 2005. *Morfološko-taksonomske Karakteristike Crvenookice - Rutilus rutilus (Linnaeus, 1758) iz Rijeke Save*. Diploma thesis. University of Sarajevo, Sarajevo.
- Gajević M. & Korjenić E. 2005. Morfološko-taksonomske karakteristike crvenookice *Rutilus rutilus* (Linnaeus, 1758) iz rijeke Save. *Veterinaria* 54 (3–4): 211–224.
- Galić P. 1986. *Dužina Digestivnog Trakta Vrste Rutilus rubilio (Bonaparte, 1837) iz Hutovog Blata*. Diploma thesis. University of Sarajevo, Sarajevo.
- Geiger M.F., Herder F., Monaghan M.T., Almada V., Barbieri R., Bariche M., Berrebi P., Bohlen J., Casal-Lopez M., Delmastro G.B., Denys G.P.J., Dettai A., Doadrio I., Kalogianni E., Kärst H., Kottelat M., Kovačić M., Laporte M., Lorenzoni M., Marčić Z... & Freyhof J. 2014. Spatial heterogeneity in the Mediterranean biodiversity hotspot affects barcoding accuracy of its freshwater fishes. *Molecular Ecology Resources* 14 (6): 1210–1221. <https://doi.org/10.1111/1755-0998.12257>
- Georgiev S. 2003. On the origin of the Balkan peninsula salmonids. *Ribarstvo* 4: 147–174.
- Georgiev S. 2011. Study of *Salmo* genus (Pisces, Salmonidae) on the Balkan Peninsula. *Ribarstvo* 69 (1): 11–20.
- Gilić N. 1983. *Morfometrijski i Meristički Karakteri Vrste Leuciscus turskyi iz Buškog Blata*. Diploma thesis. University of Sarajevo, Sarajevo.
- Gilles A., Costedoat C., Barascud B., Voisin A., Banarescu P., Bianco P.G., Economidis P.S., Marić D. & Chappaz R. 2009. Speciation pattern of *Telestes souffia* complex (Teleostei, Cyprinidae) in Europe using morphological and molecular markers. *Zoologica Scripta* 39: 225–242.
- Glamuzina B. & Bartulović V. 2006. Some characteristics of the endemic dentex trout, *Salmo dentex* (Heckel, 1851) from the Neretva River, Bosnia-Herzegovina. *Ribarstvo* 64 (2): 59–64.
- Glamuzina L. & Dobrosravić T. 2020. Summer fish migrations in the River Neretva (South-Eastern Adriatic coast, Croatia) as a consequence of salinization. *Naše More* 67: 103–116. <https://doi.org/10.17818/NM/2020/2.3>
- Glamuzina B., Tutman P. & Conides A. 2001. *Report on Ichthyological Survey on Hutovo Blato Wetlands. European Union LIFE Third Countries Program and the Ministry of Civil Engineering and Nature Protection of Neretva-Herzegovina Cantonal Government/LTCY/035/BiH project, "Development of a New Management Policy for Hutovo Blato Wetlands, Bosnia-Herzegovina"*. Institut za oceanografiju i ribarstvo, Split, Dubrovnik.
- Glamuzina B., Tutman P., Buntić I., Bartulović V. & Conides A. 2002. Kvalitativni sastav ihtiofaune Hutova Blata. In: Buntić I. (ed.) *Nova Politika Gospodarenja Vlažnim Područjima Hutova Blata. Završni Simpozij LIFE Projekta LIFECY 99/BIH/035*: 38–44. LIFE i Ministarstvo graditeljstva, prostornog uređenja i zaštite okoliša HNŽ-K, Mostar.

- Glamuzina B., Bartulović V., Dulčić J., Conides A.J., Tutman P., Matić-Skoko S., Gavrilović A., Jug-Dujaković J., Hasković E., Ivanc A. & Zovko N. 2007. Some biological characteristics of the endemic Neretvan nase, *Chondrostoma knerii* Heckel, 1843, in the Hutovo Blato wetlands (Bosnia and Herzegovina). *Journal of Applied Ichthyology* 23 (3): 221–225. <https://doi.org/10.1111/j.1439-0426.2006.00828.x>
- Glamuzina B., Dulčić J., Hasković E., Ivanc A., Mandić S., Mrdak D. & Skaramuca B. 2008a. Stanje ihtiofaune u slivovima rijeka Neretve, Trebišnjice i Morače tijekom ljetnih mjeseci 2007. In: Skaramuca B. & Dulčić J. (eds) *Zbornik Radova Znanstveno-stručnog Skupa "Ugrožene i Endemske Vrste Riba u Slivovima Rijeka Neretve, Trebišnjice i Morače"*: 21–41. Sveučilište u Dubrovniku, EastWest Institute, Dubrovnik.
- Glamuzina B., Bartulović V., Conides A. & Zovko N. 2008b. Status populacije europske jegulje, *Anguilla anguilla* (Linnaeus, 1758) na području močvare Hutovo blato, Bosna i Hercegovina. In: Pospišil M. (ed.) *Proceedings of 43rd Croatian and 3rd International Symposium on Agriculture: 733–736*. Agronomski fakultet, Zagreb.
- Glamuzina B., Dulčić J., Tutman P., Matić-Skoko S., Tomšić S., Bartulović V. 2009. Record size of *Salmo dentex* (Heckel, 1851) (Salmonidae) from Neretva River (Bosnia and Herzegovina). *Cybium* 33 (2): 165–167. <https://doi.org/10.26028/CYBIUM/2009-332-008>
- Glamuzina B., Pavličević J., Tutman P., Glamuzina L., Bogut I. & Dulčić J. 2013. *Ribe Neretve*. Udruga Centar za zaštitu i promicanje endemičnih i autohtonih ribljih vrsta, CEAV, Modro zelena zadruga braniteljia Metković, Metković.
- Glamuzina B., Tutman P., Nikolić V., Vidović Z., Pavličević J., Vilizzi L., Copp G.H. & Simonović P. 2017. Comparison of taxon-specific and taxon-generic risk screening tools to identify potentially invasive non-native fishes in the river Neretva catchment (Bosnia and Herzegovina and Croatia). *River Research and Applications* 33 (5): 670–679. <https://doi.org/10.1002/rra.3124>
- Glamuzina B., Stanić-Koštroman S., Matić-Skoko S., Glamuzina L., Muhamedagić S., Rozić I., Weiss S. & Pavličević J. 2018. Recent status and life history traits of endangered soft-mouth trout, *Salmo obtusirostris* in the Neretva River catchment (Bosnia and Herzegovina) as a consequence of river alteration. *Journal of Applied Ichthyology* 34 (5): 1160–1168. <https://doi.org/10.1111/jai.13780>
- Glamuzina B., Dobroslačić T., Bukvić V., Tutman P. & Bartulović V. 2019. Rare records of mature Sea lamprey, *Petromyzon marinus* Linnaeus, 1758 during migration in the River Neretva (Croatia, Bosnia and Herzegovina). *Cahiers de Biologie Marine* 60: 399–402. <https://doi.org/10.21411/CBM.A.D3EF0B92>
- Glowacki J. 1896. Die Fischfauna der Save und der Isonzo. I. *Jahresbericht des kaiserlich-königlichen Staats-Untergymnasiums in Chilli 1895–1896*: 1–37.
- Golob A., Skoberne P., Milanović Đ., Drešković N., Đug S., Kovačević D., Radošević D., Todorović S., Brujić J., Stupar V., Gašić B., Stanivuković Z., Burlica Č., Pašić J., Petković D., Bokić M., Nikić D., Zafirov I., Velghe D., Ferlin F. & Andries T. 2015. *Guidelines for the Preparation of Management Plans for Natura 2000 Areas in Bosnia and Herzegovina with Indicative Management Plans for Tišina, Orjen-Bijela Gora and Vranica*. Prospect C&S, Brussels.
- Golub D., Dekić R., Šukalo G., Sidak S. & Lolić S. 2012. Ichthyofauna diversity of some Vrbas river tributaries as water quality indicator. In: Đukić A. (ed.) *Water 2012, Conference Proceedings of 41th Annual Conference of the Serbian Water Pollution Control Society*: 97–104. Srpsko društvo za zaštitu voda u saradnji sa Institutom za vodoprivredu "Jaroslav Černi", Beograd i "Vujić Valjevo", Valjevo.
- Golub D., Dekić R., Šukalo G., Sidak S. & Živković Đ. 2013. Ichthyofauna composition of artificial hydroaccumulation Sredani (Lijeveće polje) as water quality indicator. In: Đukić A. (ed.) *Water 2013, Conference Proceedings of 42th Annual Conference of the Serbian Water Pollution Control Society*: 181–186. Srpsko društvo za zaštitu voda u saradnji sa Institutom za vodoprivredu "Jaroslav Černi", Beograd.

- Golub D., Đekić T., Dekić R., Šukalo G. & Cvijić S. 2016a. Ichthyofauna Švrakava watercourse as an indicator of water quality. *Conference Proceedings of the Eighth International Conference "Ecology, Health, Work, Sport"*: 152–156.
- Golub D., Marić Ž., Šukalo G., Cvijić S. & Dekić R. 2016b. Morphological characteristics of *Cobitis elongatoides* and *Sabanejewia balcanica* (Cobitidae) from river Suturlija. *Third Symposium of biologists and ecologists of Republic of Srpska*: 139–148.
- Golub D., Bukva M. & Šukalo G. 2017a. Morphometric characteristics of digestive tract of nase (*Chondrostoma nasus*) from Vrbas and Una Rivers. *Skup* 8 (2): 27–35. <https://doi.org/10.7251/SKP170802027G>
- Golub D., Šukalo G. & Ranitović M. 2017b. Digestive tract morphometrics of some cyprinid fishes from River Sava. *Skup* 8 (1): 67–74. <https://doi.org/10.7251/SKP170801067G>
- Golub D., Dekić R., Manojlović M., Friščić J., Šukalo G., Cvijić S. & Lolić S. 2017c. Fish communities as water quality indicator of Ukrina River basin (Republic of Srpska, Bosnia and Herzegovina). In: Đukić A. (ed.) *Water 2017, Conference Proceedings of 46th Annual Conference of the Serbian Water Pollution Control Society*: 141–147. Srpsko društvo za zaštitu voda u saradnji sa Institutom za vodoprivredu "Jaroslav Černi", Belgrade.
- Golub D., Dekić R. & Šukalo G. 2018a. Ichthyofauna as water quality indicator of Jablanica River (Republic of Srpska, B&H). In: Đukić A. (ed.) *Water 2018, Conference Proceedings of 47th Annual Conference of the Serbian Water Pollution Control Society*: 195–201. Srpsko društvo za zaštitu voda u saradnji sa Institutom za vodoprivredu "Jaroslav Černi", Beograd i JKP Drugi oktobar, Vršac.
- Golub D., Lolić S., Dmitrović D., Dekić R., Šukalo G. & Cvijić S. 2018b. Physical, chemical and biological indicators of the Jablanica River water quality (Republic of Srpska, Bosnia and Herzegovina). *Water Research and Management* 8 (4): 11–18.
- Golub D., Bukva M. & Dekić R. 2019. Morphological characteristics and condition of Bleak *Alburnus alburnus* (Teleostei; Cyprinidae) from the Bosna River. *Skup* 10 (1): 3–15. <https://doi.org/10.7251/SKPEN191001016G>
- Grbelja J., Vuković T. & Guzina N. 1980. Imuni odgovor šarana i patuljastog somića na neke biljne viruse (*Cyprinus carpio* Linnaeus, 1758, *Ictalurus nebulosus* Le Sueur, 1819). *Ribarstvo Jugoslavije* 35 (1): 11–12.
- Grgurević M. 1984. *Morfološko-taksonomske Karakteristike Žutalja Rutilus rubilio rubilio (Bonaparte 1837) iz Hutovog Blata*. Diploma thesis. University of Sarajevo, Sarajevo.
- Gunther A. 1859–1870. *A Catalogue of the Fishes in the British Museum*. British Museum, London.
- Guzina N. 2000. *Morfološko-taksonomske Karakteristike Vrste Aulopyge huegeli Heckel, 1841*. PhD thesis, University of Sarajevo, Sarajevo.
- Guzina N. 2005. Osobnosti ždrijelnih zuba vrste *Aulopyge hüegeli* Heckel, 1841. *Veterinaria* 54 (3–4): 225–232.
- Guzina N. & Vuković T. 1972. Mikroelementi željezo i bakar u tkivu jetre i mišića nekih endemičnih vrsta roda *Paraphoxinus*. *Ichthyologia* 3 (1): 19–25.
- Guzina N. & Vuković T. 1977. Serološka proučavanja ciprinida Bosne i Hercegovine. In: *III simpozijum biosistematičara Jugoslavije, Rezimei referata*. Institut za biologiju, Prirodno-matematički fakultet, Novi Sad.
- Guzina N. & Vuković T. 1978. Serološka proučavanja nekih vrsta ciprinida iz voda Bosne i Hercegovine. *Ichthyologia* 10 (1): 55–65.

- Guzina N. & Vuković T. 1987. Prvi podaci o nalazu i morfologiji krljušti od *Aulopyge hügelii* Heckel, 1841 iz Buškog blata. *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 40: 27–38.
- Guzina N., Vuković T. & Miladinović Ž. 1971. Serološka istraživanja nekih vrsta riba iz familije Salmonidae i Cyprinidae (Pisces) iz voda jadranskog sliva u Bosni i Hercegovini. In: Slavnić Ž. (ed.) *I Simpozijum Biosistematičara Jugoslavije. Zbornik Radova*. Sarajevo.
- Guzina N., Vuković T. & Aganović M. 1977. Mikroelementi željezo i bakar u jetri i mišićima *Salmo gairdneri* Richardson, iz različitih ribogajilišta. *Ichthyologia* 9 (1): 75–84.
- Guzina N., Vuković T. & Grbelja J. 1980. Imuni odgovor karaša, *Carassius carassius* (Linnaeus, 1758) na virus žutog mozaika postrne repe. *Ichthyologia* 12 (1): 9–13.
- Guzina N., Vuković T. & Seratlić D. 1981. The artificial hybridization between the species *Phoxinus phoxinus* (Linnaeus) and *Leuciscus cephalus* (Linnaeus). *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 34: 39–46.
- Guzina N., Seratlić D. & Vuković T. 1986. Experimental hybridization between the species *Aulopyge hügelii* Heckel, 1841 and *Leuciscus turskyi* Heckel 1843. *Godišnjak Biološkog Instituta univerziteta u Sarajevu* 39: 39–48.
- Gvozdrenović O., Pavlović V., Kekić H., Mijatović N., Pejić K. & Ivanc A. 1978. Seasonal oscillation of concentration of glucose, lactic acid and glycogen content of some organs of the European eel (*Anguilla anguilla*). *Yugoslavian Physiological and Pharmacological Acta* 14 (1): 85–87.
- Gvozdrenović O., Kekić H., Pavlović V., Ivanc A., Mijatović N. & Pejić K. 1985. Ugljeni hidrati oštrulja (*Aulopyge hügelii* Heckel 1841) u uslovima totalnog gladovanja. *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 38: 77–85.
- Habeković D., Homen Z. & Popović J. 1981. Ihtiofauna akumulacijskog jezera “Modrac” *Croatian Journal of Fisheries: Ribarstvo* 36 (1): 4–7.
- Hadžović J. 1975. *Chondrostoma nasus* iz Rijeke Drine kod Bratunca. Diploma thesis. University of Sarajevo, Sarajevo.
- Habitats Directive 1992. Council Directive 92/43/EEC on the Conservation of natural habitats and of wild fauna and flora. Available from <https://eur-lex.europa.eu/eli/dir/1992/43/oj/eng>
- Hafner-Lahorski R. 1947. Sibirske i daleko-istočne mladice. *Ribarstvo Jugoslavije* 5: 97–99.
- Hafner-Lahorski R. 1948. O zaštiti mladice-glavatice. *Ribarstvo Jugoslavije* 6: 69–70.
- Hajdarević E. 2011. *Hematološke karakteristike Deverike Abramis brama* (Linnaeus, 1758) iz Hidroakumulacije Modrac u Sezonskom Aspektu. PhD thesis, University of Sarajevo, Sarajevo.
- Hajdarević E., Hasanović T., Omeragić A., Adrović A., Skenderović I. & Bajrić A. 2015. Karakteristike erotrocita babuške (*Carassius gibelio*) iz jezera Vidara. *XX. Savjetovanje o Biotehnologiji* 22: 435–439.
- Hajdarević E., Bajrić A., Adrović A., Hercegovac A., Skenderović I., Mehić A., Tanović E. & Mujabašić E. 2018. The hematological and morphometric characteristics of the *Squalius cephalus* of the catchment area in Modrac Lake. *XXIII. Savjetovanje o Biotehnologiji. Zbornik Radova*: 435–439.
- Halilović S.H. & Adrović A. 2015. Morphometric and meristic differences between male and female *Ameiurus melas* in the artificial lake Modrac. *Educa* 8 (8): 21–40.
- Halilović S.H. & Dekić R. 2022. Dužinsko-maseni odnosi i kondiciono stanje babuške (*Carassius gibelio*) iz tri hidroakumulacije u Bosni i Hercegovini. *Hrvatske Vode* 30 (118): 17–25.
- Hamzić A. 2002. Protection on endangered salmonid species in Bosnia and Herzegovina. *EIFAC Symposium on Inland Fisheries and the Aquatic Environment*: 32.

- Hamzić A. 2024. *Biodiverzitet Slatkovodnih Riba Bosne i Hercegovine*. Buybook, Sarajevo.
- Hamzić A. & Lelo S. 2007. Rasprostranjenost linjaka, *Tinca tinca* Linnaeus, 1758, (Actinopterygii: Cypriniformes, Cyprinidae) u Bosni i Hercegovini. *Prilozi Fauni Bosne i Hercegovine* 3: 41–45.
- Hamzić A., Muhamedagić S. & Lelo S. 2011. Peled, *Coregonus peled* (Gmelin, 1789) (Salmoniformes, Salmonidae, Coregoninae) nova vrsta u ihtiofauni Bosne i Hercegovine. *Fauna Bosne i Hercegovine – Biosistematski Prijegledi* 7: 31–36. Udruženje za inventarizaciju i zaštitu životinja, Ilijaš, Kanton Sarajevo.
- Hamzić A., Muhamedagić S. & Lelo S. 2012a. Ohridska pastrmka, *Salmo letnica* (Karaman 1924) (Salmoniformes, Salmonidae, Salmoninae). *Fauna Bosne i Hercegovine – Biosistematski Prijegledi*. Udruženje za inventarizaciju i zaštitu životinja, Ilijaš, Kanton Sarajevo.
- Hamzić A., Lelo S., Muhamedagić S. & Spasojević P. 2012b. Novi i najsjeverniji nalaz gregorca, *Gasterosteus aculeatus* Linnaeus, 1758 u Bosni i Hercegovini. *Prilozi Fauni Bosne i Hercegovine* 9: 15–19.
- Hamzić A., Ugarčin N. & Péteri A. et al. 2015. *Analysis of the Fishery and Aquaculture Sector in Bosnia and Herzegovina*. Food and Agriculture Organization of the United Nations, Regional Office for Europe and Central Asia, Quebec City.
- Hasković E., Glamuzina B., Čulin S. & Dulčić J. 2010. Hematološki parametri oštrulja *Aulopyge hügelii* (Heckel, 1841) u sezonskom aspektu. In: Bogut I., Pintarić A., Treer T., Katavić I., Teskeredžić Z., Glamuzina B., Skaramuca B., Oraić D., Gjurčević E. & Župan B. (eds) *Hrvatsko ribarstvo – kako i kuda dalje? Zbornik sažetaka* 6: 15. *Savjetovanje o Slatkovodnom Ribarstvu Republike Hrvatske s Međunarodnim Sudjelovanjem*. Vukovar.
- Hasković E., Džajić A. & Suljević D. 2011. Biohemijski status krvnog seruma kalifornijske pastrmke *Oncorhynchus mykiss* (Walbaum, 1792) u različitim uslovima držanja i ishrane. *Veterinaria* 3–4: 141–152.
- Hasković E., Mehinović L., Suljević D., Hasković D., Hajdarević E. & Glamuzina B. 2013. Differential blood count of Tench *Tinca tinca* (Linnaeus, 1758) in conditions of thermal stress. *Veterinaria* 62 (3–4): 175–184.
- Has-Schön E., Bogut I., Rajković V., Bogut S., Čačić M. & Horvatić J. 2008. Heavy metal distribution in tissues of six fish species included in human diet, inhabiting freshwaters of the Nature Park “Hutovo Blato” (Bosnia and Herzegovina). *Archives of Environmental Contamination and Toxicology* 54: 75–83. <https://doi.org/10.1007/s00244-007-9008-2>
- Hawlitcshek A. 1895. Die Salmen der österreichischen Gewässer. *Mittheilungen des Österreichischen Fischerei-Vereines* 15 (57): 27–34.
- Hawlitcshek A. 1898. Der Sterlet – *Acipenser ruthenus*. *Mittheilungen des Österreichischen Fischerei-Vereines* 18 (9): 168–175.
- Heckel J. 1843. Ichthyologie. In: Russegger J. Von (ed.) *Reisen in Europe, Asien und Africa, mit Besonderer Rücksicht auf die naturwissenschaftlichen Verhältnisse der betreffenden Länder unternommen (two words) in den Jahren 1835 bis 1841. Erster Band. Reise in Griechenland, Unterägypten, im nördlichen Syrien und südöstlichen Kleinasien. Teil 2*: 991–1099. Schweizerbart, Stuttgart.
- Heckel J. 1852. *Salar dentex*. Heft I–IV. Akademie der Wissenschaften, Wien.
- Heckel J. & Kner R. 1858. *Die Süßwasserfische der Österreichischen Monarchie*. Verlag von Wilhelm Engelmann, Leipzig.
- Hećo Dž. 2000. *Morfometrijske i Merističke Karakteristike Vrste Gasterosteus aculeatus (Linné, 1758) iz Rijeke Bune*. Diploma thesis, University of Sarajevo, Sarajevo.

- Heintz K. 1908. Die Salmoniden Bosniens und der Herzegowina. *Beilage zur Deutschen Anglerzeitung* IX (1): 114–118.
- Helfman G.S. 2007. *Fish conservation: a Guide to Understanding and Restoring Global Aquatic Biodiversity and fishery Resources*. Island Press, Washington DC.
- Hewitt G.M. 1999. Post-glacial re-colonization of European biota. *Biological Journal of Linnean Society* 68: 87–112.
- Hirc D. 1896. Kako ribare na Dunavu i drugim nekim našim vodama? *Lovačko-ribarski Vestnik* 5: 124–127.
- Hirc D. 1900. Pastrva - prinos ihtiofauni Hrvatske i Slavonije. *Lovačko-ribarski Vestnik* 9: 70–72.
- Hirc D. 1902. Glavatica i “Lachsforelle”. *Glasnik Hrvatskoga Naravoslovnoga Društva* 4–6: 455–457.
- Holčik J. & Delić A. 2000. New discovery of the Ukrainian brook lamprey in Croatia. *Journal of Fish Biology* 56: 73–86. <https://doi.org/10.1111/j.1095-8649.2000.tb02087.x>
- Holčik J. & Mrakovčić M. 1997. First record of *Lethenteron zanandreae* (Cyclostomata, Petromyzontidae) in the Adriatic basin of the Balkan peninsula and its zoogeographic consequences. *Folia Zoologica* 46: 263–271.
- Holčik J., Delić A., Kučinić M., Bukvić V. & Vater M. 2004. Distribution and morphology of the sea lamprey *Petromyzon marinus* in the Balkanian coast of the Adriatic Sea. *Journal of Fish Biology* 64: 514–527. <https://doi.org/10.1111/j.0022-1112.2004.00318.x>
- Horvat M. 1964. Rast lipljena u rijeci Uni. *Ribarstvo Jugoslavije* 4: 103–111.
- Horvat Š. 1901. *Ribe i Ribogojstvo*. Društvo sv. Jeronima, Zagreb.
- Horvatić S., Marčić Z., Mrakovčić M., Mustafić P., Buj I., Čaleta M. & Zanella D. 2017. Threatened fishes of the world: *Orsinogobius croaticus* (Mrakovčić, Kerovec, Mišetić & Schneider, 1996) (Teleostei: Gobiidae). *Croatian Journal of Fisheries, Ribarstvo* 75 (1): 30–35. <https://doi.org/10.1515/cjf-2007-0006>
- Howes G.J. 1985. A revised synonymy of the minnow genus *Phoxinus* Rafinesque, 1820 (Teleostei, Cyprinidae) with comments on its relationships and distribution. *Bulletin of the British Museum (Natural History), Zoology Series* 48: 57–74. <https://doi.org/10.5962/p.219626>
- Howes G.J. 1987. The phylogenetic position of the Yugoslavian cyprinid fish genus *Aulopyge* Heckel, 1841, with an appraisal of the genus *Barbus* Cuvier and Cloquet, 1816 and the subfamily Cyprinidae. *Bulletin of the British Museum (Natural History), Zoology Series* 52: 165–196. <https://doi.org/10.5962/p.203885>
- Ibrašimović A. 1988. *Krljušti Vrste Chondrostoma phoxinus Heckel, 1843. iz Buškog Blata*. Diploma thesis, University of Sarajevo, Sarajevo.
- Imamović V. 1986. Paraziti i parazitoze riba u nekim salmonidnim ribogojilištima u Bosni i Hercegovini I. Ihtiobodoza i Heksamitijaza. *Veterinaria* 35 (1): 47–66.
- Imamović V. 1987. Parasites and parasitoses of fishes in some salmonid fish ponds in Bosnia – Hercegovina II. Ichtyophthiriasis and chylodonellosis. *Veterinaria* 36 (2):151–158.
- Ivanc A. & Dekić R. 2012. Ekofiziološka istraživanja autohtonih vrsta riba Republike Srpske. *Skup* 4 (1): 26 – 35.
- Ivanc A., Kekić H., Lazarević-Ivanc Lj., Gvozdenović O., Pavlović V., Pejić K. & Mijatović N. 1989. Serum lipids of *Aulopyge hügelii* Heckel during starvation. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 42: 29–43.

- Ivanc A., Dekić R., Četković D., Mišćević B., Könyves T., Vukosav M., Obradović S. & Suturović E. 2011. Evaluation of environmental conditions by fish hematology. *Agrár-és Vidékfejlesztési Szemle* 6 (suppl.): 130–136.
- Ivanišević B. 1962. *Uporedba Morfologije i Rasta Scardinius erythrophthalmus scardafa (Bonaparte) u Donjoj Neretvi i u Skadarskom Jezeru*. Diploma thesis, University of Sarajevo, Sarajevo.
- Ivanković P., Treer T., Piria M. & Knezović Z. 2010. Diet and growth of endemic Neretva chub, *Squalius svallize* from the Neretva river area, Bosnia and Herzegovina. *Folia Zoologica* 59 (1): 51–56. <https://doi.org/10.25225/fozo.v59.i1.a8.2010>
- Ivanković P., Piria M., Treer T. & Knezović Z. 2011. Meristic and morphometric characteristics of endemic Neretva chub, *Squalius svallize* from the Neretva River area, Bosnia and Herzegovina. *Journal of Applied Ichthyology* 27: 1031–1032. <https://doi.org/10.1111/j.1439-0426.2011.01741.x>
- Ivanković P., Treer T., Tomljanović T., Knezović Z. & Salaj S. 2017. Back-calculating growth at age from scale readings for *Squalius microlepis* Heckel, 1843 and *Scardinius plotizza* Heckel & Kner, 1858 in the Neretva River basin (Bosnia and Herzegovina). *Journal of Applied Ichthyology* 33 (3): 546–549. <https://doi.org/10.1111/jai.13349>
- Ivanović B. 1968. Ekologija *Alburnus alburnus alborella* (Filippi). *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 20: 5–69.
- Ivanović B. 1972. *Scardinius erythrophthalmus scardafa* (Bonaparte 1832). *Ichthyologia* 4 (1): 33–39.
- Ivanović B. & Sekulović T. 1971. Taksonomske karakteristike *Leuciscus cephalus albus* Bonaparte, 1938 iz Skadarskog jezera. *Ichthyologia* 3 (1): 27–36.
- IUCN. 2025. The IUCN Red List of Threatened Species. Version 2024–2. Available at <https://www.iucnredlist.org> [22 Aug. 2025]
- Jakovlić I., Wu Q.J., Treer T., Šprem N. & Gui J.F. 2013. Introgression evidence and phylogenetic relationship among three (*Para*)*Misgurnus* species as revealed by mitochondrial and nuclear DNA markers. *Archives of Biological Sciences* 65: 1463–1467. <https://doi.org/10.2298/ABS1304463J>
- Jakovlić I., Piria M., Šprem N., Tomljanović T., Matulić D. & Treer T. 2015. Distribution, abundance and condition of invasive Ponto-Caspian gobies *Ponticola kessleri* Günther, 1861), *Neogobius fluviatilis* (Pallas, 1814), and *Neogobius melanostomus* (Pallas, 1814) in the Sava River basin, Croatia. *Journal of Applied Ichthyology* 33: 888–894. <https://doi.org/10.1111/jai.12803>
- Jamaković A. 2011. *Morfološke Karakteristike Digestivnog Trakta Smuđa - Sander lucioperca (Linnaeus, 1758) iz Nerentvanskih Hidroakumulacija*. Diploma thesis, University of Sarajevo, Sarajevo.
- Jankelić D. 1963. *Morfološke Karakteristike Rhodeus Sericeus amarus iz Okoline Banja Luke*. Diploma thesis, University of Sarajevo, Bosnia and Herzegovina.
- Janković D. 1960. *Sistematika i Ekologija Lipljena Jugoslavije*. Biološki institut, Stručno udruženje za unapređenje slatkovodnog ribarstva, Beograd.
- Janković D. 1961. Taksonomska i ekološka ispitivanja na mekousnoj pastrmci (*Salmo obtusirostris oxyrhynchus* S.) iz reke Bune. *Zbornik Radova Biološkog Instituta NR Srbije. Knjiga* 5 (4–5): 1–31. Naučno delo, Beograd.
- Jardas I. 1996. *Jadranska ihtiofauna*. Školska knjiga, Zagreb.
- Jedlička D. 1931. Pljeva i Janj. *Ribarski List* 7:1–4.
- Jedlička D. 1932. O rastu lipljena. *Ribarski List*: 11–12.
- Jedlička D. 1935. Važnije ribolovne vode Vrbaske Banovine. *Ribarski Vjesnik* 9–10: 225–230.

- Jedlička D. 1947. *Ribe Narodne Republike Bosne i Hercegovine*. Sarajevo.
- Jelić D., Duplić A., Čaleta M. & Žutinić P. 2008. *Endemske Vrste Riba Jadranskog Sliva*. AZO-Agencija za zaštitu okoliša, Zagreb.
- Jerković L., Aganović M. & Remeta D. 1973. Prilog poznavanju ishrane podbile *Chondrostoma phoxinus* Heckel. *I Europski kongres ihtiologa, Rezime referata*: 62.
- Jojić S. 1984. *Morfometrijske i Merističke Karakteristike Vrste Aulopyge hugeli*. Diploma thesis, University of Sarajevo, Sarajevo.
- Jukić M. 2008. The River Sava as a resource in spatial planning. *Geoadria* 13: 81–96.
- Juretić N., Vuković T., Grbelja J. & Guzina N. 1977. Immunogenicity of turnip yellow mosaic virus in green tench (*Tinca tinca*). *Naturwissenschaften* 64: 440–441. <https://doi.org/10.1007/BF00508708>
- Jurinac A.E. 1887. Prilog fauni zapadne Slavonije. *Glasnik Hrvatskoga Naravoslovnoga Društva* 1–3: 21–34.
- Kaćanski D. & Kosorić Đ. 1970. O ishrani potočne pastrmke (*Salmo trutta m. fario* L.) iz nekih tekućica jadranskog sliva u SRBiH. *Ichthyologia* 2 (1): 63–71.
- Kaćanski D., Kosorić Đ. & Čepić V. 1977. O ishrani nekih vrsta riba u slivu Neretve (od Uloga do Mostara). *Ichthyologia* 9 (1): 31–45.
- Kaćanski D., Jerković L., Hafner D. & Aganović M. 1978. O ishrani nekih vrsta riba iz Buškog Blata. *Ichthyologia* 10 (1): 67–75.
- Kaćanski D., Hafner D., Blagojević S. & Kosorić Đ. 1981. O ishrani nekih ciprinidnih vrsta riba iz rijeke Bosne. *Ichthyologia* 13 (1): 51–62.
- Kadribašić S. 2005. *Morfometrijske i Merističke Karakteristike zele Alburnus alburnus (Linnaeus, 1758) iz rijeke Save*. Diploma thesis, University of Sarajevo, Sarajevo.
- Kadribegović E. 2005. *Karakteristike Ishrane Potočne Pastrmke Salmo trutta m. fario (Linnaeus, 1758) iz Rijeke Rakitnice*. Diploma thesis, University of Sarajevo, Sarajevo.
- Kalamujić B. 2008. *Molekularno-genetički Diverzitet Populacija Lipljena (Thymallus thymallus L.) u Bosni i Hercegovini*. Master's Thesis, University of Sarajevo, Sarajevo.
- Kalamujić B. 2013. *Genetički Diverzitet Recentnih Populacija Salmo trutta L. Rijeke Neretve u Odnosu na Rekonstruisani Simulacioni Model Početnog Stanja*. PhD Thesis, Univerzitet u Sarajevu, Institut za genetičko inženjerstvo i biotehnologiju, Sarajevo.
- Kalamujić B., Durmić-Pašić A., Pojskić N., Škrijelj R. & Hadžiselimović R. 2007. Genetic diversity of grayling (*Thymallus thymallus* L.) populations in Bosnia and Herzegovina. In: Buj I., Zanella L., & Mrakovčić M. (eds) *12th European Congress of Ichthyology. Book of Abstracts* 13: 32. Croatian Ichthyological Society, Cavtat (Dubrovnik).
- Kalamujić B., Škrijelj R. & Pojskić N. 2015. Genetic diversity of contemporary populations of *Salmo trutta* L. in the basin of the River Neretva, Bosnia and Herzegovina. *Frontiers in Marine Science. Conference Abstract: XV European Congress of Ichthyology*. <https://doi.org/10.3389/conf.fmars.2015.03.00049>
- Kalamujić Stroil B., Mušović A., Škrijelj R., Dorić S., Đug S. & Pojskić N. 2019. Molecular-genetic diversity of the endangered Dalmatian barbelgudgeon, *Aulopyge huegelii* from the Buško Blato reservoir. *Genetica* 147: 269–280. <https://doi.org/10.1007/s10709-019-00069-z>
- Kalamujić Stroil B., Stanić M., Kalajdžić A. & Pojskić N. 2020. Diversity of brook trout in Bosnian hatcheries. *Croatian Journal of Fisheries* 78 (4): 183–194. <https://doi.org/10.2478/cjf-2020-0018>

- Kalamujić Stroil B., Ušanović L., Kalajdžić A., Lasić L., Škrijelj R. & Pojskić N. 2021. Molecular structure and hybridization patterns of *Abramis brama* × *Rutilus rutilus* hybrids from Modrac reservoir, Bosnia and Herzegovina. *Croatian Journal of Fisheries* 7: 147–156. <https://doi.org/10.2478/cjf-2021-0016>
- Kalous L., Rylkov'a K., Bohlen J., Šanda R. & Petrtyl M. 2012. New mtDNA data reveal a wide distribution of the Japanese ginbuna *Carassius langsdorfii* in Europe. *Journal of Fish Biology* 82 (2): 703–707. <https://doi.org/10.1111/j.1095-8649.2012.03492.x>
- Kaltak A. 2009. *Morfoanatomske Osobenosti Spolnih Žlijezda Sapače Barbus meridionalis petenyi (Heckel, 1847) iz Rijeka Bosne i Stavnje*. Diploma thesis, University of Sarajevo, Sarajevo.
- Kaluđerčić M., Kaluđerčić D. & Kosorić Đ. 1970. Rezultati ispitivanja serum proteina krvi *Salmo trutta m. fario* L. lovljenih u raznim lokalitetima. In: *III Simpozijum "Savremena Ihtiološka Istraživanja"*: 25–26. Yugoslav Ichthyological Society, Kotor.
- Kaluđerčić M., Kaluđerčić D., Kosorić Đ., Pavlić S. & Vuković T. 1971. Serum proteini krvi nekih slatkovodnih riba Bosne i Hercegovine. *Ichthyologia* 3 (1): 37–48.
- Kapetanović N. & Vuković T. 1968. Nalaz *Leuciscus svalizze* Heckel et Kner u estavelama Popovog polja. *Ribarstvo Jugoslavije* 3: 33–67.
- Kapetanović N., Aganović M. & Vuković T. 1966. O specifičnostima oblika otolita kod jedinki iz tri lokalne populacije oštrulja (*Aulopyge hügelii* Heckel). *Ribarstvo Jugoslavije* 21 (6): 17–22.
- Kapetina S. 1981. *Morfološke Karakteristike Vrste Gobio gobio (Linnaeus, 1758) iz Rijeke Bosne kod Donje Moštre*. Diploma thesis, University of Sarajevo, Sarajevo.
- Kaplanović Z. 1984. *Morfometrijski i Meristički Karakteri Vrste Leuciscus cephalus (Linnaeus, 1758) iz rijeke Mlječanice*. Diploma thesis, University of Sarajevo, Sarajevo.
- Karaman M. 1963. Eine neue Unterart des Steinpeitzgers, *Cobitis aurata bosniaca* n. ssp. (Cobitidae, Pisces), aus Jugoslawien. *Internationale Revue der gesamten Hydrobiologie und Hydrographie* 48 (4): 629–632.
- Karaman M.S. 1972. Süßwasserfische der Türkei. 9. Teil. Revision einiger kleinwüchsiger Cyprinidengattungen, *Phoxinellus*, *Leucaspius*, *Acanthobrama* usw. aus Südeuropa, Kleinasien, Vorderasien und Nordafrika. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 69: 115–155.
- Karaman S. 1922. Über eine neue *Cobitis*-Art aus Jugoslavien, *Cobitis balcanica* n. sp. *Glasnik Hrvatskoga Prirodoslovnoga Društva* 34: 1–4.
- Karaman S. 1923. Filogenetska važnost broja ljusaka i kralježaka kod Cyprinida. *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 25: 25–37.
- Karaman S. 1924. *Pisces Macedoniae*. Hrvatska štamparija, Split.
- Karaman S. 1926. Salmonidi Balkana, *Glasnik Skopskog Naučnog Društva* 1: 253–268.
- Karaman S. 1927. Salmonidi Balkana. *Glasnik Skopskog Naučnog Društva* 2: 253–269.
- Karaman S. 1928. Prilozi ihtiologiji Jugoslavije. I. *Glasnik Skopskog Naučnog Društva* 6 (2): 147–176.
- Karaman S. 1938. Beiträge zur Kenntnis der Süßwasserfische Jugoslaviens. *Glasnik Skopskog Naučnog Društva* 18: 131–139.
- Katurić M. 1883. Notizie Zoologische. *Estratto dal Bollettino della Societa Adriatica di Scienze Naturali* 8 (1): 1–9.
- Katurić M. 1903. Novo nalazište jesetre atlantske, *Acipenser sturio* L. *Rad Jugoslavenske Akademije Znanosti i Umjetnosti* 33: 1–6.

- Kekić H. 1967. Sezonske oscilacije broja eritrocita i koncentracije hemoglobina u krvi klena (*Leuciscus cephalus* L.) iz srednjeg toka rijeke Željeznice. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 20: 25–34.
- Kekić H. 1968. Contribution to the study of seasonal variations in the number of erythrocytes and haemoglobin concentration in the blood of *Chondrostoma nasus* L. from the middle flow of Željeznica river. *Yugoslavian Physiological and Pharmacological Acta* 4: 41–47.
- Kekić H., Ivanc A., Pavlović V., Gvozdenović O., Mijatović N. & Pejić K. 1983. The differential leukocyte count of Dalmatian barbelgudgeon (*Aulopyge hügelii*) under the conditions of prolonged starvation. *Yugoslavian Physiological and Pharmacological Acta* 19 (1): 77.
- Kerkez V., Dekić R. & Ivanc A. 2014. Physiology of Brown trout (*Salmo trutta*) nutrition from the rivers Ugar and Pliva. *Skup* 6 (1): 48–58.
- Kerovec M., Mrakovčić M., Schneider D. & Tomašković N. 1998. Occurrence of *Chondrostoma phoxinus* in Croatia. *Folia Zoologica* 47 (1): 39–43.
- Kesterčanek F.X. 1886. Naše ribarstvo. *Šumarski List* 10: 25–33.
- Keško N. 2018. Kvalitativno-kvantitativni sastav ihtiopopulacije rijeke Lepenice. *Water And We, Journal of the Agency for the Sava River Water Area* 97: 42–48.
- Ketmaier V., Bianco P.G., Cobolli M., Krivokapić M., Caniglia R. & De Matthaëis E. 2004. Molecular phylogeny of two lineages of Leuciscinae cyprinids (*Telestes* and *Scardinius*) from the peri-Mediterranean area based on cytochrome b data. *Molecular Phylogenetics and Evolution* 32 (3): 1061–1071. <https://doi.org/10.1016/j.ympev.2004.04.008>
- Ketmaier V., Bianco P.G. & Durand J.D. 2008. Molecular systematics, phylogeny and biogeography of roaches (Rutilus, Teleostei, Cyprinidae). *Molecular Phylogenetics and Evolution* 49: 362–367. <https://doi.org/10.1016/j.ympev.2008.07.012>
- Kiškarolj M. 1965. Cestodi i nematodi salmonidnih i nekih drugih riba Bosne i Hercegovine. *Veterinaria* 14 (2): 235–247.
- Kiškarolj M. & Čanković M. 1968. *Pomphorhynchus bosniacus* nov. sp. aus Barben *Barbus barbatus* (L.) des Save-Gebietes. *Zoologischer Anzeiger* 182: 69–74.
- Kiškarolj M. & Čanković M. 1969. Neke još neutvrđene vrste endohelminata slatkovodnih riba u Jugoslaviji. *Ichthyologia* 1 (1): 43–55.
- Kišpatić M. 1893. *Ribe*. Tisak Karla Albrechta, Zagreb.
- Knežević B. 1977. The fecundity of *Rutilus rubilio rubilio* (Bonaparte, 1837). *Glas Republičkog Zavoda za Zaštitu Prirode – Prirodnjačkog Muzeja* 10: 69–74.
- Knežević B. & Ivanović B. 1975. Sezonsko razviće gonada *Rutilus rubilio* (Bp). *Ichthyologia* 7: 17–24.
- Knop F. 1947. Bolen (*Aspius rapax*). *Ribarstvo Jugoslavije* 5: 78.
- Kolar C.S. & Lodge D.M. 2001. Progress in invasion biology: predicting invaders. *Trends in Ecology & Evolution* 16 (4): 199–204. [https://doi.org/10.1016/S0169-5347\(01\)02101-2](https://doi.org/10.1016/S0169-5347(01)02101-2)
- Kolombatović J. 1886. Imenik kralješnjaka Dalmacije. II Dio: Dvoživci, gmazovi i ribe. *Godišnje Izvješće Velike Realke*: 1–20.
- Kolombatović J.. 1907. Contribuzioni all fauna dei vertebrati della Dalmazia. *Glasnik Hrvatskoga Naravoslovnoga Društva* 19: 1–24.
- Kopjar N., Mustafić P., Zanella D., Buj I., Čaleta M., Marčić Z., Milić M., Dolenc Z. & Mrakovčić M. 2008. Assessment of DNA integrity in erythrocytes of *Cobitis elongata* affected by water pollution, the alkaline comet assay study. *Folia Zoologica* 57 (1–2): 120–130.

- Korić H. 1975. *Aulopyge hugeli* Šujice i Buškog Blata. Diploma thesis, University of Sarajevo, Sarajevo.
- Korjenić E. 1998. Morfološke karakteristike digestivnog trakta kod *Leuciscus souffia agassizi* iz gornjeg toka rijeke Drine. *Veterinaria* 47 (1–2): 131–142.
- Korjenić E. 1999. Morfološke karakteristike digestivnog trakta kod *Alburnus alburnus alburnus* iz gornjeg toka rijeke Drine. *Veterinaria* 48 (1–2): 99–109.
- Korjenić E. 2003. Ishrana potočne pastrmke – *Salmo trutta m. fario* Linnaeus, 1758. iz sliva rijeke Fojnice. *Radovi Poljoprivrednog Fakulteta, Univerziteta u Sarajevu* 48 (53): 19–26.
- Korjenić E. 2004a. Odnos sastava faune dna i ishrane lipljena *Thymallus thymallus* (Linnaeus, 1758) iz sliva rijeke Fojnice. *Veterinaria* 53 (1): 13–24.
- Korjenić E. 2004b. Varijacija osnovnih tjelesnih mjera nekih vrsta riba iz sliva rijeke Fojnice. *Veterinaria* 53 (1): 25–35.
- Korjenić E. 2005a. Odnos sastava faune dna i ishrana pliske *Alburnoides bipunctatus* (Bloch, 1872) iz rijeke Fojnice. *Veterinaria* 54 (1–2): 61–68.
- Korjenić E. 2005b. Populacije riba rijeke Gvožđanke. *Veterinaria* 54 (3–4): 233–244.
- Korjenić E. 2006a. Odnos sastava faune dna i ishrane evropskog kljena *Leuciscus cephalus* (Linnaeus, 1758) iz rijeke Fojnice. *Veterinaria* 55 (1–4): 101–109.
- Korjenić E. 2006b. Morfološko-taksonomske karakteristike evropskog kljena *Leuciscus cephalus* (Linnaeus, 1758) iz rijeke Bosne. *Veterinaria* 55 (1–4): 117–130.
- Korjenić E. 2007. Praćenje tempa rasta predkonzumne potočne pastrmke – *Salmo trutta m. fario* Linnaeus, 1758 na ribogojilištu “Žunovnica” u Hadžićima. *Zbornik Radova Prirodno-matematičkog Fakulteta Univerziteta u Tuzli* 3 (3–4): 97–115.
- Korjenić E., Škrijelj R. & Guzina N. 2003. Dobna i spolna struktura populacija riba hidroakumulacije Grabovica. *Veterinaria* 52 (1–4): 87–96.
- Korjenić E., Kadribašić S. & Žujo D. 2005. Morfometrijske i merističke karakteristike zeke – *Alburnus alburnus* (Linnaeus, 1758) iz rijeke Save. *Zbornik radova Prirodno-matematičkog Fakulteta Univerziteta u Tuzli Univerziteta u Tuzli* 2 (2): 81–92.
- Korjenić E., Karović S. & Žujo D. 2006. Morfometrijske i merističke karakteristike mreke – *Barbus barbus* (Linnaeus, 1758) iz rijeke Save. *Hrvatski Glasnik Sveučilišta u Mostaru* 18: 149–160.
- Korjenić E., Džano A. & Šuta N. 2009. Populacije riba rijeke Požarne. *Veterinaria* 58 (1–2): 55–62.
- Korovina V.M. & Vuković T. 1972. Anatomsko-histološke osobine srednjeg creva *Salmothymus obtusirostris oxyrhynchus* (Steindachner 1882). *Ichthyologia* 4 (1): 41–38.
- Kosorić Đ. 1965. Prvi podaci o ribarstvu i njegovom iskorišćavanju u budućoj ramskoj vodenoj akumulaciji. *Ribarski List SRBiH* 40 (1): 9–12.
- Kosorić Đ. 1969. Neki podaci o rasprostranjenju i migracijama neretvanske glavatice (*Salmo marmoratus* Cuv.) u slivu Neretve. *Glasnik Zemaljskog Muzeja* 8: 79–82.
- Kosorić Đ. 1972a. Some data on distribution and migration of the *Salmo marmoratus* Cuv. in tributary of the Neretva. *Glasnik Zemaljskog muzeja* Band II, Heft C: 89–92.
- Kosorić Đ. 1972b. Uticaj otpadnih voda na slatkovodno ribarstvo u SRBiH, sa posebnom analizom stanja u rijeci Bosni. *Vodoprivreda* 21–24: 157–177.
- Kosorić Đ. 1974. Ribe rijeke Rame. *Ichthyologia* 6 (1): 69–78.

- Kosorić Đ. 1976. Mješovite populacije riba kao biološki parametar kvaliteta vode gornjeg toka rijeke Bosne u oblasti narušenoj ekspoatacijom uglja i industrijom. *Glasnik Zemaljskog Muzeja* XV: 201–214.
- Kosorić Đ. 1977a. Populacije riba srednje Neretve, sa projekcijom razvojnih mogućnosti poslije izgradnje vodenih akumulacija. *Ichthyologia* 9 (1): 121–129.
- Kosorić Đ. 1977b. Naselje neretvanske mekousne (*Salmothymus obtusirostris oxyrhynchus* Steind.) u slivu Neretve. *Ribarski List* 4: 81–82.
- Kosorić Đ. 1978. Sastav populacije riba Hutova blata. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 31: 69–81.
- Kosorić Đ. 1981. The fish populations of the Krivaja river. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 34: 57–73.
- Kosorić Đ. & Kardoš N. 1955. Otpadne vode industrije u Zenici i njihov uticaj na riblji fond u rijeci Bosni. *Ribarski List* 30 (3): 49–53.
- Kosorić Đ. & Kačanski D. 1959. Prilog ihtio–biološkom poznavanju gornjeg sliva rijeke Usore. *Ribarstvo Jugoslavije* 14 (2): 31–35.
- Kosorić Đ. & Mikavica D. 1986. Sastav populacije riba rijeka Sanice i Korčanice. *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 39: 117–125.
- Kosorić Đ. & Vuković T. 1966a. Ribe rijeke Bune. *Glasnik Zemaljskog Muzeja Bosne i Hercegovine u Sarajevu* 5: 179–190.
- Kosorić Đ. & Vuković T. 1966b. Pokretljivosti spermatozoida babura (*Rutilus rubilio*) pri različitim salinitetima i temperaturama. *Ribarstvo Jugoslavije* 21 (6): 14–17.
- Kosorić Đ. & Vuković T. 1969. Ispitivanja mogućnosti hibridizacije salmonidnih vrsta sliva Neretve. *Ichthyologia* 1 (1): 57–67.
- Kosorić Đ. & Vuković T. 1971. O mriješćenju glavatice *Salmo marmoratus* Cuv. (Pisces, Salmonidae) u slivu Neretve. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 24: 111–115.
- Kosorić Đ., Kačanski D. & Veledar I. 1971. Sadašnje stanje ribarstva ramske vodene akumulacije i dalje mjere unapređenja. *Ribarstvo Jugoslavije* 26 (3): 54–57.
- Kosorić Đ., Kapetanović N. & Veledar I. 1975. Ribe Velikog Zlatarskog jezera. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 28: 133–141.
- Kosorić Đ., Kapetanović N. & Mikavica D. 1980. The composition and structure of Fish populations in the Bosna river from Visoko to mouth into Sava river. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 33: 99–113.
- Kosorić Đ., Vuković T., Kapetanović N., Guzina N. & Mikavica D. 1983. Sastav naselja riba rijeke Neretve u Bosni i Hercegovini. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 36: 117–129.
- Kosorić Đ., Vuković T., Guzina N., Kapetanović N. & Mikavica D. 1989. Ihtiofauna rijeke Neretve i njene promjene posljedično izgradnji hidroelektrana. *Savjetovanje o Ribarstvu na Hidroakumulacijama, Zbornik Radova*: 133–138.
- Koutsikos N., Zogaris S., Vardakas L., Tachos V., Kalogianni E., Šanda R., Chatzinikolaou Y., Giakoumi S., Economidis P.S., & Economou A.N. 2012. Recent contributions to the distribution of the freshwater ichthyofauna in Greece. *Mediterranean Marine Science* 13: 268–277. <https://doi.org/10.12681/mms.308>
- Kovačić M. 2005. A new species of *Knipowitschia* (Gobiidae) from Dalmatia, Croatia. *Cybium* 29: 275–280. <https://doi.org/10.26028/cybium/2005-293-005>

- Kovačić M. & Pallaoro A. 2003. Is *Knipowitschia caucasica*-like form from the Adriatic Sea a new goby species? Evidence from a morphological approach in the eastern Adriatic Sea. *Cybium* 27 (2): 131–136.
- Kovačević D., Jovanić D., Todorović S., Panić G., Radošević D. & Davidov B. 2018. *Stručno Mišljenje za Prethodnu Zaštitu Rijeke Une*. Republički zavod za zaštitu kulturno–istorijskog i prirodnog nasljeđa Republike Srpske, Banja Luka.
- Kottelat M. 1997. European freshwater fishes. An heuristic checklist of the freshwater fishes of Europe (exclusive of former USSR), with an introduction for nonsystematists and comments on nomenclature and conservation. *Biologia* 52 (Suppl 5): 1–271.
- Kottelat M. 2007. Three new species of *Phoxinus* from Greece and southern France (Teleostei, Cyprinidae). *Ichthyological Exploration of Freshwaters* 18: 145–162.
- Kottelat M. & Freyhof J. 2007. *Handbook of European Freshwater Fishes*. Kottelat, Cornol, and Freyhof, Berlin.
- Kujačić Lj. 1987. *Morfometrijski i Meristički Karakteri Vrste Rutilus rutilus (L. 1758) iz Rijeke Vrbas i Ljevcanice*. Diploma thesis, University of Sarajevo, Sarajevo.
- Kurtović B., Kapetanović D. & Matašin Ž. 2007. Usporedba histoloških promjena u organima klena (*Squalius cephalus*) s brojem heterotrofnih i koliformnih bakterija u vodi rijeke Save. *Ribarstvo* 4: 127–140.
- Kurtović B., Vardić I., Valić D., Kapetanović D., Teskeredžić Z. & Teskeredžić E. 2009. Zdravstveno stanje klena (*Squalius cephalus*) u odnosu na kvalitetu vode rijeke Save. *Ribarstvo* 2: 63–75.
- Landiges W. & Vogt D. 1965. *Die Süßwasserfische Europas bis zum Ural und Kaspischen Meer*. Parey, Hamburg.
- Landeka N., Podnar M. & Jelić D. 2015. New data on the taxonomic status and distribution of *Gambusia* sp. in Croatia and Bosnia and Herzegovina. *Periodicum Biologorum* 117 (3): 415–424. <https://doi.org/10.18054/pb.2015.117.3.3143>
- Langhoffer A. 1904. Popis riba koje su prispjele Narodnom zoološkom muzeju u Zagrebu do konca godine 1900. *Glasnik Hrvatskoga Naravoslovnoga Društva* 16: 148–169.
- Leiner S. 1984. Rod *Paraphoxinus* (Cyprinidae, Pisces) u vodama Jugoslavije. Reliktne ribe ponornica Jadranskog sliva. *Speleolog* 28: 13–18.
- Leiner S. 1998. Endangered ichthyocoenoses of the Sava and Odra rivers in the Turopolje area. *Natura Croatica* 1: 31–43.
- Lelo S. & Hamzić A. 2020. Largemouth black bass, *Micropterus salmoides* (La Cepède, 1802) (Chordata: Vertebrata: Actinopterygii), in Bosnia and Herzegovina. *Prilozi Fauni Bosne i Hercegovine* 16: 39–43.
- Lever C. 1996. *The Naturalised Fishes of the World*. CA Academic Press, San Diego.
- Levin B., Ermakov A., Ermakov O., Levina M., Sarycheva O. & Sarychev S. 2016. Ukrainian brook lamprey *Eudontomyzon mariae* (Berg): phylogenetic position, genetic diversity, distribution, and some data on biology. In: Orlov A. & Beamish R. (eds) *Jawless Fishes of the World, Volume 1*. Cambridge Scholars Publishing, Newcastle upon Tyne.
- Lolić S., Dekić R., Ivanc A., Manojlović M., Erić Ž., Janjić N. & Golub D. 2015. Features of *Delminichthys ghetaldii* (Steindachner, 1882) habitat. In: Đukić A. (ed.) *Water 2015. Conference Proceedings of 45th Annual Conference of the Serbian Water Pollution Control Society*: 225–230. Serbian Water Pollution Control Society, Zlatibor.
- Lolić S., Golub D., Dekić R., Manojlović M. & Paspalj J. 2019. Water quality of some tributaries of river Drina in Republic of Srpska (B&H): microbiological and ichthyological research. *Water 2019, Conference*

Proceedings of 48th Annual Conference of the Serbian Water Pollution Control Society: 205–213. Serbian Water Pollution Control Society, Zlatibor.

Lončar D. 1970. *Osnovne karakteristike hromosomske garniture "podbile" (Chondrostoma phoxinus Heckel)*. Diploma thesis, University of Sarajevo, Sarajevo.

Lucentini L., Chiesa S., Giannetto D., Pompei L., Natali M., Sala P., Volta P., Lorenzoni M. & Fontaneto D. 2014. Integrative taxonomy does not support the occurrence of two species of the *Squalius squalus* complex (Actinopterygii, Cypriniformes, Cyprinidae) in Italy. *Biochemical Systematics and Ecology* 56: 281–288. <https://doi.org/10.1016/j.bse.2014.07.005>

Ludoški J., Francuski Lj., Lukač M., Dekić D. & Milankov V. 2021. Toward the conservation of the endemic monotypic fish genus *Aulopyge* from the Balkan Dinaric karst: Integrative assessment of introduced and natural population. *Ecology and Evolution* 11 (2): 688–699. <https://doi.org/10.1002/ece3.7108>

Lukač M., Ludoški J., Dekić R. & Milankov V. 2023. Evolutionary body shape diversification of the endemic Cyprinoidei fishes from the Balkan's Dinaric karst. *Organisms Diversity & Evolution* 23: 983–1004. <https://doi.org/10.1007/s13127-023-00615-w>

Lukač M., Ludoški J., Dekić R. & Milankov V. 2024. Evolutionary history of the Balkan endemic genus *Delminichthys* (Teleostei, Leuciscidae) with an emphasis on the population structure of Southern Dalmatian minnow. *Contributions to Zoology*: 1–42. <https://doi.org/10.1163/18759866-bja10068>

Lukić T., Stanić-Koštroman S. & Herceg N. 2016. The state of biodiversity of the Sava River through comparative analysis NATURA 2000 species in Croatia and Bosnia and Herzegovina. In: Habuda-Stanić M. (ed.) *2nd International and 6th Croatian Scientific and Professional Conference Water for all*: 175–184. Faculty of food technology Osijek, Osijek.

Ljubojević M. 1989. *Morfometrijski i Meristički Karakteri Vrsta Rutilus rubilio i Rutilus rutilus*. Diploma thesis, University of Sarajevo, Sarajevo.

Machordom A. & Doadrio I. 2001. Evolutionary history and speciation modes in the cyprinid genus *Barbus*. *Proceedings of the Royal Society, Biological Sciences* 268 (1473): 1297–1306. <https://doi.org/10.1098/rspb.2001.1654>

Mäkinen H.S. & Merilä J. 2008. Mitochondrial DNA phylogeography of three-spined stickleback (*Gasterosteus aculeatus*) in Europe – Evidence for multiple glacial refugia. *Molecular Phylogenetics and Evolution* 46 (1): 167–182. <https://doi.org/10.1016/j.ympev.2007.06.011>

Mäkinen H.S., Cano J.M. & Merilä J. 2006. Genetic relationships among marine and freshwater populations of the European three-spined stickleback (*Gasterosteus aculeatus*) revealed by microsatellites. *Molecular Ecology* 15 (6): 1519–1534. <https://doi.org/10.1111/j.1365-294X.2006.02871.x>

Mäkinen H.S., Cano J.M. & Merilä J. 2008a. Identifying footprints of directional and balancing selection in marine and freshwater three-spined stickleback (*Gasterosteus aculeatus*) populations. *Molecular Ecology* 17 (15): 3565–3582. <https://doi.org/10.1111/j.1365-294X.2008.03714.x>

Mäkinen H.S., Shikano T., Cano J.M. & Merilä J. 2008b. Hitchhiking mapping reveals a candidate genomic region for natural selection in three-spined stickleback chromosome VIII. *Genetics* 178 (1): 453–465. <https://doi.org/10.1534/genetics.107.078782>

Mandić M., Dekić R., Ivanc A., Lolić S., Manojlović M., Friščić J. & Erić Ž. 2016. Leukocyte formula of *Telestes metohiensis* (Steindachner, 1901) from different habitats. *Skup* 7 (2): 129–137.

Marčić Z., Tutman P., Čaleta M., Glamuzina B. & Dulčić J. 2012. Morfometrijska analiza ždrijelnih zuba peškelja, *Scardinius plotizza* Heckel & Kner, 1858 (Pisces, Cyprinidae) sa područja močvare Hutovo blato u

- Bosni i Hercegovini. In: Jelaska S.D., Klobučar G.I.V., Šerić Jelaska L., Leljak Levanić D. & Lukša Ž. (eds) *Zbornik sažetaka 11. Hrvatskog Biološkog Kongresa*: 115. Hrvatsko Biološko društvo, Zagreb.
- Marešová E., Delić A., Kostov V., Marić S., Mendel J. & Šanda R. 2011. Genetic diversity of *Sabanejewia balcanica* (Actinopterygii: Cobitidae) in the western Balkans and comparison with other regions. *Folia Zoologica* 60 (4): 335–342.
- Marić C. 1972. *Hromosomi Vrsta Leuciscus cephalus, Leuciscus turskyi i Leuciscus svallize*. Diploma thesis, University of Sarajevo, Sarajevo.
- Marić D. 1980. Prilog poznavanju rasprostranjenja *Paraphoxinus alepidotus* (Heckel, 1843) u vodama Jugoslavije. *Glasnik Republičkog Zavoda Zaštite Prirode – Prirodnjačkog Muzeja u Titogradu* 13: 101–105.
- Marić D. 1981. Finding of *Rutilus rubilio* (Bonaparte, 1837) and *Scardinius erythrophthalmus* (Linnaeus, 1758) in the underground course Korana river by Bosansko Grahovo. *Glasnik Republičkog Zavoda Zaštite Prirode – Prirodnjačkog Muzeja u Titogradu* 14: 97–102.
- Marić D. 1983. Morfološke karakteristike *Phoxinellus alepidotus* (Heckel, 1843) (Cyprinidae) iz ponornice Korane kod Bosanskog Grahova (Jugoslavija). *Glasnik Republičkog Zavoda Zaštite Prirode – Prirodnjačkog Muzeja u Titogradu* 16: 65–73.
- Marić D. 1988. The species revision of genus *Rutilus* Rafinesque, 1820 (Pisces) from western part of Balkan peninsula. *Glasnik Republičkog Zavoda za Zaštitu Prirode – Prirodnjačkog Muzeja Titograd* 21: 55–79.
- Marić D. 1990. Razmnožavanje *Phoxinellus alepidotus* Heckel, 1843 u podzemnom toku potoka Korana kod Bosanskog Grahova. *Glasnik Republičkog Zavoda za Zaštitu Prirode Prirodnjačkog Muzeja u Titogradu* 23: 75–84.
- Marić D. 2010. *Rutilus albus* sp. n. (Teleostei, Cyprinidae) from Lake Skadar. *Periodicum Biologorum* 112 (2): 153–158.
- Marić D.S. 2019. *Fauna of Freshwater Fish (Osteichthyes) of Montenegro*. Montenegrin Academy of Sciences and Arts, Podgorica.
- Marić A., Špelić I., Radočaj T., Vidović Z., Kanjuh T., Vilizzi L., Piria M., Nikolić V., Škraba Jurlina D., Mrdak D. & Simonović P. 2022. Changing climate may mitigate the invasiveness risk of non-native salmonids in the Danube and Adriatic basins of the Balkan Peninsula (south-eastern Europe). In: Giannetto D., Piria M., Tarkan A.S. & Zięba G. (eds) *Recent Advancements in the Risk Screening of Freshwater and Terrestrial Non-native Species, NeoBiota* 76: 135–161. <https://doi.org/10.3897/neobiota.76.82964>
- Marić S., Razpet A., Nikolić V. & Simonović P. 2011. Genetic differentiation of European grayling (*Thymallus thymallus*) populations in Serbia, based on mitochondrial and nuclear DNA analyses. *Genetics Selection Evolution* 43 (2): 1–11. <https://doi.org/10.1186/1297-9686-43-2>
- Marić S., Kalamujić B., Snoj A., Razpet A., Lukić-Bilela L., Pojskić N. & Sušnik Bajec S. 2012. Genetic variation of European grayling (*Thymallus thymallus*) populations in the Western Balkans. *Hydrobiologia* 691 (1): 225–237. <https://doi.org/10.1007/s10750-012-1076-2>
- Marić S., Snoj A., Sekulić N., Krpo-Ćetković J., Šanda R. & Jojić V. 2015. Genetic and morphological variability of the European mudminnow *Umbra krameri* (Teleostei, Umbridae) in Serbia and in Bosnia and Herzegovina, a basis for future conservation activities. *Journal of Fish Biology* 86 (5): 1534–1548. <https://doi.org/10.1111/jfb.12657>
- Marić S., Stanković D., Wanzenböck J., Šanda R., Erős T., Péter T., Specziár A., Sekulić N., Bănăduc D., Čaleta M., Trombitsky I., Galambos L., Sipos S. & Snoj A. 2017. Phylogeography and population genetics of the European mudminnow (*Umbra krameri*) with a time-calibrated phylogeny for the family Umbridae. *Hydrobiologia* 792: 151–168.

- Marić S., Stanković D., Šanda R., Čaleta M., Čolić S., Šukalo G. & Snoj A. 2019. Genetic characterisation of European mudminnow (*Umbra krameri*) populations from the Sava River system. *Knowledge & Management of Aquatic Ecosystems* 420 (46): 1–9. <https://doi.org/10.1051/kmae/2019035>
- Markotić I., Bartulović V. & Glamuzina B. 2012. Sex ratio of *Phoxinellus pseudalepidotus* Bogutskaya & Zupančić 2003 (Teleostei: Cyprinidae), from Mostarsko blato area. In: Casasús I., Rogošić J., Rosati A., Štoković I. & Gabiña D. (eds) *Animal Farming and Environmental Interactions in the Mediterranean Region*: 201–204. Wageningen Academic Publishers, Wageningen.
- Markotić I., Čaleta M., Mrakovčić M. & Glamuzina B. 2013a. The weight structure of *Phoxinellus pseudalepidotus* (Cyprinidae) from the Mostarsko blato (Neretva River basin Bosnia and Herzegovina). In: Kovačević D. (ed.) *Proceedings of Fourth International Agronomic Symposium "Agrosym 2013" Sarajevo*: 988–993. Faculty of Agriculture, University of East Sarajevo, Sarajevo.
- Markotić I., Mihaljević Z., Bartulović V., Dobrosravić T. Sulić Šprem J. & Glamuzina B. 2013b. Sezonska ishrana prikanca (*Phoxinellus pseudalepidotus* Bogutskaya & Zupančić, 2003) na području Mostarskog blata (Bosna i Hercegovina). In: Marić S. & Lončarić Z. (eds) *Zbornik 48. Hrvatskog i 8. Međunarodnog Simpozija Agronoma*: 652–656. Faculty of Agriculture, University of Josip Juraj Strossmayer in Osijek, Dubrovnik, Hrvatska.
- Markotić I., Čaleta M. & Glamuzina B. 2017. Age structure of the endemic fish *Phoxinellus pseudalepidotus* (Cyprinidae) from Mostarsko Blato (Neretva River Basin Bosnia and Herzegovina). *European International Journal of Science and Technology* 6 (5): 50–56.
- Markotić I., Čaleta M. & Glamuzina B. 2018. Length-weight relationship and condition factor of endemic fish *Phoxinellus pseudalepidotus* (Cyprinidae) from Mostarsko Blato (Neretva River Basin, Bosnia and Herzegovina). *European International Journal of Science and Technology* 7 (9): 44–53.
- Markotić I., Mihaljević Z., Čaleta M. & Glamuzina B. 2019. Feeding Ecology of the Endemic *Phoxinellus pseudalepidotus* (Cyprinidae) from Mostarsko Blato (Neretva River Basin Bosnia and Herzegovina). *European International Journal of Science and Technology* 8: 5–14.
- Marková S., Šanda R., Crivelli A., Shumka S., Wilson I.F., Vukić J., Berrebi P. & Kotlík P. 2010. Nuclear and mitochondrial DNA sequence data reveal the evolutionary history of *Barbus* (Cyprinidae) in the ancient lake systems of the Balkans. *Molecular Phylogenetics and Evolution* 55: 488–500. <https://doi.org/10.1016/j.ympev.2010.01.030>
- Martinović V. 1974. *Neke taksonomske karakteristike Alburnoides bipunctatus Bloch 1872 rijeka Kasindolke i Ljubine*. Diploma thesis, University of Sarajevo, Sarajevo.
- Mateš A. 2004. *Zaneseni Ribar: Zapisi, Doživljaji i Ulovi Jednoga Ribolovnog Zanesenjaka*. J&B, Zagreb.
- Mateš A. 2008. *Zaneseni Ribar – Glavatice i Lipljani*. J&B, Zagreb.
- Matić-Skoko S., Tutman P., Dulčić J., Prusina I., Đodo Ž., Pavličević J. & Glamuzina B. 2011. Growth pattern of the endemic Neretvan roach, *Rutilus basak* (Heckel, 1843) in the Hutovo Blato wetland. *Journal of Applied Ichthyology* 27 (3): 813–819. <https://doi.org/10.1111/j.1439-0426.2010.01593.x>
- McNeely J.A., Miller K.R., Reid W.V. & Mittermeier R. 1990. *Conserving the World's Biological Diversity*. International Union for Conservation of Nature and Natural Resources Institute, Conservation International, Washington, D.C.
- Medić M. 1896. Ihtiološke bilješke. *Rad Jugoslavenske Akademije Znanosti i Umjetnosti* 126: 83–109.
- Medić M. 1898. Bodorka. *Rad Jugoslavenske Akademije Znanosti i Umjetnosti* 135: 199–208.
- Medić M. 1901. Drugo kolo ihtioloških bilježaka. *Rad jugoslavenske Akademije Znanosti i Umjetnosti* 30: 139–191.

- Medić M. 1902. Sabljarka (*Pelecus cultratus* Linné) i Platnica (*Leuciscus virgo* Heckel). *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 1: 113–122.
- Medić-Žuljević M. 1975. *Neke Taksonomske Karakteristike Gobio gobio Linne Rijeka Štira i Jador*. Diploma thesis, University of Sarajevo, Sarajevo.
- Mehanović A. 2007. *Morfološke Osobnosti Digestivnog Trakta Škobalja – Chondrostoma nasus (Linnaeus, 1758) iz Rijeka Bosna, Krivaja i Vrbas*. Diploma thesis, University of Sarajevo, Sarajevo.
- Memić K. 2006. *Morfološko-taksonomske Karakteristike Krkuše (Gobio gobio L.) iz Nekih Tekućica sa Područja Tuzlanske Regije*. Master's thesis, Prirodno-matematički fakultet Univerziteta u Sarajevu, Sarajevo.
- Memić K. 2018. Biodiverzitet tuzlanske regije, biodiverzitet slatkovodnih kolousta i riba. *Baština Sjeveroistočne Bosne Broj* 10: 213–223.
- Memić K. 2020. Dolina rijeke Turije. *Baština Sjeveroistočne Bosne broj* 12: 221–232.
- Memić K. & Adrović A. 2005. Morfometrijske i merističke karakteristike krkuše (*Gobio gobio* L.) iz rijeke Spreče. *Zbornik Radova PMF-a Tuzla II* (2): 103–109.
- Memić K. & Adrović A. 2010. Biodiversity of ichthyofauna of the Spreča river as a part of natural heritage of Tuzla area. In: Popović-Živančević M. & Marina K. (eds) *Condition of the Cultural and Natural Heritage in the Balkan Region – South East Europe 2*: 215–226. Central Institute for Conservation, Institute Goša doo, Ministry of Culture of Republic of Serbia, Belgrade.
- Mihailović M., Blagojević D., Ogrinc N., Simonović P., Simić V., Vidaković V., Dinić S., Uskoković A., Grdović N., Arambašić-Jovanović J., Đorđević M., Tolić A., Kračun-Kolarević M., Kolarević S., Piria M. & Paunović M. 2016. Biochemical indicators and biomarkers in chub (*Squalius cephalus* L.) from the Sava River. *Science of the Total Environment* 540: 368–376. <https://doi.org/10.1016/j.scitotenv.2015.06.098>
- Mihinjač T., Markotić I. & Čaleta M. 2014. Threatened fishes of the World: *Phoxinellus pseudalepidotus* Bogutskaya & Zupančić 2003 (Cyprinidae). *Croatian Journal of Fisheries* 72: 267–270. <https://doi.org/10.14798/72.4.755>
- Mijatović A. 1966. *Ishrana Lipljena u Donjem Toku Rijeke Sane (ljetna sezona)*. Diploma thesis, University of Sarajevo, Sarajevo.
- Mijatović N., Pejić K., Pavlović V., Gvozdenović O., Kekić H. & Ivanc A. 1982. Proteini seruma oštrulja (*Aulopyge hügelii*) u uslovima gladovanja. *Izvodi Saopštenja VI Kongresa Biologa*. Unija bioloških naučnih društava Jugoslavije i Unija bioloških naučnih društava Vojvodine, Novi Sad.
- Mikavica D. 1987a. Dužinski i maseni rast mladice (*Hucho hucho* L.) iz rijeke Drine. *Ribarstvo Jugoslavije* 42: 81–86.
- Mikavica D. 1987b. New data of the composition of ichthyofauna in the hydroaccumulation “Bajina bašta”. *Ribarstvo Jugoslavije* 42 (2–3): 9–12.
- Mikavica D. 1988a. Uticaj otpadnih industrijskih voda na ihtiofaunu rijeke Vrbasa (dio toka na području SO Jajce). *Ribarstvo Jugoslavije* 43 (5–6): 99–101.
- Mikavica D. 1988b. Characteristics of the population of Turskyi dace fish (*Leuciscus turskyi*) (Haeckel 1843) from lake Buško. *Ribarstvo Jugoslavije* 43 (1): 2–7.
- Mikavica D. 1998. The specific freshwater ichthyofauna of the hydrographic region of Bosnia and Herzegovina. *Ichthyologia* 30 (1): 15–30.
- Mikavica D. & Bošnjak Ž. 1989. Morfološke karakteristike digestivnog trakta nekih vrsta riba iz roda *Chondrostoma*. *Ichthyologia* 21 (1): 7–16.

- Mikavica D. & Dizdarević F. 1990. Karakteristike digestivnog trakta nekih vrsta riba iz roda *Chondrostoma*. *Veterinaria* 39 (3–4): 347–357.
- Mikavica D. & Grujić R. 1995a. Rezultati tova kalifornijske pastrmke (*Parasalmo gairdneri* Rich.) u kavezima u hidroakumulaciji Bajina Bašta. Tehnološki parametri uzgoja. *Tehnologija Mesa* 36 (5): 306–308.
- Mikavica D. & Grujić R. 1995b. Rezultati tova kalifornijske pastrmke (*Parasalmo gairdneri* Rich.) u kavezima u hidroakumulaciji Bajina Bašta. Stepen iskorištenja hrane i dnevni prirast ribe. *Tehnologija Mesa* 36 (5): 309–312.
- Mikavica D. & Kosorić Đ. 1986. Uzrasna i polna struktura, dužinsko i težinsko rasteenje podbile (*Chondrostoma phoxinus* Heckel, 1843) iz Buškog blata. *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 39: 71–81.
- Mikavica D. & Savić N. 1997. Growth of chub (*Leuciscus cephalus* L.) and barbel (*Barbus barbus* L.) in the river Drina. In: *Jugoslavenski Simpozijum sa Međunarodnim Učešćem. Ribarstvo Jugoslavije, Stručni Radovi, Saopćenja i Drugi Prilozi iz Ribarstvo*: 81–95. Cetinje – Rijeka, Crnojevića.
- Mikavica D. & Savić N. 1999. *Ribe Rijeke Drine (Drugo Dopunjeno i Prerađeno Izdanje)*. Poljoprivredni fakultet, Banja Luka.
- Mikavica D. & Vuković D. 2003. Morphologic variations in the kidney of the Carp (*Cyprinus carpio* L., 1758) from the Saničani Fishpond – Prijedor. *Ichthyologia* 1: 23–32.
- Mikavica D., Sofradžija A. & Škrijelj R. 1988. Rasprostranjenje i neke idioekološke karakteristike lipljena (*Thymallus thymallus* L.) iz rijeke Drine. *Ichthyologia* 20 (1): 27–36.
- Mikavica D., Sofradžija A. & Bošnjak Z. 1989. Idioekološke karakteristike skobalja (*Chondrostoma nasus* L.) iz rijeke Drine. In: *Zbornik radova UPI-institut*: 109–116. Sarajevo.
- Mikavica D., Sofradžija A. & Dizdarević F. 1991. Quality and quantity of the fish population in the Drina River. *Veterinaria* 40 (1–2): 133–141.
- Mikavica D., Grujić R. & Komić J. 1997. Comparative growth analysis of the nase *Chondrostoma nasus* L. 1758, chub *Leuciscus cephalus* L. 1758 and barbel *Barbus barbus* L. 1758 in the river Drina. *Ichthyologia* 29 (1): 1–17.
- Mikavica D., Muhamedagić S., Dizdarević F. & Savić N. 2001. Idioecological characteristics of Adriatic trout *Salmothymus obtusirostris oxyrhynchus* (Steindachner, 1882). In: *Symposium of Livestock Production International Participation*: 23–25. Struga.
- Mikavica D., Muhamedagić S., Dizdarević F. & Savić N. 2002. Idioecological characteristics of Adriatic trout – *Salmothymus obtusirostris oxyrhynchus* (Steindachner, 1882). In: Opačak A. (ed.) *Proceedings of the 3rd National Scientific and Professional Conference with International Participation “Fresh-water Fishery of the Republic of Croatia in the Period of Association with European Union”*: 16. The Croatian Chamber of Economy, Bizovac.
- Mikavica D., Golub D., Šukalo G., Dmitrović D., Cvijić S. & Šukalo M. 2013. Morphometric and meristic characters of Brown bullhead *Ameiurus nebulosus* (Le Sueur, 1819) on wetland ecosystem Bardača. *Skup* 5 (1): 23–28.
- Miladinović Ž., Guzina N., Vuković T. & Selimić E. 1971. Aglutinacija eritrocita *Paraphoxinus alepidotus*, *P. metohiensis* i *P. adpersus* serumima pastrmki, toplokrvnih životinja i čovjeka. *Ichthyologia* 3 (1): 49–55.
- Milanović Đ., Stupar V., Kulijer D., Kotrošan D. & Hamzić A. 2015. Natura 2000 in Bosnia and Herzegovina: where are we at the moment? *Bulletin of the Faculty of Forestry, University of Banja Luka* 23: 95–134. <https://doi.org/10.7251/GSF1523095M>

- Mišić S. 2011. *Komparativni prikaz ishrane potočne pastrmke Salmo trutta (Linnaeus, 1758) i lipljena Thymallus thymallus (Linnaeus, 1758) iz sliva rijeke Sane u općini Ključ*. Diploma thesis, University of Sarajevo, Sarajevo.
- Mitrašinović M. 2002. *Ishrana lipljena (Thymallus thymallus L.) u rijeci Uni*. Diploma thesis, University of Sarajevo, Sarajevo.
- Mitrašinović M. & Suljević D. 2009. Hematological status of Chub fish *Leuciscus cephalus* (Linnaeus, 1758.) from Krupica and Željeznica rivers. *Veterinaria* 58 (1–2): 63–76.
- Mitrašinović-Brulić M., Škrijelj R. & Šuta N. 2016. Hematological status of zander *Stizostedion lucioperca* Linnaeus, 1758 from hydro reservoirs Modrac in seasonal aspect. *Works of the Faculty of Agriculture University of Sarajevo* 61 (66/2): 131–143.
- Mojsisovics A. 1897. *Das Tierleben der österreichisch-ungarischen Tiefebene. Biologische und thiergeographische Skizzen und Bilder*. Hölder, Wien.
- Mrakovčić M., Mišetić S. & Povž M. 1995. Status of freshwater fish in Croatian Adriatic river systems. *Biological Conservation* 72: 179–185. [https://doi.org/10.1016/0006-3207\(94\)00080-A](https://doi.org/10.1016/0006-3207(94)00080-A)
- Mrakovčić M., Kerovec M., Mišetić S. & Schneider D. 1996. Description of *Knipowitschia punctatissima croatica*, (Pisces: Gobiidae), a new freshwater goby from Dalmatia, Croatia. In: Kirchhofer A. & Hefti D. (eds) *Conservation of Endangered Freshwater Fish in Europe*: 311–319. Birkhauser Verlag, Basel.
- Mrakovčić M., Schneider D., Mustafić P. & Kerovec M. 2000. Status of genus *Cobitis* and related species in Croatia. *Folia Zoologica* 49 (Suppl. 1): 113–116.
- Mrakovčić M., Mustafić P., Čaleta M., Zanella D. & Radić I. 2002. Značajke ribljih zajednica Parka prirode Lonjsko polje. *Bilten Parka Prirode Lonjsko Polje* 4: 1–56.
- Mrakovčić M., Brigić A., Buj I., Čaleta M., Mustafić P. & Zanella D. 2006. *Red Book of Freshwater Fish of Croatia*. Ministry of culture, state institute for nature protection, Republic of Croatia.
- Mrakovčić M., Duplić A., Mustafić P. & Marčić Z. 2008. Conservation status of the genus *Cobitis* and related genera in Croatia. *Folia Zoologica* 57 (1–2): 35–41.
- Mrdak D. & Milošević D. 2017. Length-weight relationship of nine fish species from Bosnia and Herzegovina. *Agriculture & Forestry* 63 (2): 157–160. <https://doi.org/10.17707/AgricultForest.63.2.13>
- Mršić V. 1935. *Iskustva sa Udomaćenjem Dužičaste Pastrve u Jugoslaviji*. Tisak Zaklade Narodnih Novina, Zagreb.
- Muhamedagić S. & Habibović E. 2013. The state and perspective of Danube huchen (*Hucho hucho*) in Bosnia and Herzegovina. *Archives of Polish Fisheries* 21: 155–160. <https://doi.org/10.2478/aopf-2013-0012>
- Muhamedagić S., Gjoen H.M. & Vegara M. 2008. Salmonids of the Neretva river basin – present state and suggested sustainable selection programme to protect and strengthen salmonid populations. In: Cowx I.G, Taylor R., Walton S., Angelopoulos N., Smith M., Twine K., Davies C. & van Anrooy R. (eds) *Proceedings of the EIFAC Symposium on Interactions Between Social, Economic and Ecological Objectives on Inland Commercial and Recreational Fisheries and Aquaculture*: 224–275. Food and Agriculture Organization, Antalya.
- Muhamedagić S., Korjenić E., Hamzić A., Drešković N., Mustafić A., Kapo I., Lepara D. & Habibović E. 2013. Ichthyofauna of Krivaja River catchment area. In: Poleksić V. (ed.) *Conference Proceedings from VI International Conference “Water & Fish”*: 310–321. University of Belgrade, Faculty of Agriculture, Belgrade.

- Muhamedagić S., Pavličević J. & Savić N. 2019. Composition of ichthyopopulations in the upper stream of the Neretva River, half a century after the construction of the first dam. *Works of the Faculty of Agriculture and Food Sciences, University of Sarajevo* 69 (1): 134–145.
- Muhanović E. 1984. *Morfološko-taksonomske Karakteristike Ukljeve (Alburnus alburnus albarella (De Filippi, 1844) iz Hutovog Blata*. Diploma thesis, University of Sarajevo, Sarajevo.
- Munjko I. 1972. O pomoru riba u rijeci Savi. *Priroda* 7: 205–206.
- Mustafić P. & Mrakovčić M. 2017. Biodiversity status of and threats to freshwater fish of Croatia and Eastern Adriatic Countries. In: Berillis P. (eds) *Trends in Fisheries and Aquatic Animal Health*: 23–41. Bentham Science Publisher, Sharjah.
- Mustafić P., Čaleta M., Mrakovčić M., Buj I., Zanella D. & Mišetić S. 2005. Distribution and status of the genus *Gobio* in Croatia. *Folia Zoologica* 54: 81–84.
- Mustafić P., Čaleta M., Marčić Z., Zanella D. & Mrakovčić M. 2015. *Ichthyological Exploration on the Impacts of the Future HPP Janjići Hydropower Plant with the Aim of Preserving the Ichthyofauna of the Bosna River*. Croatian Ichthyological Society, Croatia.
- Mušović A. 2016. *Ekološka, Morfološka i Molekularno-genetička Karakterizacija Vrste Aulopyge huegelii Heckel, 1843 (Actinopterygii: Cipriniformes) iz Buškog Jezera*. PhD thesis, University of Sarajevo, Sarajevo.
- Mušović A., Đug S., Pojskić N., Kalamujić Stroil B., Vesnić A. & Škrijelj R. 2018. Status of endangered fish species *Aulopyge huegelii* Heckel, 1843 (Teleostei: Cyprinidae) in the Buško Blato reservoir, Bosnia and Herzegovina. *Iranian Journal of Ichthyology* 5 (3): 212–231. <https://doi.org/10.22034/iji.v5i3.290>
- Mušović A., Škrijelj R., Gajević M., Kalamujić Stroil B., Vesnić A., Mitrašinović-Brulić M. & Đug S. 2020. Long-term trends in the structure and dynamics of the fish communities in Buško blato Reservoir. *Croatian Journal of Fisheries* 78: 69–78. <https://doi.org/10.2478/cjf-2020-0007>
- Mušović A., Smailagić Vesnić L., Šljuka S., Vesnić A., Gajević A., Pilić S. & Đug S. 2025. Native distribution, introductions and risk screening of *Sander lucioperca* (Linnaeus, 1758) in the freshwater ecosystems of Bosnia and Herzegovina. *Croatian Journal of Fisheries* 83: 1–8.
- Nadaždin M., Grbelja J., Vuković T., Juretić N. & Guzina N. 1980. Imuni odgovor riba na nukleoprotein Turnip Yellow mosaic virusa. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 33: 147–153.
- Naseka A.M., Bogutskaya N.G. & Bănărescu P.M. 1999. *Gobio albipinnatus* Lukasz, 1933. In: Bănărescu P.M. (eds) *The Freshwater Fishes of Europe. Cyprinidae 2/I. Vol. 5/I*: 37–68. AULA-Verlag, Wiebelsheim.
- Nalbant T.T., Ráb P., Bohlen J. & Saitoh K. 2001. Evolutionary success of the loaches of the genus *Cobitis* (Pisces: Ostariophysi: Cobitidae). *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"* 43: 277–289.
- Nedić Z., Begović S., Dogan M., Hadžiahmetović Jurida E., Ferizbegović J. & Terzić R. 2014a. Decreasing of biodiversity of fish populations from the Sava River in Bosnia and Herzegovina. *Basic Research Journal of Agricultural Science and Review* 3 (5): 35–40.
- Nedić Z., Riđanović S. & Halilović E. 2014b. Ectoparasitic trematodes on *Scardinius erythrophthalmus* from the lower flow of the Sava River. *Veterinaria* 64 (1–4): 37–44.
- Nedić Z., Skenderović I. & Riđanović S. 2014c. Skin ectoparasites of fish from the lower flow of the Sava River. *Veterinaria* 63 (1–4): 45–53.

- Nedić Z., Halilović E., Terzić R., Skenderović I., Adrović A., Hadžiahmetović Jurida E., Živković M. & Ferizbegović J. 2014d. Ectoparasitic crustaceans are not responsible for decreasing of fish population from the Sava River. *Basic Research Journal of Agricultural Science and Review* 3 (12): 161–166.
- Nedić Z., Djikanović V. & Nikolić V. 2016a. First record of *Eudiplozoon nipponicum*, Goto 1891 (Monogenea) in the Sava River, Bosnia and Herzegovina. In: Melovski Lj. (ed.) *5th Congress of Ecologists of Macedonia with International Participation*. Macedonian Ecological Society and Institute of Biology, Faculty of Natural Sciences, Skopje, Ohrid, Macedonia.
- Nedić Z., Riđanović S. & Nikolić V. 2016b. First record of ectoparasite community on fishes in lower flow of Tolisa river, Bosnia and Herzegovina. *Educa, Journal for education, Science and Culture* 9: 45–52.
- Nedić Z., Skenderović I. & Adrović A. 2018. Study of some ectoparasites of fishes from the Sava River as part of water management in Bosnia and Herzegovina. *TEM Journal* 7 (2): 391–397. <https://doi.org/10.18421/TEM72-21>
- Nedić Z., Simonović P. & Nikolić V. 2021. A note on the occurrence of *Pallisentis ussuriensis* (Kostylev, 1941) (Acanthocephala) from the brown bullhead (*Ameiurus nebulosus*) in Bosnia and Herzegovina. In: Klun I. & Djurković-Djaković O. (eds) *Book of Abstracts of the 13th European Multicollloquium of Parasitology*: 206–207. Serbian Society for Parasitology, Belgrade.
- Neilson M.E. & Stepien C.A. 2009a. Evolution and phylogeography of the tubenose goby genus *Proterorhinus* (Gobiidae: Teleostei): Evidence for new cryptic species. *Biological Journal of Linnean Society* 96: 664–684. <https://doi.org/10.1111/j.1095-8312.2008.01135.x>
- Neilson M.E. & Stepien C.A. 2009b. Escape from the Ponto-Caspian: Evolution and biogeography of an endemic goby species flock (Benthophilinae: Gobiidae: Teleostei). *Molecular Phylogenetics and Evolution* 52: 84–102. <https://doi.org/10.1016/j.ympev.2008.12.023>
- Neuburg J., Friedrich T., Haring E., Höfler S., Maric A., Marić S., Popp S., Simonović P., Snoj A., Sušnik Bajec S., Vitecek S., Wanka A., Weiss S. & Pinter K. 2023. The ichthyofauna of the upper Neretva River. *Natura Sloveniae* 25 (3):155–180. <https://doi.org/10.14720/ns.25.3.155-180>
- Nikolić V., Bilbija B., Nedić Z., Simonović P. & Djikanović V. 2018. First record of *Azygia robusta* (Odhner, 1911) (Trematoda: Digenea: Azygiidae) in Brown trout (*Salmo trutta*) in the Vrbas River. *Croatian Journal of Fisheries* 76 (2): 85–88. <https://doi.org/10.2478/cjf-2018-0011>
- Nišić E. 2008. *Morfoanatomske Karakteristike Digestivnog Trakta Vrste Leuciscus cephalus (Linnaeus, 1758) iz Rijeka Bosne i Jošanice*. Diploma thesis, University of Sarajevo, Sarajevo.
- Nuhanović F. 1989. *Ispitivanje Plodnosti Vrste Aulopyge hugeli (Heckel, 1841) (Cyprinidae) iz Buškog Jezera*. Diploma thesis, University of Sarajevo, Sarajevo.
- Obradović S. 1974. *Hromosomi Američkog Somića (Ictalurus nebulosus)*. Diploma thesis, University of Sarajevo, Sarajevo.
- Obratil S. 1982. Ekološki pristup utvrđivanju štetnosti ihtiofagnih ptica u ribnjacima Bardača. *Glasnik Zemaljskog Muzeja BiH* 19–20: 139–256.
- Official Gazette of the Federation of Bosnia and Herzegovina no 2020. Rules on protection measures for strictly protected species and subspecies and protected species and subspecies. Available from <https://fmoit.gov.ba/dokument/pravilnik-o-mjerama-zastite-za-strogo-zasticene-i-zasticene-vrste-i-podvrste-i-zasticene-vrste-i-podvrste/>
- Official Gazette of the Republic of Srpska no 65 2020. Regulation on strictly protected and protected wild species.

- Oikonomou A., Leprieur F. & Leonardos I.D. 2014. Biogeography of freshwater fishes of the Balkan Peninsula. *Hydrobiologia* 738: 205–220. <https://doi.org/10.1007/s10750-014-1930-5>
- Okuka D. 1966. *Morfološke Karakteristike Vrste Cottus gobio Linne, 1758*. Diploma thesis, University of Sarajevo, Sarajevo.
- Omeragić N., Pocrnjić Z. & Vuković T. 1980. Toplotna rezistencija skeletnih mišića potočne mreke (*Barbus meridionalis petenyi*). *Ichthyologia* 12 (1): 33–42.
- Orelj Z. 1979. *Morfološko-taksonomske Karakteristike Uklije Alburnus alburnus alburnus L. 1758 iz Gornjeg toka Rijeke Drine*. Diploma thesis, University of Sarajevo, Sarajevo.
- Ovčina A. 2007. *Biosistematske Karakteristike Vrste Cobitis elongatoides (Bacescu & Maier, 1969) iz Rijeke Bosne – Ušća u Savu*. Diploma thesis, University of Sarajevo, Sarajevo.
- Palandačić A., Zupančić P. & Snoj A. 2010. Revised classification of former genus *Phoxinellus* using nuclear DNA sequences. *Biochemical Systematics and Ecology* 38: 1069–1073. <https://doi.org/10.1016/j.bse.2010.09.021>
- Palandačić A., Matschiner M., Zupančić P. & Snoj A. 2012. Fish migrate underground: the example of *Delminichthys adspersus* (Cyprinidae). *Molecular Ecology* 21: 1658–1671. <https://doi.org/10.1111/j.1365-294X.2012.05507.x>
- Palandačić A., Bravničar J., Zupančić P., Šanda R. & Snoj A. 2015. Molecular data suggest a multispecies complex of *Phoxinus* (Cyprinidae) in the Western Balkan Peninsula. *Molecular Phylogenetics and Evolution* 92: 118–123. <https://doi.org/10.1016/j.ympev.2015.05.024>
- Palandačić A., Naseka A., Ramler D. & Ahnelt A. 2017. Contrasting morphology with molecular data: an approach to revision of species complexes based on the example of European *Phoxinus* (Cyprinidae). *BMC Evol Biol* 17: 184. <https://doi.org/10.1186/s12862-017-1032-x>
- Palandačić A., Kruckenhauser L., Ahnelt H. & Mikschi E. 2020. European minnows through time: museum collections aid genetic assessment of species introductions in freshwater fishes Cyprinidae: *Phoxinus* species complex). *Heredity* 124 (3): 410–422. <https://doi.org/10.1038/s41437-019-0292-1>
- Palandačić A., Reier S., Diripasko O.A., Jelić D., Stroj A., Wanka A., Marić D. & Bogutskaya N.G. 2024. Substygophily in Dinaric Karst: A Model Case of Locally Endemic Minnows *Phoxinellus* (Leuciscinae). *Ecology and Evolution* 14: e70648. <https://doi.org/10.1002/ece3.70648>
- Paspalj J., Pljevaljčić T. & Dekić R. 2018. Morphometric characteristics of erythrocytes of Common gudgeon (*Gobio obtusirostris*). *Skup* 9 (2): 17–24. <https://doi.org/10.7251/SKP180902017P>
- Pavić P. 2008. *Morfometrijske Mjere Turskog Klena (Telestes turskyi)*. Diploma thesis, University of Mostar, Mostar.
- Pavličević J., Petrović D., Bogut I., Stanić-Koštroman S., Dedić A. & Batinić V. 2014. *Revizija Ribolovne Osnove za Ribolovnu zonu Srednjobosanskog kantona*. Agronomski i prehrambeno-tehnološki fakultet Sveučilišta u Mostaru, Zavod za ribarstvo, zoologiju i zaštitu voda, Mostar, Bosna i Hercegovina.
- Pavličević J., Glamuzina L., Conides A., Savić N., Rozić I., Klaoudatos D., Kazić A. & Glamuzina B. 2016. Pikeperch, *Sander lucioperca* invasion in the Neretva River watershed (Bosnia and Herzegovina, Croatia) after alteration of river flow. *River Research and Applications* 32: 967–974. <https://doi.org/10.1002/rra.2923>
- Pavlović B. & Berberović Lj. 1978. Kariometrijski pristup proučavanju citotaksonomije nekih vrsta roda *Paraphoxinus* (Cyprinidae, Pisces). *Ichthyologia* 10 (1): 85–113.
- Pavlović B.P., Pavlović N., Vidović V., Vuković D., Dekić R., Filipović S., Paraš, S., Pajčin R., Šukalo G. & Dmitrović D. 2009. Endemic species and subspecies of animals important for the preservation of biotic diversity of Republic Srpska. In: Predrag I. (ed.) *Collection of papers from the Scientific-Professional*

conference with international participation „Safety and Health in Work and Environmental Protection, 24–26 June: 501–517. Institute of Protection, Ecology and Informatics, Banja Luka.

Pavlović V., Vuković T., Mladenović O. & Kekić H. 1960. Prilog poznavanju sedimentacije eritrocita, broja eritrocita i koncentracije hemoglobina nekih vrsta riba sliva reke Bosne. *Godišnjak Biološkog instituta Univerziteta u Sarajevu* 13 (1–2): 107–115.

Pavlović V., Mladenović O., Kekić H. & Vuković T. 1962. Sedimentacija eritrocita, broj eritrocita i koncentracija hemoglobina potočne pastrmke (*Salmo trutta m. fario* L.) i lipljena (*Thymallus thymallus* L.) iz izvorišnog toka reke Bosne u sezonskim i ekološkim uslovima. *Godišnjak Biološkog instituta Univerziteta u Sarajevu* 15 (1–2): 55–76.

Pažur K. 1993. Ribarenje na Uni na području Kostajnice. *Ribarstvo* 3: 111–112.

Pažur K., Habeković D. & Popović J. 1982. Dinamika rasta mladice (*Hucho hucho* L. 1758) u vodama SR Hrvatske. *Ichthyologia* 14 (2): 161–169.

Pejić K., Gvozdrenović O., Mijatović N., Pavlović V., Kekić H., Hadžiselimović R., Sofradžija A. & Ivanc A. 1980. Sezonske promjene relativnih aktivnosti izoenzima malat dehidrogenaze (1.1.1.37) iz ekstrakta mišića zelenike (*Alburnus alburnus alburnus*). *Ichthyologia* 12 (1): 43–50.

Petrosino G., Tancioni L., Turani M., Rakaj A., Ciuffardi L. & Rossi A.R. 2022. Phylogeography of *Sarmarutilus rubilio* (Cypriniformes: Leuciscidae): complex genetic structure, clues to a new cryptic species and further insights into roaches phylogeny. *Genes* 13: 1071. <https://doi.org/10.3390/genes13061071>

Perea S., Böhme M., Zupančič P., Freyhof J., Šanda R., Özüluğ M., Abdoli A. & Doadrio I. 2010. Phylogenetic relationships and biogeographical patterns in Circum-Mediterranean subfamily Leuciscinae (Teleostei, Cyprinidae) inferred from both mitochondrial and nuclear data. *BMC Evolutionary Biology* 10: 265. <https://doi.org/10.1186/1471-2148-10-265>

Pereira A.M., Levy A., Vukić J., Šanda R., Levin B.A., Freyhof J., Geiger M., Choleva L., Francisco S.M. & Robalo J.I. 2021. Putting European lampreys into perspective: A global-scale multilocus phylogeny with a proposal for a generic structure of the Petromyzontidae. *Journal of Zoological Systematics and Evolutionary Research* 59: 1982–1993. <https://doi.org/10.1111/jzs.12522>

Pešić P. 1987. *Mogućnosti Ispitivanja Starosti Jedinki Vrste Aulopyge hugeli Heckel, 1841*. Diploma thesis, University of Sarajevo, Sarajevo.

Pietroock M., Ritterbusch D., Lewin W.C., Shumka S., Spirkovski Z., Ilik-Boeva D., Brämick U. & Peveling R. 2022. The fish community of the ancient Prespa Lake (Southeast Europe): Non-indigenous species take over. *Fisheries & Aquatic Life* 30: 112–124. <https://doi.org/10.2478/aopf-2022-0011>

Pikula P. 1977. *Hromosomi Karaša (Carrasius carassius) i Srebrenog Karaša (Carrasius auratus gibelio)*. Diploma thesis, University of Sarajevo, Sarajevo.

Pilić P., Mušović A., Gajević M., Škrijelj Š., Đug S. & Vesnić A. 2020. Inter-population variability in morphology and reproduction of *Alburnus alburnus* (Linnaeus, 1758) from Bosnia and Herzegovina. *Genetics & Applications* 4 (1): 18–29. <https://doi.org/10.31383/ga.vol4iss1pp18-29>

Pipić Š. 1966. *Variranje Dužine i Širine Otolita kod Oštrulja i Turskog Klana*. Diploma thesis, University of Sarajevo, Sarajevo.

Piria M., Šprem N., Jakovlić I., Tomljanović T., Matulić D., Treer T., Aničić I. & Safner R. 2011a. First record of round goby, *Neogobius melanostomus* (Pallas, 1814) in the River Sava, Croatia. *Aquatic Invasions* 6 (1): 153–157. <https://doi.org/10.3391/ai.2011.6.S1.034>

- Piria M., Treer T., Tomljanović T., Šprem N., Matulić D., Aničić I. & Safner R. 2011b. First record of monkey goby, *Neogobius fluviatilis* (Pallas, 1814) in the barbel zone of the River Sava, Croatia. *Journal of Applied Ichthyology* 27: 1383–1384. <https://doi.org/10.1111/j.1439-0426.2011.01800.x>
- Piria M., Jakšić G., Jakovlić I. & Treer T. 2016a. Dietary habits of invasive Ponto-Caspian gobies in the Croatian part of the Danube River basin and their potential impact on benthic fish communities. *Science of the Total Environment* 540: 386–395. <https://doi.org/10.1016/j.scitotenv.2015.05.125>
- Piria M., Tomljanović T., Treer T., Safner R., Aničić I., Matulić D. & Vilizzi L. 2016b. The common carp *Cyprinus carpio* in Croatia (Danube and Adriatic basins): a historical review. *Aquaculture International* 24 (6): 1527–1541. <https://doi.org/10.1007/s10499-016-0029-6>
- Piria M., Simonović P., Kalogianni E., Vardakas L., Koutsikos N., Zanella D., Ristovska M., Apostolou A., Adrović A., Mrdak D., Tarkan A.S., Milošević D., Zanella L.N., Bakiu R., Ekmekçi F.G., Povž M., Korro K., Nikolić V., Škrijelj R., Kostov V., Gregori A., Joy M.K. 2018. Alien freshwater fish species in the Balkans – vectors and pathways of introduction. *Fish and Fisheries* 19 (1): 138–169. <https://doi.org/10.1111/faf.12242>
- Plančić J. 1923a. Ribe Save. *Lovačko-ribarski Vjesnik* 32: 74–79.
- Plančić J. 1923b. Ribe Save. *Lovačko-ribarski Vjesnik* 32: 90–99.
- Plančić J. 1967. Srebrnasti karas-*Carassius auratus gibelio* Bloch nova vrsta naše ihtiofaune. *Ribarstvo Jugoslavije* 22: 155–156.
- Pleše S. & Buj I. 2024. Tackling the phylogeny of lampreys—insight from the Croatia’s Danube Basin. *Water* 16: 1153. <https://doi.org/10.3390/w16081153>
- Pocrnjić Z. & Šolaja M. 1988. Thyroid activity in relation to temperature adaptation and heat resistance in the chub (*Leuciscus cephalus* L., 1758). *Ichthyologia* 20 (1): 37–45.
- Pocrnjić Z., Kapetanović N. & Vuković T. 1979. Toplotna rezistencija mišićnog tkiva vrsta roda *Paraphoxinus*. *Ichthyologia* 11 (1): 19–24.
- Pocrnjić Z., Šolaja M., Zubović I. & Skrobić M. 1984. Uticaj aktivnosti tireoideje na kompenzaciju potrošnje kiseonika klena (*Leuciscus cephalus*). *Ichthyologia* 16 (1–2): 103–109.
- Pojškić N. & Kalamujić B. 2015. Simulations based on molecular genetic data in detection of expansion *Salmo trutta* allochthonous population in the Neretva River’s tributaries. In: Visconti P., Game E., Mathevet R. & Wilkerson M. (eds) *Proceedings of the 27th International Congress for Conservation Biology and 4th European Congress for Conservation Biology*: 539–540. ICCB/ECCB, Montpellier.
- Pojškić N., Durmić-Pašić A., Škrijelj R. & Hadžiselimović R. 2007. Brown trout (*Salmo trutta* m. fario l.) population structure in Sava watershed inferred from nuclear and mitochondrial dna markers. *Book of Abstracts of the 12th European Congress of Ichthyology ECI XII* 13: 32. Faculty of Science, University of Zagreb, Cavtat (Dubrovnik).
- Popović J. 2010. *Gospodarenje Ribolovnim Vodama*. Hrvatski športsko ribolovni savez, Zagreb.
- Povž M., Leiner S., Mrakovčić M. & Popović J. 1990. Rare and endangered fishes from Yugoslavian Adriatic rivers. *Journal of Fish Biology* 37 (Suppl. A): 247–249. <https://doi.org/10.1111/j.1095-8649.1990.tb05052.x>
- Povž M., Gregori A. & Gregori M. 2015. *Statkovodne Ribe in Piškurji v Sloveniji*. Zavod Umbra, Ljubljana.
- Pranjić V. 1969. *Hromozomska Garnitura Sitnoljuskavog Klena (Leuciscus turskii)*. Diploma thesis, University of Sarajevo, Sarajevo.
- Proha M. 1997. *Biosistematika i Idioekologija Sapače (Barbus meridionalis petenii Heckel 1847) iz Nekih Pritoka Gornjeg Toka Rijeke Bosne*. PhD thesis, University of Sarajevo, Sarajevo.

- Protić Đ. 1924. Hydrobiological and plankton study on lakes in Bosnia and Herzegovina. *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 36: 39–68.
- Protić Đ. 1933. Ispitivanja o biološkom samočišćenju reke Miljacke (kod Sarajeva). *Glasnik Zemaljskog Muzeja u Bosni i Hercegovini* 45: 41–56.
- Prusina I., Tutman P. & Glamuzina B. 2009. Morphological and meristical properties of endemic Neretvan rudd, *Scardinius plotizza* Heckel & Kner, 1858 (Actinopterygii, Cyprinidae) from the Hutovo Blato wetland, Neretva River basin, Bosnia and Herzegovina. In: Kontautas A. (ed.) *Proceedings of the 13th European Congress of Ichthyology*: 95–96. Klaipėdos Universitetas, Klaipėda.
- Pustovrh G., Sušnik Bajec S. & Snoj A. 2011. Evolutionary relationship between marble trout of the northern and the southern Adriatic basin. *Molecular Phylogenetics and Evolution* 59 (3): 761–766. <https://doi.org/10.1016/j.ympev.2011.03.024>
- Pustovrh G., Snoj A. & Sušnik Bajec S. 2014a. Molecular phylogeny of *Salmo* of the western Balkans, based on multiple nuclear loci. *Genetics Selection Evolution* 46 : e7. <https://doi.org/10.1186/1297-9686-46-7>
- Pustovrh G., Snoj A. & Bajec S.S. 2014b. Correction: Molecular phylogeny of *Salmo* of the western Balkans, based upon multiple nuclear loci. *Genetics Selection Evolution* 46: e12. <https://doi.org/10.1186/1297-9686-46-21>
- Radević M. 1999. Growth of the huchen (*Hucho hucho* L.) from the Vrbas River. *Ichthyologia* 31 (1): 53–59.
- Radević M. 2000. *Ekološki i Cenotički Odnosi Faune Riba u Srednjem i Donjem toku Vrbasa i Ribnjaku Bardači*. Monografija, Prirodno-matematički fakultet, Banja Luka.
- Radević M. 2001. Diverzitet ihtiofaune srednjeg i donjeg toka rijeke Vrbas. *Savremena poljoprivreda* 50 (3–4): 365–369.
- Radević M. 2002. Rast i razmnožavanje nekih ciprinidnih vrsta riba u srednjem i donjem toku reke Vrbas. *Zbornik Prirodno-matematičkih Nauka* 2 (2–3): 129–161.
- Ratković V. & Mikavica D. 1984. On nutrition of some Cyprinidae and zoobenthos in Jablanica Lake. *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 37: 111–121.
- Razpet A., Sušnik S., Jug T. & Snoj A. 2007. Genetic variation among trout in the River Neretva basin, Bosnia and Herzegovina. *Journal of Fish Biology* 70: 94–110. <https://doi.org/10.1111/j.1095-8649.2007.01392.x>
- Reier S., Bogutskaya N. & Palandačić A. 2022. Comparative Phylogeography of *Phoxinus*, *Delminichthys*, *Phoxinellus* and *Telestes* in Dinaric Karst: which factors have influenced their current distributions? *Diversity* 14 (7): 526. <https://doi.org/10.3390/d14070526>
- Ravnateljstvo Hrvatskog Zemaljskog Zoologiškog Muzeja (RHZZM) 1908. Popis riba koje su prispjele hrvatskom Zemaljskom Zoologiškom muzeju od g. 1901. do konca g. 1905. *Glasnik Hrvatskog Naravoslovnog Društva* 20: 114–126.
- Ribeiro F. & Leunda P. 2012. Non-native fish impacts on Mediterranean freshwater ecosystems: Current knowledge and research needs. *Fisheries Management and Ecology* 19: 142–156. <https://doi.org/10.1111/j.1365-2400.2011.00842.x>
- Ribić E. 1901. Ribarstvo i ribogojstvo u Bosni i Hercegovini. *Lovačko-ribarski Viestnik* 10: 97–98.
- Riđanović S., Nedić Z. & Riđanović L 2015. First observation of fish condition from Sava river in Bosnia and Herzegovina. *Journal of Survey in Fisheries Sciences* 1 (2): 27–32.
- Ristić M.Đ. 1963. Ribarsko biološka studija Acipenseridae-a jugoslavenskog dela donjeg Dunava. *Ribarstvo Jugoslavije* 3: 73–80.

- Ristić M.Đ. 1972. Biologija, uzrasna struktura, rastenje i intenzitet ribolova deverike (*Abramis brama* L.) u rekama crnomorskog sliva. *Ribarstvo Jugoslavije* 1: 1–13.
- Ristić M. 1977. *Ribe i Ribolov u Slatkim Vodama*. Nolit, Beograd.
- Robalo J.I., Almada V.C., Levy A. & Doadrio I. 2007. Re-examination and phylogeny of the genus *Chondrostoma* based on the mitochondrial and nuclear data and the definition of 5 new genera. *Molecular Phylogenetics and Evolution* 42: 362–372. <https://doi.org/10.1016/j.ympev.2006.07.003>
- Rojnik Lj. 1969. *Sezonalna Ishrana Potočne Pastrmke (Salmo trutta m. fario L.) iz Sliva Rijeke Gostovića*. Diploma thesis, University of Sarajevo, Sarajevo.
- Rončević G. 1997. *Biosistematske Karakteristike Leuciscus cephalus cephalus (L. 1758) iz Rijeke Čehotine (Akumulacija "Otilovići")*. Master's thesis, University of Sarajevo, Sarajevo.
- Rukavina J. & Ćorić B. 1968. Revizija položaja *Pseudorhadinorhynchus salmothymi* (Šinžar 1956) u redu Polymorphida Petročenko 1956. *Veterinaria* 17.
- Rukavina J. & Delić S. 1957. Prilog poznavanju raširenosti *Azygia lucii* (Müller, 1776) Luhe 1909, kod riba Bosne i Hercegovine. *Veterinaria*, Sarajevo 6 (2–3): 1–4.
- Rukavina J. & Delić S. 1959. *Cyatocephalus truncatus* kod riba u nekim vodama Bosne i Hercegovine. *Veterinaria* 5 (3–4): 547.
- Rüber L., Gandolfi A., Foresti D., Paltrinieri L., Splendiani A. & Seehausen O. 2023. Phylogenetic and biogeographic history of brook lampreys (Lampetra: Petromyzontidae) in the river basins of the Adriatic Sea based on DNA barcode data. *Ecology and Evolution* 13 (9): e10496. <https://doi.org/10.1002/ece3.10496>
- Rylková K., Kalous L., Bohlen J., Lamatsch D.K. & Petrýl M. 2013. Phylogeny and biogeographic history of the cyprinid fish genus *Carassius* (Teleostei: Cyprinidae) with focus on natural and anthropogenic arrivals in Europe. *Aquaculture* 380–383: 13–20. <https://doi.org/10.1016/j.aquaculture.2012.11.027>
- Sabioncello I. 1967. Sistematika slatkovodnih riba. In: Livojević Z. & Bojčić C. (eds) Priručnik za slatkovodno ribarstvo. *Posebna Izdanja Agronomskog Glasnika*: 21–90.
- Sabolić M., Buj I., Čaleta M. & Marčić Z. 2021. Morphological diversity and relationships of *Scardinius dergle* and *Scardinius plotizza* populations (Actinopteri; Cypriniformes) from Croatia and Bosnia and Herzegovina. *Journal of Central European Agriculture* 22: 713–734. <https://doi.org/10.5513/JCEA01/22.4.3189>
- Sadović S. 1986. *Morfološko-taksonomske Karakteristike Alburnus alburnus L., 1758 i Scardinius erythroptalmus L., 1758 iz Hutovog Blata*. Diploma thesis, University of Sarajevo, Sarajevo.
- Safin B. 1997. *Morfološko-biosistematske Odlike Carassius auratus gibelio (Bloch, 1783) na Ribogojištu Bardača*. Diploma thesis, University of Sarajevo, Bosnia and Herzegovina.
- Savić N. 2022. Condition factor and length-weight relationship of huchen (*Hucho hucho*) fingerlings in cultivated conditions. *Agriculture and Forestry* 68 (1): 173–181. <https://doi.org/10.17707/AgricultForest.68.1.10>
- Savić N., Babić R., Ostojić A. & Vaško Ž. 2018. *Dužičasta pastrmka – Gajenje, Tržište i Ekonomika*. Poljoprivredni fakultet Univerziteta u Banjoj Luci, Banja Luka.
- Schneider D., Mrakovčić M., Mustafić P. & Kerovec M. 2000a. Morphological difference in some *Cobitis* populations from Croatia. *Folia Zoologica* 49 (Suppl. 1): 227–234.
- Schneider D., Mustafić P., Mrakovčić M. & Mihaljević Z. 2000b. Some aspects of the biology of the Neretvan spined loach. *Folia Zoologica* 49 (Suppl. 1): 159–165.

- Schöffmann J. & Marić S. 2024. Salmonid fish species: opportunities for sustainable use under multiple pressures and current climatic change. In: Simić V., Simić S. & Pešić V. (eds) *Ecological Sustainability of Fish Resources of Inland Waters of the Western Balkans. Freshwater Fish Stocks, Sustainable Use and Conservation*. *Fish & Fisheries* 43: 375–410. Springer Nature, Switzerland.
- Schönhuth S., Vukić J., Šanda R., Yang L. & Mayden R. 2018. Phylogenetic relationships and classification of the Holarctic family Leuciscidae (Cypriniformes: Cyprinoidei). *Molecular Phylogenetics and Evolution* 127: 781–799. <https://doi.org/10.1016/j.ympev.2018.06.026>
- Seeley H.G. 1886. *The Freshwater Fishes of Europe*. Cassell and Company Limited, London.
- Segherloo I.H., Freyhof J., Berrebi P., Ferchaud A.L., Geiger M., Laroche J., Levin B.A., Normandeau E. & Bernatchez L. 2021. A genomic perspective on an old question: *Salmo* trouts or *Salmo trutta* (Teleostei: Salmonidae)? *Molecular Phylogenetics and Evolution* 162: e107204. <https://doi.org/10.1016/j.ympev.2021.107204>
- Sekulić N., Marić S., Galambos L., Radošević D. & Krpo-Ćetković J. 2013. New distribution data and population structure of the European mudminnow *Umbra krameri* in Serbia and Bosnia and Herzegovina. *Journal of Fish Biology* 83: 659–666. <https://doi.org/10.1111/jfb.12183>
- Seratlić-Savić D. & Vuković T. 1969. Promjenjivost broja kičmenih pršljenova i brahiospina u različitim vrsta iz familije Cyprinidae. In: Martinčić A. (ed.) *III Kongres Biologa Jugoslavije*: 287. Ljubljana.
- Seratlić-Savić D. & Vuković T. 1971. Broj kičmenih pršljenova kod nekih vrsta ciprinida iz voda Bosne i Hercegovine. *Ribarstvo Jugoslavije* 26 (3): 45–49.
- Seratlić D. & Vuković T. 1979. Morfološke karakteristike hibrida *Scardinius erythrophthalmus* (Linnaeus, 1758) × *Leuciscus turskyi* (Heckel, 1843). *Ichthyologia* 11 (1): 25–31.
- Seratlić D., Gužina N., Vuković N. & Vuković T. 1978. Description of some morphological characteristics of the hybrid *Scardinius erythrophthalmus* (Linnaeus, 1758) × *Chondrostoma kneri* Heckel, 1843. *Godišnjak Biološkog instituta Univerziteta u Sarajevu* 31: 159–167.
- Serdarević E. 2012. *Tempo Dužinskog Rasta Predkonzumne Kalifornijske Pastrmke – Oncorhynchus myskiss Walbaum, 1792 na Ribogojilištu UGSR Fojnica*. Diploma thesis, University of Sarajevo, Sarajevo.
- Shubina T.N., Popova A.A. & Vasil'ev V.P. 1989. *Acipenser stellatus* Pallas, 1771. In: Holčík J. (ed.) *The Freshwater Fishes of Europe. Vol 1, Part II, General Introduction to Fishes, Acipenseriformes*: 395–443. AULA-Verlag, Wiesbaden.
- Simić S. 1963. *Morfološke Karakteristike Nemachilus barbatulus L. iz Okoline Banja Luke*. Diploma thesis, University of Sarajevo, Bosnia and Herzegovina.
- Simonović P. 2001. *Ribe Srbije*. NNK Internacional, Beograd, Biološki fakultet Univerziteta u Beogradu, Beograd, Zavod za zaštitu prirode Srbije, Novi Beograd.
- Simonović P.D., Marić S. & Nikolić V. 2000. Growth characteristics of huchen *Hucho hucho* (L.) from Rivers Drina, Una and Sana. *Acta Biologica Iugoslavica – Ekologija Belgrade* 35: 123–126.
- Simonović P.D., Nikolić V.P., Tošić A.D. & Marić S.P. 2011. Length-weight relationship in adult huchen *Hucho hucho* (L., 1758) from Drina River, Serbia. *Biologia (Section Zoology)* 66 (1): 156–159. <https://doi.org/10.2478/s11756-010-0135-2>
- Simonović P.D., Povž M., Piria M., Treer T., Adrović A., Škrijelj R., Nikolić V. & Simić V. 2015. Ichthyofauna of the River Sava system. In: Milačić R., Ščančar J. & Paunović M (eds) *The Sava River. The Handbook of Environmental Chemistry* 31: 361–400. Springer, Berlin-Heidelberg. https://doi.org/10.1007/978-3-662-44034-6_14

Simonović P.D., Tošić A., Škraba Jurlina D., Nikolić V., Piria M., Tomljanović T., Šprem N., Mrdak D., Milošević D., Bećiraj A., Dekić R. & Povž M. 2017. Diversity of Brown trout *Salmo cf. trutta* in the River Danube Basin of Western Balkans as assessed from the structure of their mitochondrial control region haplotypes. *Journal of Ichthyology* 57 (4): 603–616. <https://doi.org/10.1134/S0032945217040154>

Skalin B. 1990. Mladica kao ugrožena riblja vrsta. *Ribarstvo Jugoslavije* 1–2: 22–30.

Skenderović I. 2015. *Biodiverzitet Parazita Riba iz Hidroakumulacije Modrac*. “OFF-SET” Tuzla, Tuzla.

Skenderović I. & Adrović A. 2008a. Ecological characteristics of Bream from Spreča River. In: Dikić B. (ed.) *Second International Conference “Ekologija, Zdravlje, Rad, Sport”*: 202–205. Udruženje Zdravlje za sve, Banja Luka.

Skenderović I. & Adrović A. 2008b. Population of Cyprinidae fish in the upper flow of the river Spreča. In: Dikić B. (ed.) *Drugi međunarodni kongres “Ekologija, Zdravlje, Rad, Sport”*: 198–201. Udruženje Zdravlje za sve, Banja Luka.

Skenderović I. & Adrović A. 2012. Istraživanje sezonske dinamike nekih helminata zele (*Alburnus alburnus*) i šarana (*Cyprinus carpio*) iz hidroakumulacije Modrac. In: Redžić S. (ed.) *Structure and Dynamics of the Dinarid Ecosystem – State, Possibilities and Perspectives. Special editions/CXLIX Zbornik radova/Proceedings* 23: 305–314. Academy of Sciences and Arts of Bosnia and Herzegovina, Department of Natural and Mathematical Sciences, Sarajevo.

Skenderović I. & Adrović A. 2016. Ektoparaziti riba iz akumulacije Modrac i Ramičko jezero. Voda i mi. *Journal of the Agency for the Sava River Water Area* 93: 16–21.

Skenderović I. & Adrović A. 2019. The Diversity of Trematodes in Fishes from the Middle Course of the River Sava (Bosnia and Herzegovina). *Fondeko Science* 1 (1): 86–95.

Skenderović I., Adrović A. & Škrijelj R. 2006. Biodiverzitet ihtiofaune rijeke Spreče uzvodno od hidroakumulacije Modrac. In: Opačak A. & Florijančić T. (eds) *Zbornik Priopćenja, Ribarski Dani u Osijeku, IV. Nacionalno Znanstveno Stručno Savjetovanje s Međunarodnim Sudjelovanjem*: 41–52. Poljoprivredni fakultet Osijek, lipanj.

Skenderović I., Adrović A. & Terzić R. 2010. Study of seasonal dynamics at found trematoda of cyprinid fish in hydro-accumulation Modrac. In: Biočanin R. (ed.) *Naučno-stručna Konferencija sa Međunarodnim Učešćem, Rizici i Eko-bezbednost u Postmodernom Ambijentu, Eko-Undp* 2010: 41–46. State University in Novi Pazar, Novi Pazar.

Skenderović I., Žujo D., Adrović A. & Marković G. 2011a. Trematode fauna of some fish species in the river Spreča. In: Marković Z. (ed.) *V International Conference “Aquaculture & Fishery”, Conference Proceedings*: 294–300. Faculty of Agriculture, Belgrade-Zemun.

Skenderović I., Jažić A., Adrović A., Terzić R., Hajdarević E., Lonić E. & Bajrić A. 2011b. Fauna trematoda i cestoda – parazita nekih ciprinidnih riba iz hidroakumulacije Modrac (Bosna i Hercegovina). In: *International Conference Nature protection in XXIst Century. Proceedings of the Conference (Book No 2)*: 397–401. Zavod za zaštitu prirode Crne Gore, Ministarstvo održivog razvoja i turizma, Žabljak.

Skenderović I., Adrović A., Hajdarević E., Hadžiahmetović Jurida E., Čekmić M. & Bajrić A. 2015. Parazitski rakovi (Crustacea) ciprinidnih riba iz hidroakumulacije Modrac. *Proceedings of Scientific Conference, Lukavac*: 669–676.

Skenderović I., Adrović A., Hajdarević E., Hadžiahmetović Jurida E., Jažić A. & Krčmar S. 2016. Frequency of emergence of parasitic protozoa on *Ameiurus melas* in lake Modrac (Bosnia and Herzegovina). *Veterinary Journal of Republic of Srpska* 16 (2): 152–159.

- Skenderović I., Adrović A., Hajdarević E. & Bajrić A. 2020a. *Comparative Composition of Spreča River Fauna (Bosnia and Herzegovina)*. In: Karabegović I (ed.) *International Conference “New Technologies, Development and Applications” III. Lecture Notes in Networks and Systems* 128: 795–802. https://doi.org/10.1007/978-3-030-46817-0_90
- Skenderović I., Adrović A., Jažić A., Zuko A. & Hadžimustafić E. 2020b. Review of freshwater fish parasitofauna of Bosnia and Herzegovina. *Biologia* 76 (3): 475–515. <https://doi.org/10.2478/s11756-020-00565-0>
- Skenderović I., Nedić Z. & Adrović A. 2021. The diversity of trematodes in fishes from the middle course of the river Sava (Bosnia and Herzegovina). *Works of the Faculty of Forestry University of Sarajevo* 50 (1): 12–18. <https://doi.org/10.54652/rsf.2020.v50.i1.37>
- Sket B. 1967. *Ključni za Določevanje Živali*. I Slatkovodne ribe. Pisces. Inštitut za biologijo Univerze v Ljubljani. Ljubljana.
- Skoulikidis N.T., Zogaris S. & Karaouzas I. 2022. *Rivers of the Balkans*. In: Tockner K., Zarfl C. & Robinson C.T. (eds) *Rivers of Europe* (2nd edition): 595–655. Elsevier, Amsterdam. <https://doi.org/10.1016/B978-0-08-102612-0.00015-8>
- Smith K.G. & Darwall W.R.T. 2006. *The Status and Distribution of Freshwater Fish Endemic to the Mediterranean Basin*. IUCN, Switzerland.
- Smlatić S. 1961. Una – jedinstvena rijeka. *Priroda* 8: 225–229.
- Snoj A., Melkič E., Sušnik S., Muhamedagić S. & Dovč P. 2002a. DNA phylogeny supports revised classification of *Salmothymus obtusirostris*. *Biological Journal of the Linnean Society* 77: 399–411. <https://doi.org/10.1046/j.1095-8312.2002.00130.x>
- Snoj A., Sušnik S., Odak T., Safner R. & Dovč P. 2002b. Phylogenetic relationship between two subspecies of soft-mouth trout *Salmothymus obtusirostris oxyrhynchus* and *S. obtusirostris salonitana*. In: Heller S.R. (ed.) *Plant and Animal Genomes XI Conference, Book of Abstracts*: 116. Scherago International, San Diego.
- Snoj A., Glamuzina B., Razpet A., Zabolcki J., Bogut I., Lerceteau-Kohler E., Pojskić N. & Sušnik S. 2010. Resolving taxonomic uncertainties using molecular systematics, *Salmo dentex* and the Balkan trout community. *Hydrobiologia* 651 (1): 199–212. <https://doi.org/10.1007/s10750-010-0297-5>
- Sofradžija A. 1977. Kariologija i citotaksonomija vrsta roda *Leuciscus* iz voda Bosne i Hercegovine. *Godišnjak Biološkog instituta Univerziteta u Sarajevu* 30: 113–211.
- Sofradžija A. 1979. The chromosomes of *Hucho hucho* (L.). In: Kullander S. (ed.) *Third European Ichthyological Congress, Warszawa*: 166. EIC, Warsaw.
- Sofradžija A. 1982a. Hromosomi potočne pastrmke (*Salmo trutta* m. *fario* i *Salmo gairdneri*). *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 35: 143–153.
- Sofradžija A. 1982b. Hromosomi *Silurus glanis* L. 1758 (Siluridae, Pisces). *Genetika* 14 (2): 103–110.
- Sofradžija A. 2009. *Slatkovodne Ribe Bosne i Herecegovine*. Vijeće kongresa bošnjačkih intelektualaca, Sarajevo.
- Sofradžija A. & Berberović Lj. 1972. Uperedna kariološka istraživanja vrsta *Paraphoxinus alepidotus*, *P. adspersus*, *P. pstrossi*, *P. metohiensis* i *P. croaticus*. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 25: 135–173.
- Sofradžija A. & Berberović Lj. 1973. The chromosome number of *Barbus meridionalis petenyi* Heckel (Cyprinidae, Pisces). *Bulletin Scientifique* 18 (4–6): 77–78.

- Sofradžija A. & Berberović Lj. 1975. Osnovni podaci o hromosomskoj garnituri vrste *Gobio gobio* (Linnaeus, 1758): Cyprinidae, Pisces. *Ichthyologia* 7 (1): 53–59.
- Sofradžija A. & Berberović Lj. 1978. Diploidno-triploidni seksualni dimorfizam u vijuna (*Cobitis taenia taenia* L., Cobitidae, Pisces). *Genetika* 10 (3): 389–397.
- Sofradžija A. & Hadžiselimović R. 1981a. Chromosome complement of the species (*Chondrostoma nasus*, Cyprinidae, Pisces). *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 34: 107–115.
- Sofradžija A. & Hadžiselimović R. 1981b. Chromosomes of Yugoslav freshwater cyclostomata and pisces. *Godišnjak biološkog Instituta Univerziteta u Sarajevu* 34: 117–152.
- Sofradžija A. & Kundurović Z. 1988. Cytogenetical characteristics of the species *Aulopyge hügelii* Heckel, 1841 (Cyprinidae, Pisces). *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 41: 63–68.
- Sofradžija A. & Vuković T. 1979. Hromosomi brkice *Nemachilus barbatus* (Linnaeus, 1758): Cobitidae, Pisces. *Ichthyologia* 11 (1): 5–13.
- Sofradžija A., Hadžiselimović R. & Berberović Lj. 1969. Hromozomski broj vrste *Paraphoxinus alepidotus*. III Kongres Biologa Jugoslavije, Knjiga Plenarnih Referata: 242. Martinčić A., Ljubljana.
- Sofradžija A., Berberović Lj. & Pavlović B. 1972. Broj hromosoma u ćelijama sjemenika vrste *Paraphoxinus pstrossi* Steind. *Genetika* 4 (1): 17–23.
- Sofradžija A., Hadžiselimović R. & Marić C. 1975. Basic data about the chromosome complement of *Rhodeus sericeus amarus* (Bloch 1872): Cyprinidae, Pisces. *Bulletin Scientifique, Section A* 20 (1–2): 7–8.
- Sofradžija A., Berberović Lj. & Hadžiselimović R. 1977. Hromosomi običnog i srebrenog karaša (*Carassius carassius* i *Carassius auratus gibelio*). In: Naučni Skup "Struktura, dinamika i zaštita ekosistema i Njihovih Komponenata na Dinaridima", Rezimeji Referata: 35–36. Biološki institut Univerziteta u Sarajevu, Sarajevo.
- Sofradžija A., Berberović Lj. & Hadžiselimović R. 1978a. Hromosomske garniture karaša (*Carassius carassius*) i babuške (*Carassius auratus gibelio*). *Ichthyologia* 10 (1): 135–142.
- Sofradžija A., Berberović Lj. & Šahinović N. 1978b. Hromosomi crvenperke (*Scardinius erythrophthalmus* (L. 1758) Cyprinidae, Pisces). *Rad Akademije Nauka Umjetnosti BiH, Sekcija Prirodno Matematskih Nauka* 18: 131–142.
- Sofradžija A., Berberović Lj. & Hadžiselimović R. 1979. Hromosomske garniture vrsta *Alburnus alburnus* (L. 1758) i *Alburnoides bipunctatus* (Bloch 1782), Cyprinidae, Pisces. *Ichthyologia* 11 (1): 33–41.
- Sofradžija A., Vuković T. & Hadžiselimović R. 1980. Efekti nekih pesticida i deterdženata u hromosomskim garniturama riba – rezultati preliminarnih istraživanja. *Ichthyologia* 12 (1): 71–76.
- Sofradžija A., Vuković T. & Hadžiselimović R. 1984. Ichthyofauna of river Mlječanica on a wide area of the future thermal spring touristic center of Međuvođe (Bos. Dubica). *Ribarstvo Jugoslavije* 39 (2): 25–29.
- Sofradžija A., Hadžiselimović R., Škrijelj R., Guzina N. & Korjenić E. 2005. Biodiverzitet ihtiopopulacija sliva gornjeg toka rijeke Krivaje. *Veterinaria* 54 (3–4): 245–258.
- Sokolov L.I. & Vasil'ev V.P. 1989a. *Acipenser ruthenus* Linnaeus, 1758. In: Holčík J (ed.) *The freshwater fishes of Europe. Vol. 1, Part II, General Introduction to Fishes, Acipenseriformes*: 227–262. AULA-Verlag, Wiesbaden.
- Sokolov L.I. & Vasil'ev V.P. 1989b. *Acipenser nudiiventris* Lovetsky, 1828. In: Holčík J (ed.) *The Freshwater Fishes of Europe. Vol 1, Part II, General Introduction to Fishes, Acipenseriformes*: 206–226. AULA-Verlag, Wiesbaden

- Spasojević P. 2014. *Procjena genetičkog Diverziteta Vrste Salmo trutta (Linnaeus, 1758) u Rijeci Radobolji, u Cilju Revitalizacije Populacija*. PhD thesis, University of Sarajevo, Sarajevo.
- Stanić M. 2013. *Molekularno-genetička Obilježja Jedinki Introducirane Vrste Salvelinus fontinalis (Mitchill, 1848) u Bosni i Hercegovini*. PhD thesis, University of Sarajevo, Sarajevo.
- Stanković D., Crivelli A.J. & Snoj A. 2015. Rainbow trout in Europe: introduction, naturalization, and impacts. *Reviews in Fisheries Science & Aquaculture* 23 (1): 39–71. <https://doi.org/10.1080/23308249.2015.1024825>
- Stanković D., Stephens M.R. & Snoj A. 2016. Origin and introduction history of self-sustaining rainbow trout populations in Europe as inferred from mitochondrial DNA and a Y-linked marker. *Hydrobiologia* 770 (1): 129–144. <https://doi.org/10.1007/s10750-015-2577-6>
- Steindachner F. 1882a. Beiträge zur Kenntnis der Fische Afrikas (II.) und Beschreibung einer neuen *Paraphoxinus* Art aus den unterirdischen Gewässern in der Herzegowina. Sitzung der mathematisch-naturwissenschaftlichen Classe vom 9. Februar 1882. *Anzeiger Kaiserliche Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse* 19 (5): 41–43.
- Steindachner F. 1882b. Ichthyologische Beiträge (XII.) und berichtet in derselben Sitzung der mathematisch-naturwissenschaftlichen Classe vom 22. Juni 1882. *Anzeiger Kaiserliche Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse* 19 (16): 139–143.
- Steindachner F. 1882c. Beiträge zur Kenntnis der Fische Afrikas (II.) und Beschreibung einer neuen *Paraphoxinus* Art aus der Herzegowina. *Anzeiger Kaiserliche Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse* 45: 1–18.
- Steindachner F. 1883. Ichthyologische Beiträge (XII.). *Anzeiger Kaiserliche Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse* 86 (Jahrgang 1882): 61–82.
- Steindachner F. 1901. Bericht über das Vorkommen einer bisher noch unbeschriebenen *Paraphoxinus* – Art. *Anzeiger der Kaiserlichen Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse* 38: 197–198.
- Stepanek O. 1953. Još nešto o ishrani lipljena i načinu njegova života. *Ribarski List, Sarajevo* 6: 115–118.
- Stierandová S., Vukić J., Vasil'eva E.D., Zogaris S., Shumka S., Halačka K., Vetešník L., Švátora M., Nowak M., Stefanov T., Koščo J. & Mendel J. 2016. A multilocus assessment of nuclear and mitochondrial sequence data elucidates phylogenetic relationships among European spirrins (Alburnoidea, Cyprinidae). *Molecular Phylogenetics and Evolution* 94: 479–491. <https://doi.org/10.1016/j.ympev.2015.10.025>
- Suljević D. 2012. *Biohemijski Status Vrsta Salmo trutta fario (Linnaeus, 1758) i Salmo farioides (Karaman, 1938) Tokom i Nakon Mrijesta u Centru za Ribarstvo Neretva – Konjic*. PhD thesis, University of Sarajevo, Sarajevo.
- Sušnik S., Snoj A., Wilson I.F., Mrdak D. & Weiss S. 2007a. Historical demography of brown trout (*Salmo trutta*) in the Adriatic drainage including the putative *S. letnica* endemic to Lake Ohrid. *Molecular Phylogenetics and Evolution* 44: 63–76. <https://doi.org/10.1016/j.ympev.2006.08.021>
- Sušnik S., Weiss S., Odak T., Dellling B., Treer T. & Snoj A. 2007b. Reticulate evolution: Ancient introgression of the Adriatic brown trout mtDNA in softmouth trout *Salmo obtusirostris* (Teleostei: Salmonidae). *Biological Journal of the Linnean Society* 90 (1): 139–152. <https://doi.org/10.1111/j.1095-8312.2007.00717.x>
- Suić J., Šarić M., Homen Z., Jahutka I. & Mišura A. 2008. Gospodarski ribolov na slatkim vodama Republike Hrvatske u 2006. godini. *Ribarstvo* 1: 17–24.
- Šanda R. & Kovačić M. 2009. Freshwater gobies in the Adriatic drainage basin of the Western Balkans. *Annales, anali za istrske in mediteranske študije. Series Historia Naturalis* 19 (1): 1–10.

- Šanda R. & Vukić J. 2009. Agnatha in the ichthyologic collection of the National Museum in Prague. *Bulletin Lampetra* 6: 56–65.
- Šanda R., Bogut I. & Vukić J. 2008a. Novi podaci o ihtiofauni slijeva donje Neretve i okolnih kraških polja u Bosni i Hercegovini. Uzgoj slatkovodne ribe, stanje i perspektive. *Hrvatska Gospodarska Komora*: 119–125.
- Šanda R., Bogut I., Doadrio I., Kohout J., Perdices A., Perea S., Šedivá A. & Vukić J. 2008b. Distribution and taxonomic relationships of spined loaches (Cobitidae, Cobitis) in the River Neretva basin, Bosnia and Herzegovina. *Folia Zoologica* 57 (1–2): 20–25.
- Šanda R., Vukić J., Marić D. & Bogut I. 2009. Sastav populacija riba Mostarskog blata. Uzgoj slatkovodne ribe u otvorenim vodama – stanje i perspektive. *Hrvatska Gospodarska Komora*: 117–125.
- Šenk O. 1953. Ispitivanje rasteinje i spolnog sazrijevanja lipljenova (*Thymallus thymallus* L.) iz rijeke Drine, Bosne i Vrbasa. *Veterinaria* 2 (4): 14–22.
- Šenk O. 1954. Ispitivanje razvoja krljušti kod mlađi potočnih pastrmki. *Ribarstvo Jugoslavije* 3: 50–57.
- Šenk O. 1956a. Prilog poznavanju režima ishrane lipljenova izvorišnog dijela rijeke Bosne. *Ribarstvo Jugoslavije* 3: 50–57.
- Šenk O. 1956b. Prilog poznavanju režima ishrane potočnih pastrmki (*S. trutta fario*) izvorskog dijela rijeke Bosne. *Ribarstvo Jugoslavije* 11 (3–4): 57–65.
- Šenk O. 1962. Prilog poznavanju razvitka krljušti kod *Phoxinus phoxinus* L. (gaga, pijor). *Radovi Poljoprivrednog Fakulteta u Sarajevu* 13.
- Šenk O. 1963. Prilog poznavanju krljušti *Barbus meridionalis petenyi*. *Ribarstvo Jugoslavije* 2: 32–40.
- Šenk O. 1969. Aberacije u razvitku krljušti kod nekih vrsta riba. *Ichthyologia* 1 (1): 69–81.
- Šenk O. & Aganović M. 1968. Prilog ispitivanju ishrane riba rijeke Vrbanje. *Ribarstvo Jugoslavije* 23 (4): 77–83.
- Šenk O. & Kaluđerčić M. 1963. Prilog poznavanju krljušti *Gobio gobio*. *Ribarstvo Jugoslavije* 18 (5): 139–147.
- Šenk O. & Kosorić Đ. 1968. Ispitivanje razvitka krljušti kod mlađi neretvanske mekousne (*Salmothymus obtusirostris oxyrhynchus* Steind. *Radovi Poljoprivrednog Fakulteta u Sarajevu* 17 (19): 93–109.
- Šenk O. & Telalbašić R. 1963. Prilog ispitivanju varijacija oblika krljušti kod *Barbus meridionalis petenyi* (Heckel). *Radovi Poljoprivrednog Fakulteta u Sarajevu* 12 (14): 213–228.
- Šinžar D. 1955. Prilog poznavanju entoparazita pastrmke *Salmo trutta* L. *Glasnik. Prirodnjačkog Muzeja Srpske zemlje Serie B, VII* (4): 223–229.
- Škraba D., Bećiraj A., Šarić I., Ićanović I., Džaferović A., Piria M., Dekić R., Tošić A., Nikolić V. & Simonović P. 2017. Haplotype diversity of brown trout (*Salmo trutta* L.) populations from Una River drainage area in Bosnia and Herzegovina: implications for conservation and fishery management. *Acta Zoologica Bulgarica* 69 (1): 25–30.
- Škraba Jurlina D., Marić A., Mrdak D., Kanjuh T., Špelić I., Nikolić V., Piria M. & Simonović P. 2020. Alternative life-history in native trout (*Salmo* spp.) suppresses the invasive effect of alien trout strains introduced into streams in the western part of the Balkans. *Frontiers in Ecology and Evolution* 8:188. <https://doi.org/10.3389/fevo.2020.00188>
- Škrijelj R. 1991. Srebrni karaš-babuška (*Carassius auratus gibelio* Bloch) – nova vrsta u ihtiofauni akumulacionog jezera “Jablanica”. *Biološki List* 38: 148–150.
- Škrijelj R. 2002. *Populacije Riba Neretvanskih Jezera*. Ihtiološka monografija. Prirodoslovno-matematički fakultet Univerziteta u Sarajevu, Sarajevo.

- Škrijelj R. & Sofradžija A. 1990. Nalaz vrste (*Phoxinellus alepidotus* Heck.) u Jablaničkom jezeru. *Ribarstvo Jugoslavije* 45 (4): 71–72.
- Škrijelj R. & Mašović M. 2001. Populacija smuđa (*Stizostedion lucioperca* L.) u neretvanskim hidroakumulacijama. *Ribarstvo* 59 (2): 57–69.
- Škrijelj R. & Mašović M. 2008. Uticaj formiranja hidroakumulacija na rijeci Neretvi na dinamiku populacija vrsta strugača *Leuciscus svalizze svalizze* (Heckel et Kner, 1858) i bijelog kljena *Leuciscus cephalus albus* (Bonaparte, 1838). *Veterinaria* 57 (3–4): 207–218.
- Škrijelj R., Korjenić E. & Guzina N. 2003. Populacije riba hidroakumulacije Grabovica. *Veterinaria* 52 (1–4): 1–10.
- Škrijelj R., Sofradžija A., Hadžiselimović R., Korjenić E., Guzina N., Adrović A., Muhamedagić S., Mihaljica M., Džano A. & Šljuka S. 2005a. *Gospodarenje Slatkovodnim Ribljim Resursima na Pograničnim Rijekama. Pilot Studija s Cjelovitim Regionalnim Pristupom*. Federalno ministarstvo poljoprivrede, vodoprivrede i šumarstava, Sarajevo.
- Škrijelj R., Sofradžija A., Hadžiselimović R., Spahić M., Korjenić E., Guzina N., Muhamedagić S., Muzaferović Š., Adrović A. & Drešković N. 2005b. Hydrological and ichthyological characteristics of Sniježnica Lake. *Works of the Faculty of Agriculture, University of Sarajevo* 50 (56): 63–74.
- Škrijelj R., Sofradžija A., Hadžiselimović R., Korjenić E., Trožić-Borovac S., Guzina N., Đug S., Čekić A. & Šljuka S. 2006. Ichthyofauna of the river Una – current state and sustainable management. In: *Zbornik Priopćenja, Ribarski Dani u Osijeku, IV. Nacionalno Znanstveno Stručno Savjetovanje s Međunarodnim Sudjelovanjem*: 29–40. Poljoprivredni fakultet Osijek, Osijek.
- Škrijelj R., Mitrašinović M., Ivanc A., Krnjić J., Đug S., Šljuka S. & Mušović A. 2011. Ecological, morphological-taxonomic and chematological characterisation of pike perch (*Sander lucioperca*, Linnaeus 1758) from reservoirs on the river Neretva. *Ribarstvo* 69 (2): 71–79.
- Škrijelj R., Lelo S., Sofradžija A., Trožić-Borovac S., Korjenić E., Lukić-Bilela L., Mitrašinović-Brulić M., Kotrošan D., Šljuka S., Gajević M., Drešković N. & Karačić J. 2013. *Red List of Endangered Plants, Animals and Mushrooms in the Federation of Bosnia and Herzegovina. Red List of Fauna of the Federation of Bosnia and Herzegovina, Book 3*. The Federal Ministry of Environment and Tourism, EU “Greenway” Sarajevo and PMF University in Sarajevo, Sarajevo.
- Šlechtová V., Bohlen J., Freyhof J., Persat H. & Delmastro G. 2004. The Alps as barrier to dispersal in cold-adapted freshwater fishes? Phylogeographic history and taxonomic status of the bullhead in the Adriatic freshwater drainage. *Molecular Phylogenetics and Evolution* 33: 225–39. <https://doi.org/10.1016/j.ympev.2004.05.005>
- Šljuka S. 2011. *Morfološko-taksonomske i Ekološka Karakterizacija Vrste Sander Lucioperca (Linnaeus 1758), Percidae, Perciformes iz Hidroakumulacija na Rijeci Neretvi i Rami*. Master’s thesis, University of Sarajevo, Sarajevo.
- Šljuka S. 2016. *Komparativna Analiza Odabranih Merističkih Značajki i Molekularnih Markera u Određivanju Strukture i Genetičkog Diverziteta Populacija Sander lucioperca (Linnaeus, 1758) u slivu Rijeka Neretve i Save (Linnaeus 1758)*. PhD thesis, Prirodno-matematički fakultet Univerziteta u Sarajevu.
- Šolaja M. & Pocrnjić Z. 1983a. Uticaj termičke adaptacije (aklimacije) na potrošnju kiseonika potočne mreše (*Barbus meridionalis petenyi*). *Ichthyologia* 15 (1): 37–42.
- Šolaja M. & Pocrnjić Z. 1983b. Otpornost skeletnog mišićnog tkiva kljena (*Leuciscus cephalus*) na povišene temperature. *Ichthyologia* 15 (2): 47–53.

- Šorić V. 1992. Osteological characteristics in six species of the genus *Phoxinellus* Heck, 1843. *Ichthyologia* 24 (1): 43–56.
- Šorić V. 1998. New data on the taxonomy and distribution of lampreys (Petromyzontidae) in southern Europe. *Ichthyologia* 30 (1): 71–82.
- Šorić V.M. 2007. Taxonomic status of the species of the genus *Leuciscus* (Pisces, Cyprinidae) from the Adriatic and Aegean basins. *Kragujevac Journal of Science* 29: 155–169.
- Šorić V. & Janković D. 1989. Characteristics of growth and sexual maturity of *Barbus meridionalis*. *Acta Biologica Iugoslavica-Ichthyologia* 21: 27–37.
- Šorić V.M. & Bănărescu P.M. 1999. *Aulopyge* Heckel, 1841. In: Bănărescu PM (ed.) *The Freshwater Fishes of Europe Vol. 5/I, Cyprinidae* 2: 369–381. AULA-Verlag, Wiebelsheim.
- Šukalo G., Đorđević S., Dmitrović D. & Tomović Lj. 2012. Introduced fish *Ameiurus nebulosus* (Le Sueur, 1819): hazard to the Grass snake, *Natrix natrix* (Laurenti, 1768). *Hyla* 2: 41–42.
- Šukalo G., Dmitrović D. & Golub D. 2018. First record of weatherfish *Misgurnus fossilis* (Linnaeus, 1758) from Adriatic Sea catchment area in Bosnia and Herzegovina. *Ecologica Montenegrina* 18: 126–128. <https://doi.org/10.37828/em.2018.18.12>
- Švob T. & Kilalić T. 1972. Prilog morfologiji digestivnog sistema nekih salmonida i ciprinida rijeke Neretve. *Ichthyologia* 4 (1): 83–88.
- Tadić A. 1931. Pastruga. *Ribarski List* 7–10: 114.
- Tadić A. 1932. Zoološki rariteti u Podunavlju. *Priroda* 9–10: 288–289.
- Tadić A. 1966. Riblji svijet i đerdapska hidroelektrana. *Priroda* 9: 273.
- Takács P., Bihari P., Erős T., Specziár A., Szivák I., Bíró P. & Csoma E. 2014. Genetic heterogeneity reveals on-going speciation and cryptic taxonomic diversity of stream-dwelling gudgeons (Teleostei, Cyprinidae) in the Middle Danubian hydrosystem (Hungary). *PLoS ONE* (9): e97278. <https://doi.org/10.1371/journal.pone.0097278>
- Taler Z. 1932. Slatkovodne ribe. In: Thaller Z. (ed.) *Vode i Ribe Jugoslavije u Slici i Riječi*: 59–65. Tipografija, Zagreb.
- Taler Z. 1944. *Lipljen Thymallus thymallus* L, Njegovo Životno Područje u Hrvatskoj i na Balkanu, te Njegova Gospodarska Vrijednost. Tisak „Hrvatske tiskare“, Zagreb.
- Taler Z. 1945a. *Mladica-glavatica (Salmo hucho L.) i Neretvanska glavatica (Salmo marmoratus Cuv.)*: Zoogeografski Podatci i Gospodarska Vrijednost. Pučka knjižara Priroda, Zagreb.
- Taler Z. 1945b. *Podust – Chondrostoma nasus (Linne)*. Zoogeografski Podaci, Prehrambena i Gospodarska Vrijednost te Razna Opažanja. Nakladni zavod Hrvatske, Zagreb.
- Taler Z. 1948. Kečiga, jesetra i njeni srodnici. *Ribarstvo Jugoslavije* 7–8: 57.
- Taler Z. 1950. Salmonidi koji žive u Jugoslaviji i koji se uzgajaju. *Ribarstvo Jugoslavije* 3–4: 58–60.
- Taler Z. 1951a. Podzemne ribe u našem kršu. *Slatkovodno Ribarstvo Jugoslavije* 6 (4): 107–109.
- Taler Z. 1951b. O poribljavanju. *Ribarstvo Jugoslavije* 5: 127–130.
- Taler Z. 1951c. Zaštita naših rijetkih i neobičnih riba. *Slatkovodno Ribarstvo Jugoslavije* 6 (10): 224–227.
- Taler Z. 1951d. Međusobni odnosi između urođenih potočnih pasrva i novonaseljenih pastrva prigodom poribljavanja. *Ribarstvo Jugoslavije* 12: 285–287.
- Taler Z. 1952. Trebišnjica. *Ribarstvo Jugoslavije* 7 (9–10): 160–162.

- Taler Z. 1953a. Rasprostranjenje i popis slatkovodnih riba Jugoslavije. *Glasnik Prirodnjačkog Muzeja Srpskih Zemalja (serija B)* 5–6: 425–455.
- Taler Z. 1953b. Znamenite i osobite ribe u Bosni i Hercegovini. *Ribarski List Sarajevo* 28 (4): 56–57.
- Taler Z. 1953c. Značaj kalifornijske pastrve u Trebišnjici. *Ribarstvo Jugoslavije* 1: 11–14.
- Taler Z. 1954. Kalifornijska pastrva i njeno udomaćenje u Jugoslaviji. *Ribarstvo Jugoslavije* 4: 82–89.
- Tanović E., Hajdarević E., Adrović A., Skenderović I., Bajrić A. & Tanović A. 2022. Length-weight relationships and condition factor of chub (*Squalius cephalus* Linnaeus, 1758) from the area of northeastern Bosnia and Herzegovina. *TEM Journal* 11 (1): 169–173. <https://doi.org/10.18421/TEM111-20>
- Tanović E., Hajdarević E., Adrović A. & Skenderović I. 2023. Differential blood picture of chub (*Squalius cephalus* Linnaeus, 1758) from the water courses of northeast Bosnia. In: Bajrović K. (ed.) *International Conference of Biochemists and Molecular Biologists in Bosnia and Herzegovina – ABMBBIH. Genetics & Applications ABMBBIH, Special Edition, Book of Abstract, Poster presentation S5–08*: 142. Institute for Genetic Engineering and Biotechnology, University of Sarajevo, Sarajevo.
- Taylor J.N., Courtenay W.R. & Jr McCann J.A. 1984. Known impacts of non-native fishes in the continental United States. In: Courtenay W.R. & Jr Stauffer J.R. (eds) *Distribution, Biology and Management of Non-native Fishes*: 322–373. John Hopkins University Press, Baltimore, MD.
- Thacker C.E., Gkenas C., Triantafyllidis A., Malavasi S. & Leonardos I. 2019. Phylogeny, systematics, and biogeography of the European sand gobies (Gobiiformes: Gobionellidae). *Zoological Journal of the Linnean Society* 185 (1): 212–225. <https://doi.org/10.1093/zoolinnean/zly026>
- Tinjak E. 2005. *Morfološko-taksonomske karakteristike klijena Leuciscus cephalus (Linnaeus, 1758) iz rijeke Bosne*. Diploma thesis. University of Sarajevo, Sarajevo.
- Tomljanović T., Piria M., Treer T., Safner R., Šprem N., Aničić I., Matulić D. & Kordić V. 2011. Morfološke osobine šaranskih populacija (*Cyprinus carpio carpio* L.) u Republici Hrvatskoj. *Ribarstvo* 69 (3): 81–93.
- Tomljanović T., Treer T., Čubrić V.Č., Safner R., Šprem N., Piria M., Matulić D., Safner R. & Aničić I. 2013. Microsatellite-based genetic variability and differentiation of hatchery and feral common carp *Cyprinus carpio* L (Cyprinidae; Cypriniformes) populations in Croatia. *Archives of Biological Sciences Belgrade* 65 (2): 577–584. <https://doi.org/10.2298/ABS1302577T>
- Tortonese E. 1970. *Osteichthyes (Pesci ossei). Fauna d'Italia. Parte Prima*. Edizioni Calderini, Bologna.
- Tougard C., Vukić J., Ahnelt H., Buj I., Kovačić M., Moro G.A., Tutman P. & Šanda R. 2021. Quaternary climatic cycles promoted (re)colonization and diversification events in Adriatic sand gobies. *Journal of Zoological Systematics and Evolutionary Research* 59: 1037–1052. <https://doi.org/10.1111/jzs.12468>
- Treer T., Habeković D., Aničić I., Safner R. & Piria M. 2000. Salmonidne vrste i njihov uzgoj. *Agriculturae Conspectus Scientificus* 65 (3): 175–180.
- Trewavas E. 1971. The type-species of the genera *Phoxinellus*, and *Paraphoxinus* (Pisces, Cyprinidae). *Bulletin of the British Museum (Natural History). Zoology Series* 21: 359–361.
- Trgovčević L. 1905. *Paraphoxinus* Bikr. i *Telestes* Bonap. u vodama Like i Krbave. *Nastavni Vjesnik Zagreb* 14: 1–23.
- Trojanović S. 1934. O rečnim ribama. *Priroda* 1: 13–19.
- Trožić-Borovac S. 2002. Prehrana potodne pastrve *Salmo trutta* morfa *fario* L, u rijeci Uni. *Ribarstvo* 3: 83–104.
- Trožić-Borovac S. & Škrijelj R. 2007. The nutrition of a pikeperch *Stizostedion lucioperca* Linnaeus, 1758 in the river Neretva. *Ribarstvo* 65: 61–74.

- Trožić-Borovac S., Sofradžija A., Hadžiselimović R., Škrijelj R., Guzina N., Korjenić E. & Hamzić A. 2003. Pojava sunčanice *Lepomis gibbosus* Linnaeus, 1758 u hidroakumulaciji Jablanica kao rezultat slučajne introdukcije. *Radovi Poljoprivrednog Fakulteta Univerziteta u Sarajevu* 53: 13–18.
- Trožić-Borovac S., Škrijelj R., Trakić-Juvan S. & Bakrač-Bećiraj A. 2005. Ishrana potočne pastrmke (*Salmo trutta* morfo *fario* L.) u rijeci Bosni. *Radovi Poljoprivrednog fakulteta Univerziteta u Sarajevu* 5 (55): 22–32.
- Trožić-Borovac S., Škrijelj R., Mitrašinović-Brulić M., Muhamedagić S., Hamzić A., Đug S. & Šljuka S. 2013. Diet and condition coefficient of the Grayling (*Thymallus thymallus* L.) from the Una river basin. *Radovi Poljoprivredno-prehrambenog fakulteta Univerziteta u Sarajevu* 63 (1): 7–17.
- Tutman P., Glamuzina B., Bartulović V. & Buntić I. 2002. Endemična ihtiofauna močvare Hutovo Blato. In: Buntić I. (ed.) *Nova politika gospodarenja vlažnim područjima Hutova Blata. Završni simpozij LIFE Projekta LIFE CY 99/BIH/035*: 45–48. LIFE and Ministarstvo graditeljstva, prostornog uređenja i zaštite okoliša HNŽ-K, Mostar.
- Tutman P., Matić-Skoko S., Glamuzina B., Bartulović V. & Dulčić J. 2006. Some biological parameters of Neretvan spined loach, *Cobitis narentana* (Karaman 1928) from Hutovo blato wetland, Bosnia and Herzegovina. In: Mrakovčić M., Buj I. & Zanella L. (eds) *3rd International Conference, Loaches of the genus Cobitis and Related Genera, Biology, Systematics, Genetics, Distribution, Ecology and Conservation*: 64. Hrvatsko ihtiološko društvo, Zagreb.
- Tutman P., Glamuzina B., Bartulović V. & Dulčić J. 2007. A new maximum length for *Anguilla anguilla* (Anguillidae). *Cybium* 31 (4): 485–486.
- Tutman P., Glamuzina B. & Dulčić J. 2008a. Threatened fishes of the world, *Salmo dentex* (Heckel 1852) (Salmonidae). *Environmental Biology of Fishes* 82 (2): 175–176. <https://doi.org/10.1007/s10641-007-9273-y>
- Tutman P., Glamuzina B. & Dulčić J. 2008b. Threatened fishes of the world, *Chondrostoma knerii* Heckel, 1843 (Cyprinidae). *Environmental Biology of Fishes* 83 (2): 209–210. <https://doi.org/10.1007/s10641-007-9320-8>
- Tutman P., Dulčić J. & Glamuzina B. 2009a. First record of Po brook lamprey, *Lethenteron zanandreae* (Cephalaspidomorphi, Petromyzontiformes, Petromyzontidae) in the Hutovo blato wetland, Bosnia and Herzegovina. *Acta Ichthyologica et Piscatoria* 39 (1): 55–58. <https://doi.org/10.3750/AIP2009.39.1.11>
- Tutman P., Glamuzina B., Dulčić J., Bartulović V. & Hasković E. 2009b. Endemic fish fauna of the Hutovo Blato wetland (Neretva River basin, Bosnia and Herzegovina) and their conservation status. In: Kontautas A. (ed.) *13th European Congress of Ichthyology*: 69. Klaipedos universitetas, Klaipeda.
- Tutman P., Glamuzina B., Dulčić J. & Zovko N. 2012a. Ihtiofauna močvare Hutovo blato (donji tok rijeke Neretve, Bosna i Hercegovina): stanje i ugroženost. *Croatian Journal of Fisheries (Ribarstvo)* 70 (4): 169–185.
- Tutman P., Burić M. & Skaramuca B. 2012b. First substantiated record of the black-striped pipefish, *Syngnathus abaster* (Actinopterygii, Syngnathiformes, Syngnathidae) in the freshwaters of Bosnia and Herzegovina. *Acta Ichthyologica et Piscatoria* 42 (3): 259–262. <https://doi.org/10.3750/AIP2011.42.3.11>
- Tutman P., Čaleta M., Glamuzina B. & Dulčić J. 2012c. Biološko-ekološke osobitosti, rasprostranjenost i stanje zaštite peškelja, *Scardinius plotizza* (Heckel i Kner, 1858) (Pisces, Cyprinidae) na području močvare Hutovo blato u Bosni i Hercegovini. *Croatian Journal of Fisheries (Ribarstvo)* 70 (Supplement 1): 15–28.
- Tutman P., Šanda R., Glamuzina B. & Dulčić J. 2013. First confirmed record of *Pomatoschistus canestrinii* (Ninni, 1883) (Gobiidae) from Bosnia and Herzegovina. *Journal of Applied Ichthyology* 29: 937–939. <https://doi.org/10.1111/jai.12226>

- Tutman P., Freyhof J., Dulčić J., Glamuzina B. & Geiger M. 2017a. *Lampetra soljani*, a new brook lamprey from the southern Adriatic Sea basin (Petromyzontiformes: Petromyzontidae). *Zootaxa* 4273: 531–548. <https://doi.org/10.11646/zootaxa.4273.4.4>
- Tutman P., Buj I., Čaleta M., Hamzić A., Korjenić S., Adrović A. & Glamuzina B. 2017b. Status and distribution of spined loaches (Cobitidae) and stone loaches (Nemacheilidae) in Bosnia and Herzegovina. *Folia Zoologica* 66 (4): 211–227. <https://doi.org/10.25225/fozo.v66.i4.a2.2017>
- Tutman P., Hamzić A., Džano A. & Dulčić J. 2017c. The first record of non-native largemouth black bass *Micropterus salmoides* (Actinopterygii: Perciformes: Centrarchidae), in Bosnia and Herzegovina. *Acta Ichthyologica et Piscatoria* 47 (3): 317–320. <https://doi.org/10.3750/AIEP/02187>
- Tutman P., Matić-Skoko S., Hamzić A., Dulčić J. & Glamuzina B. 2018. Life history traits of Neretva roach *Rutilus basak* (heckel, 1843) (Pisces, Cyprinidae): biological and ecological contribution for biodiversity conservation of freshwater fish. *Croatian Journal of Fisheries* 76 (2): 66–71. <https://doi.org/10.2478/cjf-2018-0008>
- Tutman P., Glamuzina B., Matić-Skoko S., Dulčić J., Pavličević J. & Hamzić A. 2019. An outline of the biology, distribution, and conservation of Neretva nase, *Chondrostoma knerii* (Actinopterygii: Cypriniformes: Cyprinidae), endemic to the Neretva River Basin (Croatia and Bosnia and Herzegovina). *Acta Ichthyologica et Piscatoria* 49 (2): 147–157. <https://doi.org/10.3750/AIEP/02517>
- Tutman P., Buj I., Čaleta M., Marčić Z. & Adrović A. 2020a. Review of the lampreys (Petromyzontidae) in Bosnia and Herzegovina: a current status and geographic distribution. *Journal of Vertebrate Biology* 69 (1): 1–13. <https://doi.org/10.25225/jvb.19046>
- Tutman P., Zanella D., Horvatić S., Hamzić A., Adrović A., Dulčić J. & Glamuzina B. 2020b. Freshwater gobies (Gobiidae) of Bosnia and Herzegovina: a review of the current status and distribution. *Journal of Vertebrate Biology* 69 (49): 1–15. <https://doi.org/10.25225/jvb.20046>
- Tutman P. & Glamuzina B. 2021. Detection and confirmation of freshwater blenny *Salaria fluviatilis* (Actinopterygii: Blenniidae) in Bosnia and Herzegovina. *Croatian Journal of Fisheries* 79: 75–82. <https://doi.org/10.2478/cjf-2021-0009>
- Tutman P., Dragičević B., Dulčić J., Glamuzina L., Bukvić V. & Vekić J. 2021. Unusual records of marine organisms in the Neretva River (Croatia and Bosnia and Herzegovina). *Acta Adriatica* 62 (2): 139–148. <https://doi.org/10.32582/aa.62.2.2>
- Tutman P., Kennedy B., Matić-Skoko S., Ugarković P., Hamzić A., Bolotin J. & Glamuzina B. 2023. Unusual findings of softmouth trout *Salmo obtusirostris* (Actinopteri: Salmonidae) in the marine environment along the middle Eastern Adriatic coast as a consequence of extreme weather events. *Naše more* 70 (3): 171–176. <https://doi.org/10.17818/NM/2023/SI5>
- Uršić S. 1975. *Morfološke i Taksonomske Karakteristike Krkuš – Gobio gobio L. 1758 Sa Gornjeg toka Rijeke Bosne*. Diploma thesis. University of Sarajevo, Sarajevo.
- Van der Laan R., Fricke R. & Eschmeyer W.N. (eds). 2025. Eschmeyer's catalog of fishes: classification. Available from <https://www.calacademy.org/scientists/catalog-of-fishes-classification/> [accessed 3 Mar. 2025].
- Vegara M., Muhamedagić S., Hamzić A., Katica V., Savić M., Đug S., Omanović-Miklićanin E., Lelo S., Drešković N., Muminović M. & Šljuka S. 2009. *Mid-term Improvement Program of Fisheries for the Fishing Area of Kalinovik Municipality*. Norwegian University of Life Sciences (UMB), Sarajevo, NORAGRIC, Aas.
- Veladžić M., Džaferović A., Bećiraj A., Makić H., Dekić R. & Dedić S. 2017. Heavy Metals in Water and Muscle Tissue of Trout (*Salmo trutta*) in the River Una, *Technologica Acta* 10 (1): 45–50.

- Veledar I. 1971. *Određivanje Starosti Podbile (Chondrostoma phoxinus Heckel 1843) iz Voda Livanjskog Polja*. Diploma thesis. University of Sarajevo, Sarajevo.
- Veledar I. & Kosorić Đ. 1972. Dužinsko i težinsko rastenje podbile (*Chondrostoma phoxinus* Heck. 1843) iz voda Livanjskog polja. *Ichthyologia* 4 (1): 89–95.
- Veledar I. & Kosorić Đ. 1974. Metodi određivanja starosti podbile (*Chondrostoma phoxinus* Heck. 1843) iz voda Livanjskog polja. *Ichthyologia* 6 (1): 93–109.
- Veljović P. 1982. Beitrag zum Studium der Ichthyofauna des Save flusses. *Internationale Arbeitsgemeinschaft Donauforschung der Societes Internationalis Limnologie, Wien*: 210–213.
- Veljović P. 1985. Spektar ishrane *Esox lucius* L., *Stizostedion lucioperca* L. i *Silurus glanis* L. u srednjem toku reke Save. *Ribarstvo Jugoslavije* 2–3: 51–55.
- Vesnić A., Gajević M., Mušović A., Đug S., Drešković N., Korjenić E. & Škrijelj R. 2018. Trophic interactions of *Lepomis gibbosus* (Linnaeus, 1758) (Actinopterygii: Perciformes) in Jablanica reservoir on the river Neretva. *Works of the Faculty of Agriculture University of Sarajevo* 68 (1): 66–69.
- Vicković Đ. 1979. *Diferencijalna Krvna Slika Mužjaka i Ženki Kalifornijske Pastrmke (Salmo gairdneri) iz Ribogojilišta Blagaj*. Diploma thesis. University of Sarajevo, Sarajevo.
- Višnjčić Ž., Jarić I., Jovanović Lj., Škorić S., Smederevac-Lalić M., Nikčević M. & Lenhardt M. 2010. Heavy metal and trace element accumulation in muscle, liver and gills of the Pontic shad (*Alosa immaculata* Bennet 1835) from the Danube River (Serbia). *Microchemical Journal* 95: 341–344. <https://doi.org/10.1016/j.microc.2010.02.004>
- Vlasenko A.D., Pavlov A.V., Sokolov L.I. & Vasil'ev V.P. 1989. *Acipenser gueldenstaedti* Brandt, 1833. In: Holčík J. (ed.) *The Freshwater Fishes of Europe: Vol 1, Part II, General Introduction to Fishes, Acipenseriformes*: 294–344. AULA-Verlag, Wiesbaden.
- Volić J. 1984. *Eksperimentalna Hibridizacija Oštrulja (Aulopyge hugeli, Heckel, 1843) i Sitnojstavog Kljena (Leuciscus turskyi, Heckel, 1843) iz Buškog Blata*. Diploma thesis. University of Sarajevo, Sarajevo.
- Vucić M., Sučić I. & Jelić D. 2017. New distribution data for *Alburnus sava* Bogutskaya, Zupančić, Jelić, Diripasko & Naseka, 2017 and *Telestes souffia* (Risso, 1827) in the Western Balkans. *Croatian Journal of Fisheries* 75: 137–142. <https://doi.org/10.1515/cjf-2017-0017>
- Vucić M., Jelić D., Žutinić P., Grandjean F. & Jelić M. 2018. Distribution of Eurasian minnows (*Phoxinus*: Cypriniformes) in the Western Balkans. *Knowledge & Management of Aquatic Ecosystems* 419: 1–11. <https://doi.org/10.1051/kmae/2017051>
- Vujanović T., Radević M., Manojlović M., Ivanc A. & Dekić R. 2011. Dužinsko-težinski odnosi kod skobalja (*Chondrostoma nasus*). *Skup 1. Savremena Univerzitetska Nastava* 3: 57–66.
- Vukić J., Eliášová K., Marić D. & Šanda R. 2019. Occurrence of alien spiralin (*Alburnoides* sp.) in the Neretva river basin. *Knowledge & Management of Aquatic Ecosystems* 420: 15. <https://doi.org/10.1051/kmae/2019007>
- Vuković D., Tursi A., Carlucci R. & Dekić R. 2008. Ichthyofauna of the wetland ecosystem in the Bardača area. *Ribarstvo* 66: 89–103.
- Vuković N. 1979. Morphological-taxonomic characteristics of *Leuciscus souffia agassizi* (Cyprinidae, Pisces) from the river Drina. In: Kullander S. (ed.) *Third European Ichthyological Congress*: 240–240. EIC, Warsaw.
- Vuković N. 1982a. The external brain morphology of some Salmonoid fish species in Bosnia and Herzegovina. *Godišnjak Biološkog instituta Univerziteta Sarajevo* 43: 3–152.

- Vuković N. 1985. *Ekološke i Biosistematske Karakteristike Leuciscus souffia Risso, 1826 iz Gornjeg toka Rijeke Drine*. PhD thesis. University of Sarajevo, Sarajevo.
- Vuković N. 1986. Ishrana jelšovke (*Leuciscus souffia agassizi* Valen. 1844) iz gornjeg toka reke Drine. *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 39: 175–187.
- Vuković N. 1987. Morpho-taxonomic characteristics *Leuciscus souffia agassizi* Valenciennes, 1844, from the upper reaches of the river Drina. *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 40: 117–136.
- Vuković N. 1990. Morpho-taxonomic characteristics *Leuciscus souffia agassizi* Valenciennes from the upper reaches of the river Drina. *Godišnjak Biološkog Instituta Univerziteta Sarajevo* 43: 63–82.
- Vuković N. & Vuković T. 1974. Morfologija mozga nekih ciprinida iz voda Jugoslavije. *Ichthyologia* 6 (1): 125–141.
- Vuković N. & Vuković T. 1977. Osobnosti spoljašnje morfologije mozga glavatiće (*Salmo marmoratus* Cuvier, 1817) i mekousne pastrmke (*Salmothymus obtusirostris oxyrhynchus* Steindachner, 1882). *Ichthyologia* 9 (1): 15–20.
- Vuković N., Vuković T. & Sekulović T. 1970a. Težina srca i mozga kod nekih vrsta ciprinida iz Skadarskog jezera i reke Bune. *Ichthyologia* 2 (1): 143–154.
- Vuković T. 1959. Stvaranje godišnjeg prstena na krljuštima klena (*Squalius cephalus* L) iz Izvorskog Dela Reke Bosne. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 12 (1–2): 193–199.
- Vuković T. 1960. Die Bildung des Jahresringes an den Schuppen des Döbels (*Squalius cephalus* L.) aus dem Quellgebiet der Bosna. *Bulletin Scientifique* 5 (3): 76.
- Vuković T. 1961. Populacije i mrešćenje *Alosa falax nilotica* (Geoffroy) u vodama Neretve i Skadarskog jezera. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 14 (1–2): 85–178.
- Vuković T. 1962. Novi podaci o prezimljavanju u slatkoj vodi i rastenju mladih ćepa (*Alosa falax nilotica* (Geoffroy) u vodama donje Neretve. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 15 (1–2): 141–145.
- Vuković T. 1963a. *Ribe Bosne i Hercegovine (Ključ za Određivanje)*. Zavod za izdavanje udžbenika, Sarajevo.
- Vuković T. 1963b. Prilog poznavanju rasprostranjenja *Leuciscus souffia* Risso u vodama Jugoslavije i opis podvrste *Leuciscus souffia montenegrinus* n. ssp. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 16 (1–2): 205–207.
- Vuković T. 1963c. Nalaz križanca gagice i sapače (*Phoxinus phoxinus* Linné × *Barbus meridionalis petenyi* Heckel) u okolini Banja Luke. *Ribarstvo Jugoslavije Zagreb* 17 (5): 147–148.
- Vuković T. 1964. Prilog poznavanju prirodne hibridizacije ciprinida u vodama Livanjskog polja. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 17: 199–206.
- Vuković T. 1966a. Dužine crijevnog trakta sapača (*Barbus meridionalis petenyi*, Heck.) i krkuša (*Gobio gobio* Linnaeus) iz različitih lokaliteta. *Ribarstvo Jugoslavije* 21 (6): 132–134.
- Vuković T. 1966b. Variranje relativne dužine crevnog trakta kod gagica (*Phoxinus phoxinus* Linné) iz različitih voda u Bosni i Hercegovini. *Veterinaria* 15 (4): 523–526.
- Vuković T. 1967a. Analiza nekih taksonomskih karaktera *Phoxinus phoxinus* (Liné) iz Bregave i sliva gornjeg toka rijeke Bune. *Glasnik Zemaljskog Muzeja Bosne i Hercegovine* 6: 127–139.
- Vuković T. 1967b. Nalaz *Paraphoxinus ghetaldi* Stein. u slivu gornjeg toka reke Bosne. *Glasnik Zemaljskog Muzeja Bosne i Hercegovine* 6: 219–222.

- Vuković T. 1968a. Nalaz hibrida *Rutilus rubilio* (Bonaparte) × *Alburnus alburnus alborella* (Filipi) u slivu reke Neretve. *Glasnik Zemaljskog Muzeja Bosne i Hercegovine* 7: 237–241.
- Vuković T. 1968b. Variranje dužine crevnog trakta nekih ciprinidnih vrsta iz pritoka gornjeg toka reke Bosne. *Ribarstvo Jugoslavije* 23 (4): 84–86.
- Vuković T. 1970. Contributo alla cognizione di alcuni caratteri tassonomici delle popolazioni del *Rutilus rubilio* (Bonaparte) [Pisces, Cyprinidae] delle regione confluyente del fiume Po. *Annali del Museo civico di Storia naturale di Genova*: 12–24.
- Vuković T. 1977a. *Ribe Bosne i Hercegovine*. IGKRO „Svjetlost“, Sarajevo.
- Vuković T. 1977b. *Phoxinellus alepidotus* (Heckel, 1843). Novi podaci o rasprostranjenju u vodama Jugoslavije. *Ribarstvo Jugoslavije* 32 (3–4): 15–22.
- Vuković T. 1982b. Sistematika riba. In: Bojčić I. (ed.) *Slatkovodno Ribarstvo*: 99–168. Poslovna zajednica slatkovodnog ribarstva Jugoslavije, Ribozajednica Zagreb, JUMENA, Zagreb.
- Vuković T. & Aganović M. 1971. Analiza morfološko-taksonomskih karakteristika vrste *Rutilus rutilus* Linnaeus (Pisces, Cyprinidae). In: Slavnić Ž. (ed.) *Zbornik referata sa I Simpozijuma Sistematičara Jugoslavije*: 207–213. Sarajevo.
- Vuković T. & Ivanišević B. 1962. Postojanje dviju morfološki različitih populacija *Scardinius erythrophthalmus scardafa* (Bonaparte) u donjoj Neretvi i Skadarskom jezeru. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 15 (1–2): 141–145.
- Vuković T. & Ivanović B. 1970. Taksonomske karakteristike *Rutilus rubilio* (Bonaparte) iz Skadarskog jezera i sliva Neretve. *Ribarstvo Jugoslavije* 25 (1): 18–20.
- Vuković T. & Ivanović B. 1971. *Slatkovodne Ribe Jugoslavije*. Zemaljski Muzej Sarajevo, Sarajevo.
- Vuković T. & Kosorić Đ. 1967. Uzrasno variranje dužine dorzalnih bodlji kod *Gasterosteus aculeatus* Linné (koljuška, bodonja) iz reke Bune. *Ribarstvo Jugoslavije* 22 (1): 7–9.
- Vuković T. & Kosorić Đ. 1978. Efekti introdukcije ribljih vrsta u vode Jugoslavije i mogući uticaji rekonstrukcije ihtiofaune. *Croatian Journal of Fisheries: Ribarstvo* 33 (4): 92–95.
- Vuković T. & Miladinović Ž. 1969. Serološka proučavanja nekih vrsta roda *Paraphoxinus*. *Ichthyologia* 1 (1): 83–88.
- Vuković T. & Prolić F. 1966. Prilog poznavanju taksonomskih karaktera *Gasterosteus aculeatus* Linne iz donje Neretve. *Glasnik Zemaljskog Muzeja Bosne i Hercegovine* 5: 175–178.
- Vuković T. & Seratlić-Savić D. 1967. Broj brahiospina na škržnim lucima ukljeve *Alburnus albidus alborella* (Filippi) i masnice *Rutilus rubilio* (Bonaparte) iz Skadarskog jezera i reke Bune. *Glasnik Zemaljskog Muzeja Bosne i Hercegovine* 6: 211–214.
- Vuković T. & Seratlić-Savić D. 1970. Promjenljivost broja kičmenih pršljenova u populacijama vrsta *Phoxinus phoxinus* (Linnaeus) i *Gobio gobio* (Linnaeus) u vodama Bosne i Hercegovine. In: *III Simpozijum Jugoslavenskog Ihtiološkog Društva*: 27. Yugoslav Ichthyological Society, Kotor.
- Vuković T. & Sofradžija A. 1987. Endemična ihtiofauna Bosne i Hercegovine i problemi njene zaštite. In: Vuković T. (ed.) *Zaštita Endema u Živom Svijetu Jugoslavije*: 39–43. Posebna izdanja Odjeljenja prirodnih i matematičkih nauka, Akademije nauka i umjetnosti Bosne i Hercegovine, Sarajevo.
- Vuković T. & Veleđar I. 1974. Opis krljušti hibrida *Scardinius erythrophthalmus* (Linnaeus, 1758) × *Leuciscus turskyi* (Heckel, 1843). *Croatian Journal of Fisheries: Ribarstvo* 29 (4): 75–78.

- Vuković T. & Vuković N. 1968a. Neke taksonomske karakteristike vrste *Rutilus rubilio* (Bonap.) [Pisces, Cyprinidae] iz voda Jugoslavije i severne Italije. *Godišnjak Biološkog Instituta Univerziteta u Sarajevu* 21: 129–135.
- Vuković T. & Vuković N. 1968b. Dužina crijevnog trakta klijenova (*Leuciscus cephalus* Linnaeus) i pliski (*Alburnoides bipunctatus* Bloch) iz potoka Suturlije. *Ribarstvo Jugoslavije* 23 (6): 147–149.
- Vuković T. & Vuković N. 1970. Proučavanje taksonomskih karakteristika vrste *Rutilus rubilio* (Bonaparte). In: *III simpozijum "Savremena ihtiološka istraživanja"*: 59. Yugoslav Ichthyological Society, Kotor.
- Vuković T. & Žnidaršić-Krzyk S. 1969. Broj eritrocita, broj leukocita i koncentracija hemoglobina u nekih ciprinidnih vrsta riba. *Ribarstvo Jugoslavije* 1: 10–11.
- Vuković T., Šestak V. & Vuković N. 1962. Prilog proučavanju metodike određivanja starosti i porasta klena *Leuciscus cephalus* L. *Veterinaria* 11 (4): 5217–521.
- Vuković T., Seratlić-Savić D. & Karanac V. 1970b. Neke morfološke karakteristike hibrida *Chondrostoma phoxinus* (Heckel) × *Paraphoxinus alepidotus* (Heckel). *Ichthyologia* 2 (1): 155–169.
- Vuković T., Karanac V. & Seratlić-Savić D. 1971a. Description of the new hybrid *Paraphoxinus alepidotus* Heck. × *Scardinius erythrophthalmus* L. from the waters of Livanjsko polje. *Bulletin scientifique, Section A* (16): 11–12.
- Vuković T., Seratlić-Savić D. & Karanac V. 1971b. Opis novog hibrida *Scardinius erythrophthalmus* L. × *Paraphoxinus alepidotus* Heck. *Ichthyologia* 3 (1): 79–88.
- Vuković T., Guzina N. & Grbelja J. 1980. Immunological reactivity of some fish species (Pisces, Cyprinidae) to plant viruses. *Ichthyologia* 12 (1): 97–98.
- Vuković T., Seratlić-Savić D., Guzina N. & Vuković N. 1977. Problemi biosistematskog statusa vrste *Salmo marmoratus* Cuvier, 1817. In: *Naučni Skup "Struktura, Dinamika i Zaštita Ekosistema i Njihovih Komponenta na Dinaridima"*. Institute of Biology, University of Sarajevo, Sarajevo.
- Vuković T., Grbelja J., Guzina N. & Juretić N. 1979. Imunološka reaktivnost babuške, *Carassius auratus gibelio* (Bloch, 1783) na virus žutog mozaika postrne repe (turnip yellow mosaic virus) i virus šarenila velebilja (belladonna mottle virus). *Ichthyologia* 11 (1): 57–61.
- Vuković Teslić B. 1950. Iz ribarstva NR Bosne i Hercegovine. *Ribarstvo Jugoslavije* 9: 214–215.
- Vuletić A. 1969. *Chondrostoma Nasus sa Ušća Rijeke Bosne*. Diploma thesis. University of Sarajevo, Sarajevo.
- Winterhalter M., Jadrić S. & Perić B. 1971. Elektroforetska analiza izoenzima LDH u oštrulje (*Aulopyge hügelii* Heck.) na gelu akrilamida. *Ichthyologia* 3 (1): 89–93.
- Weiss S. & Schenekar T. 2016. Genetic evaluation of the self-sustaining status of a population of the endangered Danube salmon, *Hucho hucho*. *Hydrobiologia* 775: 153–165. <https://doi.org/10.1007/s10750-016-2726-6>
- Weiss S., Marić, S. & Snoj A. 2011. Regional structure despite limited mtDNA sequence diversity found in the endangered Huchen, *Hucho hucho* (Linnaeus, 1758). *Hydrobiologia* 658: 103–110. <https://doi.org/10.1007/s10750-010-0453-y>
- Wisniewski L.W. 1932. *Cyatocephalus truncatus* Pallas I. Die Postembryonalentwicklung und Biologie. *Bulletin International de l'Academie Polonaise des Sciences et des Lettres: Classe des Sciences Mathematiques et Naturelles, Serie B*: 237–252.

- Witkowski A., Bajić A., Treer T., Hegediš A., Marić S., Šprem N., Piria M. & Kapusta A. 2013. Past and present of and perspectives for the Danube huchen, *Hucho hucho* (L.), in the Danube basin. *Archives of Polish Fisheries* 21: 129–142. <https://doi.org/10.2478/aopf-2013-0010>
- Zanandrea G. 1959. Lamprede parasite e non parasite nel bacino del Danubio e la nuova entita sistematica *Eudontomyzon danfordi vladykovi*. *Archivio Zoologico Italiano* 44: 215–250.
- Zanella D., Mrakovčić M., Schneider D., Mustafić P., Čaleta M. & Radić I. 2003. Growth of *Cobitis narentana*, 1928 in the Neretva River, Croatia. *Folia Biologica (Kraków)* 51: 154–157.
- Zanella D., Mrakovčić M., Mustafić P., Čaleta M. & Marčić Z. 2009. Threatened fishes of the world, *Cobitis narentana* Karaman 1928 (Cobitidae). *Environmental Biology of Fishes* 86 (4): 523–524. <https://doi.org/10.1007/s10641-009-9559-3>
- Zanella D., Marčić Z., Čaleta M., Buj I., Zrnčić S., Horvatić S. & Mustafić P. 2017. Early development of the freshwater goby *Orsinogobius croaticus* endemic to Croatia and Bosnia–Herzegovina. *Cybium* 41 (4): 335–342.
- Zanella L.N., DeFaveri J., Zanella D., Merilä J., Šanda R. & Mrakovčić M. 2015. Does predation drive morphological differentiation among Adriatic populations of the three-spined stickleback? *Biological Journal of the Linnean Society* 115: 219–240. <https://doi.org/10.1111/bij.12491>
- Zangl L., Daill D., Gessl W., Friedrich T. & Koblmüller S. 2020. Austrian gudgeons of the genus *Gobio* (Teleostei: Gobionidae): A mixture of divergent lineages. *Journal of Zoological Systematics and Evolutionary Research* 58: 327–340. <https://doi.org/10.1111/jzs.12340>
- Zaplata R. 1932. Platnica (*Leuciscus virgo* Heckel). *Ribarski list* 9–10: 102–104.
- Zaplata R. & Taler Z. 1932. Ribe Sarajeva i okoline. *Glasnik Zemaljskog muzeja u Bosni i Hercegovini* 44 (1): 1–32.
- Zorlak Dž. 2005. *Analiza Morfološko-taksonomskih Karakteristika Vrste Gobio gobio (Linnaeus, 1758) iz Rijeke Save*. Diploma thesis. University of Sarajevo, Sarajevo.
- Zrnčić S., Oraić D., Šoštarić B., Čaleta M., Buj I., Zanella D. & Šurmanović D. 2009. Occurrence of parasites in Cobitidae from Croatian rivers draining into two different watersh. *Journal of Applied Ichthyology* 25: 447–450. <https://doi.org/10.1111/J.1439-0426.2009.01192.X>
- Zuliani T., Vidmar J., Drinčić A., Ščančar J., Horvat M., Nečemer M., Piria M., Simonović P., Paunović M. & Milačić R. 2019. Potentially toxic elements in muscle tissue of different fish species from the Sava River and risk assessment for consumers. *Science of the Total Environment* 650: 958–969. <https://doi.org/10.1016/j.scitotenv.2018.09.083>
- Zupančić P. 2008. *Rijetke i Ugrožene Slatkovodne Ribe Jadranskog Slijeva Hrvatske, Slovenije i Bosne i Hercegovine*. Narodna in univerzitetna knjižnica, Ljubljana.
- Zupančić P. & Bogutskaya N.G. 2000. Description of a new species, *Phoxinellus dalmaticus* (Cyprinidae, Leuciscinae): from the Čikola River in the Krka River system, Adriatic basin (Croatia). *Natura Croatica* 9 (2): 67–81.
- Zupančić P. & Bogutskaya N.G. 2002. Description of two new species, *Phoxinellus krbavensis* and *P. jadovensis*, re-description of *P. fontinalis* Karaman 1972, and discussion of the distribution of *Phoxinellus* species (Teleostei, Cyprinidae) in Croatia and Bosnia and Herzegovina. *Natura Croatica* 11 (4): 411–437.
- Žitnan R., Kiškaroļj M. & Čanković M. 1969. Epizootiološki odnosi parazitoza na ribama jednog šaranskog gazdinstva u Bosni i Hercegovini. *Ichthyologia* 8 (1): 99–108.
- Žujo D. 2004. *Biosistematska Istraživanja Vrste Chondrostoma phoxinus Heckel, 1843 iz Buškog jezera*. Master's thesis, University of Sarajevo, Sarajevo.

- Žujo-Zekić D. 2012. The nourishment of the endemic species *Chodrostoma phoxinus* Heckel, 1843 (Teleostei: Ostaryophysii, Cyprinidae) in the waters of Buško Lake (Bosnia and Herzegovina). In: Milošević D. (ed.) *Zbornik radova XVII Savetovanja o Biotehnologiji sa Međunarodnim Učešćem, Čačak, Srbija* 17 (19): 337–341. Agronomski fakultet u Čačku, Čačak.
- Žujo-Zekić D. 2013. Age structure of population of endemic species *Chondrostoma phoxinus* Heckel, 1843 (Teleostei: Ostaryophysii, Cyprinidae) from Buško Lake. In: Petrović M. & Poleksić V. (eds) *VI International Conference “Water & Fish”, Conference Proceedings*: 331–341. Faculty of Agriculture, Belgrade-Zemun.
- Žujo-Zekić D. 2021. *Ihtiološka raznolikost Buškog jezera, jučer – danas – sutra*. Univerzitet “Džemal Bijedić” u Mostaru, organizacijska jedinica Nastavnički fakultet, Mostar.
- Žujo-Zekić D., Kašić-Lelo M., Lelo S. & Abaza A. 2016. Sex ratio in local population of species *Chondrostoma phoxinus* Heckel, 1843 (Teleostei: Ostaryophysii, Cyprinidae) from Buško Lake. *Proceedings of the VII International Scientific Agriculture Symposium “Agrosym 2016”*: 2043–2051. University of East Sarajevo, Faculty of Agriculture, Bosnia and Herzegovina, Jahorina. <https://doi.org/10.7251/AGRENG1607306>
- Žujo-Zekić D., Ademović E., Riđanović S., Riđanović L., Spasojević P. & Boškailo A. 2018. Biodiversity of the Buško Lake freshwater fish as a part of diversity of ichthyofauna in the karst fields of Bosnia and Herzegovina. *Proceedings of the IX International Agricultural Symposium “Agrosym 2018”*: 1410–1417. University of East Sarajevo, Faculty of Agriculture, Bosnia and Herzegovina, Jahorina.
- Žujo-Zekić D., Riđanović S. & Spasojević P. 2020. First finding of species *Alburnus arborella* (Bonaparte, 1841) syn. *Alburnus alborella* (De Filippi, 1844) (Actinoptery GII: Cyprinidae, Alburninae) in Buško Lake. *Journal of Survey in Fisheries Sciences* 6 (2): 79–92. <https://doi.org/10.18331/SFS2020.6.2.9>
- Žuljević S. 1967. *Neke taksonomske karakteristike Alburnus albidus arborella donje Neretve*. Diploma thesis. University of Sarajevo, Sarajevo.
- Žuljević M. 1970. *Neke Taksonomske Karakteristike Alburnoides bipunctatus (Bloch) Rijeka Crnomorskog Sliva Bosne i Hercegovine*. Diploma thesis. University of Sarajevo, Sarajevo.

Printed versions of all papers are deposited in the libraries of two of the institutes that are members of the *EJT* consortium: Muséum national d’Histoire naturelle, Paris, France and Royal Museum for Central Africa, Tervuren, Belgium. The other members of the consortium are: Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Meise Botanic Garden, Meise, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic; The Steinhardt Museum of Natural History, Tel Aviv, Israël.

Supplementary file

Supp. file 1. List of all lampreys and other fish species recorded in freshwaters in Bosnia and Herzegovina according to literature to date. <https://doi.org/10.5852/ejt.2026.1042.3201.14209>