















Received: 2 September 2025 • Accepted: 13 February 2026 • Published: 10 April 2026

Topic editor: Magalie Castelin • Section editor: Arnaud Henrard • Desk editor: Pepe Fernández

### Research article

[urn:lsid:zoobank.org:pub:362F99E7-9407-44AD-A608-D4FEDEA33657](https://zoobank.org/pub:362F99E7-9407-44AD-A608-D4FEDEA33657)

## Discoveries of two new *Mogrus* species (Araneae: Salticidae) and notable records from India

Rishikesh TRIPATHI<sup>1,\*</sup>, Gautam KADAM<sup>2</sup>, Theresa Joy ASHA<sup>3</sup>,  
Atharva KULKARNI<sup>4</sup>, Ashish Kumar JANGID<sup>5</sup>,  
Dhruv A. PRAJAPATI<sup>6</sup> & Ambalaparambil Vasu SUDHIKUMAR<sup>7</sup>

<sup>1,3,7</sup>Centre for Animal Taxonomy and Ecology, Department of Zoology, Christ College, Irinjalakuda, Thrissur, Kerala 680 125, India.

<sup>1</sup>Wildlife Conservation Trust, 11<sup>th</sup> floor, Mafatlal Centre, Nariman Point, Mumbai, Maharashtra 400021, India.

<sup>2</sup>Division of Arachnology, Department of Zoology, Sacred Heart College, Thevara, Cochin, Kerala 682 013, India.

<sup>4</sup>Snehal Kunj Apartment, Sant Eknath Nagar, Bibwewadi, Pune, Maharashtra 411037, India.

<sup>5</sup>Wildlife Institute of India, Dehradun, Uttarakhand 248 001, India.

<sup>6</sup>Sardar Patel Zoological Park, Statue of Unity Road, Ektanagar, Gujarat 393151, India.

\*Corresponding author: [rishikesh.tripathi14@gmail.com](mailto:rishikesh.tripathi14@gmail.com)

<sup>2</sup>Email: [gautamkadam7wild@gmail.com](mailto:gautamkadam7wild@gmail.com)

<sup>3</sup>Email: [ashatheresajoy@gmail.com](mailto:ashatheresajoy@gmail.com)

<sup>4</sup>Email: [atharvakulkarni1919@gmail.com](mailto:atharvakulkarni1919@gmail.com)

<sup>5</sup>Email: [ashishjangid22@gmail.com](mailto:ashishjangid22@gmail.com)

<sup>6</sup>Email: [dhruvspidy215@gmail.com](mailto:dhruvspidy215@gmail.com)

<sup>7</sup>Email: [spidersudhi@gmail.com](mailto:spidersudhi@gmail.com)

**Abstract.** This study provides taxonomic contributions to the genus *Mogrus* Simon, 1882 from India, with the descriptions of two new species: *Mogrus shushka* Tripathi, Kadam & Prajapati sp. nov. and *Mogrus pune* Tripathi, Kulkarni & Kadam sp. nov. Furthermore, the male of *Mogrus rajasthanensis* Caleb, Chatterjee, Tyagi, Kundu & Kumar, 2017 is described for the first time, while *Mogrus larisae* Logunov, 1995 is newly reported from India. We provide detailed morphological descriptions and images, as well as discuss variations in female genitalia within the genus.

**Keywords.** Jumping spiders, genitalia variation, semi-arid ecosystems, species record, taxonomy.

Tripathi R., Kadam G., Asha T.J., Kulkarni A., Jangid A.K., Prajapati D.A. & Sudhikumar A.V. 2026. Discoveries of two new *Mogrus* species (Araneae: Salticidae) and notable records from India. *European Journal of Taxonomy* 1048: 223–244. <https://doi.org/10.5852/ejt.2026.1048.3249>

## Introduction

*Mogrus* Simon, 1882 is a genus of medium-sized jumping spiders widely found in arid and semi-arid regions of Africa, Europe, the Middle East, and Asia (WSC 2026). The genus is mostly found in southern Palaeartic regions, ranging from the Mediterranean to the Arabian Peninsula to Mongolia and western China, with additional occurrences documented in South Asia (Andreeva *et al.* 1981; Caleb *et al.* 2017; Kanesharatnam & Benjamin 2018).

*Mogrus* has been revised twice since its initial description (Andreeva *et al.* 1981; Logunov 1995), with an earlier classification in the subfamily Pelleninae Petrunkevitch, 1928, and later in Plexippinae Petrunkevitch, 1928 by Prószyński (1976). However, this categorisation was somewhat contested based on morphological discrepancies, especially the lack of a membranous conductor in *Mogrus* male palps compared to kindred genera (Logunov 1995). Molecular phylogenetic research has placed *Mogrus* within the subfamily Salticinae Blackwall, 1841 (Maddison 2015). Further refinement grouped them within the supergroup Hylloida Prószyński, 2016 characterized by a long embolus that originates laterally on the bulbus and parallels its margin, an important diagnostic trait for the genus (Prószyński 2016). In this paper, we describe and illustrate two new species of the genus *Mogrus* from India, and reporting the expanded distribution of the genus.

## Material and methods

All measurements are in millimeters (mm). Lengths of palp and leg segments are given as: total (femur, patella, tibia, metatarsus (except for palp), tarsus). The micrographic images were taken with a Leica DFC500 digital camera attached to a Leica M205A stereo microscope with the software package Leica Application Suite (LAS ver. 3.8) for stacking images taken at different focal planes. The terminology for genitalia mostly follows Logunov (1995) and Tripathi *et al.* (2023). The types and studied materials are deposited at NRC-AA (NCBS Research Collection, National Centre for Biological Sciences, Bengaluru, India) and the reference collection of Web of Nature (WON) Research Foundation, Gujarat, India.

## Abbreviations used for morphology

ALE = anterior lateral eye  
AME = anterior median eye  
PLE = posterior lateral eye  
PME = posterior median eye  
RTA = retrolateral tibial apophysis  
I–IV = 1<sup>st</sup> to 4<sup>th</sup> leg

## Acronyms of repositories

ISEN = Zoological Museum, Institute for Systematics and Ecology of Animals, Novosibirsk, Russia  
NCBS = National Centre for Biological Sciences, Bengaluru, Karnataka, India  
NZC-ZSI = National Zoological Collections, Zoological Survey of India, Kolkata, India  
WON = Web of Nature Research Foundation, Gujarat, India  
ZSI-CDT = Centre for DNA Taxonomy, Zoological Survey of India, Kolkata, India

## Results

### *Keys to Indian species of Mogrus*

1. Males ..... 2  
– Females ..... 5
2. Protuberance of bulb triangular, situated prolaterally, 11 o'clock position .....  
..... *M. shushka* Tripathi, Kadam & Prajapati sp. nov.  
– Protuberance of bulb heart-shaped, situated retrolaterally, 12–2 o'clock position ..... 3
3. RTA long, broad at base, gradually narrowing, bent tip in ventral view .....  
..... *M. pune* Tripathi, Kulkarni & Kadam sp. nov.  
– RTA moderately long, arched dorsally, rounded tip in retrolateral view .....  
..... *M. rajasthanensis* Caleb, Chatterjee, Tyagi, Kundu & Kumar, 2017
4. RTA slender, extending from base to distal end with nearly parallel margins, terminating in a flattened tip ..... *M. larisae* Logunov, 1995  
– RTA long, gradually tapering toward the tip, slightly bent in ventral view ..... 5
5. Epigynal depression deep, semi-circular-shaped, occupying central region of epigyne .....  
..... *M. shushka* sp. nov.  
– Epigynal depression shallow, heart-shaped concavity that tapers anteriorly ..... 6
6. Copulatory openings large, situated mediolaterally ..... *M. pune* sp. nov.  
– Copulatory openings small, situated posterolaterally ..... 7
7. Copulatory ducts distinct and small; spermathecae circular ..... *M. rajasthanensis*  
– Copulatory ducts indistinct, large, and spermathecae C-shaped ..... *M. larisae*

### *New species*

Class Arachnida Lamarck, 1801  
Order Araneae Clerck, 1757  
Family Salticidae Blackwall, 1841  
Subfamily Salticinae Blackwall, 1841  
Tribe Salticini Blackwall, 1841  
  
Genus *Mogrus* Simon, 1882

### **Type species**

*Mogrus fulvovittatus* Simon, 1882, by original designation.

### **Diagnosis**

For diagnostic features of the genus, see Andreeva *et al.* (1981) and Logunov (1995).

### **Distribution**

Africa, the Middle East, Central Asia, Europe, and South Asia.

*Mogrus shushka* Tripathi, Kadam & Prajapati sp. nov.  
[urn:lsid:zoobank.org:act:EF44D5D2-8082-48A7-B24B-233F4AB5E4BB](https://doi.org/10.21203/rs.3.rs-1234567/v1)

Figs 1–4

### Diagnosis

Males of *M. shushka* sp. nov. are similar to those of *M. cognatus* Wesolowska & van Harten, 1994, in having a long and broad cymbium, a protuberance positioned slightly at the prolateral side but can be distinguished by a triangular-shaped protuberance (vs thumb-shaped in *M. cognatus*); RTA gradually narrows towards the tip (vs pronounced curve and extended tip in *M. cognatus*) (compare Figs 3A–C, 4A–B with Wesolowska & van Harten 1994: figs 112–114; 2010: figs 37–39). Females are similar to those of *M. macrocephalus* Lawrence, 1927 in having a centrally positioned depression; copulatory ducts membranous, initially broad, gradually narrowing and becoming coiled but can be distinguished from the latter by differences in the shape of the depression and the coiling of the copulatory duct. Fertilization ducts are short and closely spaced (vs large and widely spaced in *M. macrocephalus*) compare Figs 3D–H, 4C–D with Azarkina & Foord (2013: figs 51–53).


### Etymology

The specific epithet ‘*shushka*’ is a noun in apposition, derived from the Sanskrit word *Śuṣka*, meaning ‘dry/arid’. It refers to the species’ distribution in the drylands, highlighting its habitat preference for arid and open natural ecosystems.

### Type material

#### Holotype


INDIA • ♂; Rajasthan, Beawar, Chhatrasagar, Nimaaj; 26°06'32.2" N, 74°01'08.6" E; elev. 320 m; 28 Sep. 2024; R. Tripathi and G. Kadam leg.; found on a tree branch, by hand; NCBS, NRC-AA-0640.




**Fig. 1.** Field photographs of *Mogrus shushka* Tripathi, Kadam & Prajapati, 2026 sp. nov. **A–D.** Holotype, ♂ (NRC-AA-0640). **E–F.** Paratype, ♀ (NRC-AA-0641).

**Paratypes**


INDIA • 2 ♀♀; same data as for holotype; NCBS, NRC-AA-0641, NRC-AA-0642 • 1 ♂, 1 ♀; Rajasthan, Jaisalmer, Desert National Park, Loharki area; 27°08'21.9" N, 71°45'58.2" E; elev. 234 m; 20 Aug. 2022; R. Tripathi leg.; NCBS, NRC-AA-0643, NRC-AA-0644 • 1 ♂, 2 ♀♀; Gujarat, Kutch, Chaari; 23°33'45.02" N, 69°24'44.10" E; elev. 28 m; 7 Dec. 2023; D. Prajapati leg.; found on a tree branch, by hand; WON104360A.



**Fig. 2.** *Mogrus shushka* Tripathi, Kadam & Prajapati sp. nov. A–C. Holotype, ♂ (NRC-AA-0640). D–F. Paratype, ♀ (NRC-AA-0641). A, D. Habitus, dorsal view. B, E. Same, lateral view. C, F. Same, frontal view. Scale bars = 2 mm.



**Fig. 3.** *Mogrus shushka* Tripathi, Kadam & Prajapati sp. nov., genitalia. A–C. Holotype, ♂ (NRC-AA-0640). D–E. Paratype, ♀ (NRC-AA-0641). F. Paratype, ♀ (NRC-AA-0642). G. Paratype, ♀ (NRC-AA-0643). H. Paratype, ♀ (NRC-AA-0644). A–C. Male left palp. A. Ventral view. B. Retrolateral view. C. Dorsal view. D–H. Epigyne/vulva. D. Ventral view. E–H. Dorsal view. Scale bars = 0.2 mm.



**Fig. 4.** *Mogrus shushka* Tripathi, Kadam & Prajapati sp. nov., genitalia. **A–B.** Holotype, ♂ (NRC-AA-0640). **C–D.** Paratype, ♀ (NRC-AA-0641). **A–B.** Male left palp. **A.** Ventral view. **B.** Retrolateral view. **C–D.** Epigyne/vulva. **C.** Ventral view. **D.** Dorsal view. Scale bars = 0.2 mm.

## Description

### Male (holotype, Figs 1A–D, 2A–C)

COLOURATION IN ALCOHOL. Carapace, eye region, clypeus, chelicerae, endites, labium, legs, opisthosoma, sternum chocolate brown (Fig. 2A–C). Carapace with two prominent, nearly uniform white bands originating behind AMEs, extending posteriorly, and merging at the anterior part to form distinct U-shape (Figs 1A, 2A). Eye field with recumbent black flattened setae reflects in alcohol; anterior eyes encircled by tiny dirt whitish hairs (Fig. 2A, C). White hairs cover clypeal region; thin brown stripes arise beneath ALE extending laterally (Fig. 2C). Abdomen ovoid, with dark brown median region featuring chevron-shaped markings, bordered by thick white longitudinal stripes on either side (Fig. 2A).

MEASUREMENTS. Total length: 7.28. Carapace: length 3.52, width 2.57. Abdomen: length 3.76, width 2.21. Length of chelicerae: 1.73. Clypeus height: 0.18. Eye diameters: AME 0.61, ALE 0.31, PME 0.09, PLE 0.31; AME–AME: 0.05. Eye interdistances: AME–ALE 0.17, PME–PLE 0.36, ALE–ALE 1.66, PME–PME 2.12, ALE–PME 0.37, PLE–PLE 2.16. Measurements of palp and legs: palp: 2.88 (1.04, 0.35, 0.28, 1.21), leg I 7.66 (2.18, 1.26, 2.01, 1.42, 0.79), II 6.17 (1.88, 1.04, 1.49, 1.04, 0.72), III 6.12 (2.02, 0.86, 1.41, 1.18, 0.65), IV 5.98 (1.92, 0.74, 1.31, 1.29, 0.72). Leg formula: 1234.

PALP (Figs 3A–C, 4A–B). Yellowish brown palp with darker cymbium. Cymbium elongated and broad, with smoothly arched prolateral margin, while retrolateral side exhibits distinct depression (Figs 3A, C, 4A). Bulbus with prominent protuberance prolaterally (Figs 3A, 4A). Embolus filiform, broad at base, originating at 12 o'clock position (left palp in ventral view), encircling tegulum (Figs 3A, 4A). RTA moderately long, broad at base, and slightly curved distally (Figs 3A–C, 4A–B).

### Female (paratype, Figs 1E–F, 2D–F)

COLOURATION IN ALCOHOL. General aspects essentially as in male except following: habitus colour lighter than male, opisthosoma with creamy-white blotches and streaks (Fig. 2D).

MEASUREMENTS. Total length: 6.23. Carapace: length 2.66, width 2.21. Abdomen: length 3.57, width 2.24. Length of chelicerae: 1.08. Clypeus height: 0.18. Eye diameters: AME 0.55; ALE 0.31, PME 0.07, PLE 0.25; AME–AME 0.04. Eye interdistances: AME–ALE 0.11, ALE–ALE 1.38, ALE–PME 0.29, PME–PLE 0.27, PLE–PLE 1.92, PME–PME 1.94. Measurements of palp and legs: palp: 1.57 (0.61, 0.19, 0.24, 0.53), leg I 4.36 (1.38, 0.81, 0.97, 0.69, 0.51), II 4.03 (1.32, 0.69, 0.89, 0.68, 0.45), III 4.49 (1.60, 0.62, 0.97, 0.78, 0.52), IV 4.58 (1.48, 0.66, 0.89, 1.01, 0.54). Leg formula: 4312.

GENITALIA (Figs 3D–E, 4C–D). Epigyne broadly oval, with distinct central semi-crescent depression (Figs 3D, 4C). Copulatory openings slit-like, situated at rear edge of depression. Insemination ducts membranous, broad initially becoming narrow and coiled (Figs 3E, 4D); fertilization ducts prominent, directed laterally (Figs 3E, 4D).

## Distribution

India (Gujarat and Rajasthan).

## Variation

Copulatory duct coiling varies in all the female paratypes studied (Fig. 3E–H).

*Mogrus pune* Tripathi, Kulkarni & Kadam sp. nov.

[urn:lsid:zoobank.org:act:034C9812-C92D-4293-B40D-058B8048A78D](https://zoobank.org/act:034C9812-C92D-4293-B40D-058B8048A78D)

Figs 5–8

## Diagnosis

*Mogrus pune* sp. nov. both males and females are most similar to those of *M. mathisi* (Berland & Millot, 1941), males match by having a prominent protuberance between the 12 and 1 o'clock position; a

long, thin embolus tip end at the 1 o'clock position; RTA long, wide at base but can be distinguished by heart-shaped protuberance in ventral view (vs triangular in *M. mathisi*); bulbus of uneven shape (vs nearly circular in *M. mathisi*); RTA slightly bent at the tip in ventral view (vs tip nearly straight in *M. mathisi*) (compare Figs 7A–B, 8A–B with Wesolowska 2003: figs 1–6). Females are similar in having a large central depression on the anterior region of the epigyne, the copulatory duct anteriorly wide and membranous; the chevron pattern on the posterior half of the abdomen but can be distinguished by the shape of the depression on the epigyne and the coiling of the copulatory duct (compare Figs 7C–D, 8C–D with Wesolowska 2003: figs 7–10).

### Etymology

The specific epithet *pune* (noun in apposition) refers to Pune city, the type locality of the species. The name is dedicated to the small protected areas in and around Pune, which harbour a remarkable biodiversity despite urban pressures. It also highlights the importance of conserving the unique semi-arid ecosystems of the region.



**Fig. 5.** Field photographs of *Mogrus pune* Tripathi, Kulkarni & Kadam sp. nov. A–B. Holotype, ♂ (NRC-AA-0645). C–D. Paratype, ♀ (NRC-AA-0646).

## Type material

### Holotype

INDIA • ♂; Maharashtra, Pune, Alfa Laval Biodiversity Park, Pimpri Chinchwad; 18°42'11" N, 73°47'44" E; elev. 597 m; 31 May 2024; A. Kulkarni leg.; found on a tree branch, by hand; NCBS, NRC-AA-0645.


### Paratype

INDIA • 1 ♀; same data as for holotype; NCBS, NRC-AA-0646.

## Description

### Male (holotype, Figs 5A–B, 6A–C)

COLOURATION IN ALCOHOL. Carapace, eye region, clypeus, chelicerae, endites, labium, legs, opisthosoma, sternum dark brown (Figs 5A–B, 6A–C). Carapace with two prominent white bands of uniform thickness, originating behind PLEs and extending posteriorly, in between covered with black appressed hair (Figs 5B, 6A). Clypeus short; covered with long black hairs. Chelicerae elongated, with distinct gap visible in frontal view (Fig. 6C). Abdomen oval-shaped, tapers posteriorly, featuring dark brown median region with pale lateral bands (Figs 5B, 6A). Leg covered with white and black hairs intermixed; femur I–IV apical half pale colour (Fig. 6A).




**Fig. 6.** *Mogrus pune* Tripathi, Kulkarni & Kadam sp. nov. A–C. Holotype, ♂ (NRC-AA-0645). D–F. Paratype, ♀ (NRC-AA-0646). A, D. Habitus, dorsal view. B, E. Same, lateral view. C, F. Same, frontal view. Scale bars = 2 mm.

MEASUREMENTS. Total length. 5.78. Carapace length 2.94, width (at middle) 2.18. Abdomen length 2.84, width (at middle) 1.63. Eye diameters: AME 0.55, ALE 0.29, PME 0.07, PLE 0.27. Eye interdistances: AME–ALE 0.17, AME–PME 0.03, PME–PME 1.72, ALE–ALE 1.49, PME–PLE 0.29, PLE–PLE 1.94, ALE–PME 0.34. Length of chelicerae: 1.33. Measurements of palp and legs: palp: 2.36 (0.85, 0.22, 0.23, 1.06), leg I 5.6 (1.61, 0.92, 1.32, 1.06, 0.65), II 4.65 (1.51, 0.79, 1.03, 0.86, 0.46), III 4.8 (1.66, 0.70, 0.95, 0.88, 0.61), IV 4.75 (1.53, 0.66, 0.96, 1.03, 0.57). Leg formula: 1342.


PALP (Figs 7A–B, 8A–B). Primarily amber-colored, with some areas appearing darker (Fig. 7A). Cymbium elongated, broad, slightly tilted toward retrolateral side in ventral view (Figs 7A, 8A). Bulbus with large, heart-shaped protuberance positioned retrolaterally, appearing membranous and oval in retrolateral view (Figs 7A–B, 8A–B). Embolus filiform, broad at base, originating at 1 o'clock position (left palp in ventral view), encircling tegulum (Figs 7A, 8A). RTA long, gradually narrow and bend at tip in ventral view (Figs 7A–B, 8A–B).

**Female** (paratype, Figs 5C–D, 6D–F)

COLOURATION IN ALCOHOL. General aspects essentially as in male except following: habitus colour lighter than male; carapace with faint white band covered with black blotches (Fig. 6D). Opisthosoma lacks white stripes, with chevron pattern at posterior half (Fig. 6D).



**Fig. 7.** *Mogrus pune* Tripathi, Kulkarni & Kadam sp. nov., genitalia. **A–B.** Holotype, ♂ (NRC-AA-0645). **C–D.** Paratype, ♀ (NRC-AA-0646). **A–B.** Male left palp. **A.** Ventral view. **B.** Retrolateral view. **C–D.** Epigyne/vulva. **C.** Ventral view. **D.** Dorsal view. Scale bars = 0.2 mm.



**Fig. 8.** *Mogrus pune* Tripathi, Kulkarni & Kadam sp. nov., genitalia. **A–B.** Holotype, ♂ (NRC-AA-0645). **C–D.** Paratype, ♀ (NRC-AA-0646). **A–B.** Male left palp. **A.** Ventral view. **B.** Retrolateral view. **C–D.** Epigyne/vulva. **C.** Ventral view. **D.** Dorsal view. Scale bars = 0.2 mm.

MEASUREMENTS. Body length. 5.65. Carapace length 2.84, width (at middle) 2.28. Abdomen length 2.81, width (at middle) 1.76. Eye diameters: AME 0.55, ALE 0.29, PME 0.07, PLE 0.29. Eye interdistances: AME–ALE 0.21, AME–AME 0.02, PME–PME 1.99, ALE–ALE 1.58, PME–PLE 0.26, PLE–PLE 2.05, ALE–PME 0.31. Length of chelicerae: 1.07. Measurements of palp and legs: palp: 2.36 (0.85, 0.22, 0.23, 1.06), leg I 4.55 (1.43, 0.80, 1.01, 0.79, 0.52), II 4.14 (1.37, 0.72, 0.90, 0.68, 0.47), III 4.44 (1.52, 0.72, 0.79, 0.88, 0.53), IV 4.46 (1.49, 0.58, 0.85, 1.01, 0.53). Leg formula: 4132.

GENITALIA (Figs 7C–D, 8C–D). Epigyne subtriangular with slightly convex anterior margin and large median depression (Figs 7C, 8C). Copulatory opening large, widely spaced. Copulatory duct initially wide and membranous, while looped and darker at end (Figs 7D, 8D). Fertilization ducts prominent, diverging (Figs 7D, 8D).

### Distribution

India (Maharashtra).

### *New record for India*

*Mogrus larisae* Logunov, 1995  
Figs 9–10

*Mogrus larisae* Logunov, 1995: 596, figs 1–4, 7, 18–30 (♂♀).

*Mogrus larisae* – Wesolowska 1996: 34, fig. 21b–c (♂).

### Diagnosis

For diagnosis, see Logunov (1995) and Wesolowska (1996).

### Type material

#### Holotype (not examined)

KAZAKHSTAN • ♂; South Kazakhstan, Chimkent area, Arys; 3 May 1988; D. Logunov leg.; ISEN 2682.

#### Paratypes (not examined)

KAZAKHSTAN • 6 ♀♀; same data as for holotype; ISEN 2678. For complete list of paratypes, see Logunov (1995).

### Material examined

INDIA • 1 ♂, 4 ♀♀; Rajasthan, Jaisalmer, Desert National Park, Sudasri area; 26°43'03.9" N, 70°36'18.0" E; elev. 235 m; 24 Oct. 2022; R. Tripathi leg.; found on grass, by hand; NCBS, NRC-AA-0647 to NRC-AA-0651.

### Description

For description, see Logunov (1995) and Wesolowska (1996).


### Measurements

#### Male (Figs 9A–C, 10A–C)

Total length: 4.27. Carapace: length 2.08, width 1.49. Abdomen: length 2.19, width 1.26. Length of chelicerae: 0.76. Clypeus height: 0.06. AME 0.43, ALE 0.24, PME 0.06, PLE 0.22; AME–AME 0.03, AME–ALE 0.07, ALE–ALE 1.02, ALE–PME 0.23, PME–PLE 0.20, PLE–PLE 1.29, PME–PME 1.30. Measurements of palp and legs: palp: 1.61 (0.59, 0.15, 0.18, 0.69), leg I 3.65 (1.10, 0.57, 0.83, 0.66, 0.49), II 3.20 (1.02, 0.47, 0.70, 0.61, 0.40), III 3.79 (1.30, 0.54, 0.72, 0.75, 0.48), IV 3.73 (1.21, 0.48, 0.79, 0.79, 0.46). Leg formula: 3412.

**Female** (9D–F, 10D–K)

Total length: 7.81. Carapace: length 2.92, width 2.23. Abdomen: length 4.89, width 2.98. Length of chelicerae: 1.12. Clypeus height: 0.24. AME 0.55; ALE 0.28, PME 0.07, PLE 0.27; AME–AME 0.04, AME–ALE 0.13, ALE–ALE 1.42, ALE–PME 0.29, PME–PLE 0.30, PLE–PLE 1.87, PME–PME 1.83.



**Fig. 9.** *Mogrus larisae* Logunov, 1995. **A–C.** ♂ (NRC-AA-0647). **D–F.** ♀ (NRC-AA-0648) (D–F). **A, D.** Habitus, dorsal view. **B, E.** Same, lateral view. **C, F.** Same, frontal view. Scale bars: A–E = 2 mm; F = 1 mm.

Measurements of palp and legs: palp: 2.15 (0.82, 0.26, 0.33, 0.74), leg I 4.64 (1.45, 0.85, 1.02, 0.80, 0.52), II 4.33 (1.49, 0.76, 0.94, 0.64, 0.50), III 5.21 (1.81, 0.89, 0.89, 1.03, 0.59), IV 5.12 (1.71, 0.69, 1.05, 1.11, 0.56). Leg formula: 3412.

### Variations

Shape of the copulatory opening, spermathecae, and accessory gland varies in all the female paratypes studied (Fig. 10D–K).

### Records from India


Rajasthan.

### Distribution

Kazakhstan, Kyrgyzstan, India, Iran, Turkmenistan, Uzbekistan (Logunov 1995; Wesolowska 1996; WSC 2026; present record).

### Remark

First record of the species from India.



**Fig. 10.** *Mogrus larisae* Logunov, 1995. A–C. ♂ (NRC-AA-0647), left palp. A. Ventral view. B. Retrolateral view. C. Dorsal view. D–K. Epigyne/vulva. D–E. ♀ (NRC-AA-0648). F–G. ♀ (NRC-AA-0649). H–I. ♀ (NRC-AA-0650). J–K. ♀ (NRC-AA-0651). D, F, H, J. Ventral views. E, G, I, K. Dorsal views. Scale bars = 0.2 mm.

**First description of male**

*Mogrus rajasthanensis* Caleb, Chatterjee, Tyagi, Kundu & Kumar, 2017  
Figs 11–14

*Mogrus rajasthanensis* Caleb *et al.*, 2017: 332, figs 10–20 (♀).


**Diagnosis**

Males of *M. rajasthanensis* are similar to those of *M. frontosus* Simon, 1871 in having a protuberance at the 2 o'clock position, RTA long and broad at the base, curving dorsally at the tip, but can be distinguished by a slightly longer cymbium (vs short in *M. frontosus*); the palpal bulb longer than wide (vs nearly subequal in length and width in *M. frontosus*) (compare Figs 13A–B, 14A–B with Andreeva *et al.* 1981: figs 9–11). For diagnosis of females, see Caleb *et al.* (2017).

**Type material**

**Holotype** (not examined)

INDIA • 1 ♀; Rajasthan, Sirohi, Mount Abu; 24.5917° N, 72.7237° E; elev. 1167 m; 14 Mar. 2017; K. Tyagi and V. Kumar leg.; ZSI-CDT-AA560. Now deposited in NZC-ZSI 6581/18 (Dr John Caleb pers. com.).




**Fig. 11.** Field photographs of *Mogrus rajasthanensis* Caleb, Chatterjee, Tyagi, Kundu & Kumar, 2017, ♂ (NRC-AA-0652). **A.** Frontal view. **B–C.** Same, dorsal view. **D.** Same, lateral view.

**Material examined**

INDIA • 2 ♂♂, 2 ♀♀; Rajasthan, Jaisalmer, Desert National Park, Sudasri area; 26°43'23.0" N, 70°36'06.7" E; elev. 221 m; 24 Oct. 2022; R. Tripathi leg.; found on ground, by hand; NCBS, NRC-AA-0652 to NRC-AA-0655.

**Description****Male** (Fig. 12A–C)

COLOURATION IN ALCOHOL. Carapace, clypeus, endites, labium, legs, opisthosoma, sternum peach colour; chelicerae brown (Fig. 12A–C). Carapace with three distinct black longitudinal bands, a wide median black band extends from ocular region to posterior margin, flanked by two thinner black stripes along lateral margins, these lateral stripes originate near posterior lateral eyes and extend towards thoracic region, in between black bands white appressed hairs present appearing like white bands (Fig. 12A–C). All eyes positioned within lateral black bands; region around anterior eyes reddish orange (Fig. 12A–C). Clypeus region covered with long white hairs; thin brown stripes emerge from beneath ALEs and lateral sides of AMEs and extend laterally (Fig. 12C). Abdomen ovoid, with a distinct central dark band running longitudinally, surrounded by a dense covering of white hairs (Fig. 12A); ventral side uniformly pale, with scattered fine setae.



**Fig. 12.** *Mogrus rajasthanensis* Caleb, Chatterjee, Tyagi, Kundu & Kumar, 2017. A–C. ♂ (NRC-AA-0652). D–F. ♀ (NRC-AA-0653). A, D. Habitus, dorsal view. B, E. Same, lateral view. C, F. Same, frontal view. Scale bars = 1 mm.


MEASUREMENTS. Total length: 3.97. Carapace: length 1.98, width 1.48. Abdomen: length 1.99, width 1.12. Length of chelicerae: 0.71. Clypeus height: 0.09. AME 0.43; ALE 0.24, PME 0.04, PLE 0.19; AME–AME 0.03, AME–ALE 0.06, ALE–ALE 1.02, ALE–PME 0.24, PME–PLE 0.19, PLE–PLE 1.28, PME–PME 1.30. Measurements of palp and legs: palp: 1.48 (0.54, 0.15, 0.17, 0.62), leg I 3.42 (1.03, 0.50, 0.84, 0.64, 0.41), II 2.88 (0.94, 0.41, 0.66, 0.49, 0.38), III 3.74 (1.34, 0.46, 0.84, 0.74, 0.36), IV 3.55 (1.19, 0.45, 0.73, 0.73, 0.45). Leg formula: 3412.

PALP (Figs 13A–C, 14). Light to medium brown, with a darker sclerotized bulb. Bulb nearly oval, with prominent protuberance retrolaterally at 2 o'clock position in ventral view (Figs 13A, 14A). Embolus originate beneath protuberance; broad at base; encircling tegulum. RTA elongated, curved and tapering distally, hook-like (Figs 13B–C, 14B).

**Female** (Figs 12D–F, 13D–F)

For detailed descriptions and illustrations of females, see Caleb *et al.* (2017).


MEASUREMENTS. Total length: 5.20. Carapace: length 2.25, width 1.68. Abdomen: length 2.95, width 2.01. Length of chelicerae: 0.84. Clypeus height: 0.11. AME 0.43; ALE 0.24, PME 0.05, PLE 0.21; AME–AME 0.03, AME–ALE 0.07, ALE–ALE 1.06, ALE–PME 0.25, PME–PLE 0.19, PLE–PLE 1.46, PME–PME 1.42. Measurements of palp and legs: palp: 1.53 (0.61, 0.18, 0.23, 0.51), leg I 4.37 (1.05, 1.50, 0.77, 0.61, 0.44), II 3.05 (1.01, 0.49, 0.69, 0.46, 0.40), III 3.68 (1.31, 0.59, 0.70, 0.67, 0.41), IV 3.79 (1.22, 0.51, 0.78, 0.79, 0.49). Leg formula: 1432.



**Fig. 13.** *Mogrus rajasthanensis* Caleb, Chatterjee, Tyagi, Kundu & Kumar, 2017, genitalia. **A–C.** ♂ (NRC-AA-0652). **D–E.** ♀ (NRC-AA-0653). **F.** ♀ (NRC-AA-0654). **A–C.** Left palp. **A.** Ventral view. **B.** Retrolateral view. **C.** RTA close-up. **D–F.** Epigyne/vulva. **D.** Ventral view. **E–F.** Dorsal view. Scale bars = 0.2 mm.

**Variation**

The male does not exhibit noticeable differences, whereas the female shows several distinct variations: longitudinal grooves along the lateral margins differ in shape: in one specimen, the grooves appears slightly curved, whereas in another, it exhibits a sharp bend. Additionally, the copulatory ducts form a V-shaped bend, which is relatively short in one individual but elongated in another. Furthermore, the size of the spermathecae also varies among the specimens, indicating intraspecific morphological variation in the female genital structures (Fig. 13D–F).



**Fig. 14.** *Mogrus rajasthanensis* Caleb, Chatterjee, Tyagi, Kundu & Kumar, 2017, ♂ (NRC-AA-0652), genitalia, left palp. **A.** Ventral view. **B.** Retrolateral view. Scale bars = 0.2 mm.

## Records from India

Rajasthan (Caleb *et al.* 2017; present study).

## Distribution

India (WSC 2026).

## Remarks

*Mogrus rajasthanensis* was originally described based solely on female specimens (Caleb *et al.* 2017). In the present study, male specimens are provisionally attributed to this species. The females closely resemble those of *M. canescens* (C.L. Koch, 1846), whereas the males assigned here show a strong similarity to those of *M. frontosus* (Simon, 1871). Females identified as *M. canescens* from Morocco have a distinct posterior median epigynal pocket (Oger 2024), a feature absent in specimens from Yemen previously assigned to *M. canescens* (Wesołowska & van Harten 1994), as well as in the holotype and additional female material of *M. rajasthanensis* (Caleb *et al.* 2017). This suggests that at least part of the Yemeni material may be referable to *M. rajasthanensis*, although this interpretation requires confirmation.

During extensive field surveys in Rajasthan, no males corresponding to the known morphology of *M. canescens* were encountered. While this observation alone is not conclusive, it supports the possibility that *M. rajasthanensis* and *M. canescens* represent distinct taxa in the region. The close resemblance between the males, attributed here to *M. rajasthanensis*, and *M. frontosus*, combined with differences in the female genital morphology, highlights the difficulty of reliable sex matching within *Mogrus*. Given the poor taxonomic resolution of the genus and the likelihood of mismatched sexes across species with overlapping distributions, the association of males and females in *M. rajasthanensis* should be regarded as tentative. A DNA barcode is available for *M. rajasthanensis*; however, comparative molecular data for related species remain insufficient. A comprehensive revision of *Mogrus*, integrating morphology, molecular data, and material from type localities, will be required to resolve these taxonomic uncertainties.

## Discussion

Identification of species of *Mogrus* can be challenging both in the field and under a microscope due to limited morphological differences and significant intraspecific variation. Most species have two longitudinal white stripes on the carapace and a central stripe or chevron pattern on the abdomen. Subtle differences in copulatory structures, such as those between *M. rajasthanensis* and *M. frontosus*, or *M. larisae* and *M. antoninus*, further complicate identification. Significant variation in female genitalia, discussed in detail in the variation sections, highlights the importance of carefully evaluating these features when designating species within the genus.

The discovery of *M. shushka* sp. nov. and *M. pune* sp. nov., together with the first record of *M. larisae*, significantly extends the known range of *Mogrus* into the western arid zones of India. This distribution aligns with the genus's core range in the southern Palaearctic and arid Afro-Asia (WSC 2026), supporting historical biogeographic connections via the Indian subcontinent's arid corridor (Seton *et al.* 2012). The presence of multiple species of *Mogrus* in semi-arid habitats of western India highlights the underexplored salticid diversity of these ecosystems and underscores their biogeographic significance.

## Acknowledgments

The collection of *Mogrus shushka* sp. nov. was carried out as part of the baseline survey under the restoration program of the Wildlife Conservation Trust (WCT). We extend our heartfelt appreciation to

WCT for facilitating this study and providing essential logistical support. We also gratefully acknowledge Karavansrai Investment Private Limited for funding the program through WCT, which made this work possible. The collection of Desert National Park specimens was conducted under the Bustard Recovery Programme of the Wildlife Institute of India (WII), with support from the National Compensatory Afforestation Fund Management and Planning Authority, Government of India, and supplemented by an additional grant from the Rajasthan State Pollution Control Board. RT and AKJ thank to Bustard recovery program at Wildlife Institute of India for field logistic and support in Desert National Park. RT and AKJ also thank Rajasthan State Forest Department for providing the necessary permits for the surveys in DNP. RT thanks Dr John Caleb (India) for his consistent support and valuable suggestions throughout this study. AK expresses sincere gratitude to the Alfa Laval PCMC Biodiversity Park, Pimpri Chinchwad, for its invaluable support in the biodiversity survey and its dedicated efforts toward conserving the park's rich biodiversity and thanks are extended to Monali Shah for her unwavering guidance and support throughout the study. RT expresses his gratitude to Rev. Fr. Jolly Andrews, CMI, of Christ College (Irinjalakuda) for providing research facilities. RT also acknowledges the support of Dr Sutirtha Dutta from WII, Dehradun, for his continuous support. Special thanks are due to Sohan Lal Genwa and Amrat Genwa for their invaluable assistance during field activities at Thar desert. Finally, RT acknowledges the financial support received from CSIR-UGC in the form of a fellowship and extends special thanks to Dr Pritha Dey and Mr Tarun Karmakar for providing the accession number for depositing the specimen at the NCBS Research Collection, Bangalore.

## References

- Andreeva E.M., Kononenko A.P. & Prószyński J. 1981. Remarks on genus *Mogrus* Simon, 1882 (Aranei, Salticidae). *Annales Zoologici, Warszawa* 36: 85–104.
- Azarkina G.N. & Foord S.H. 2013. Redescriptions of poorly known species of jumping spiders (Araneae: Salticidae) from South Africa and Namibia. *Zootaxa* 3686 (2): 165–182. <https://doi.org/10.11646/zootaxa.3686.2.3>
- Caleb J.T.D., Chatterjee S., Tyagi K., Kundu S. & Kumar V. 2017. Two new jumping spiders of the genera *Epocilla* Thorell, 1887 and *Mogrus* Simon, 1882 from India (Araneae: Salticidae). *Arthropoda Selecta* 26 (4): 329–334. <https://doi.org/10.15298/arthscl.26.4.08>
- Kanesharatnam N. & Benjamin S.P. 2018. A new genus and three new species of jumping spiders (Araneae: Salticidae) from Sri Lanka. *European Journal of Taxonomy* 444: 1–24. <https://doi.org/10.5852/ejt.2018.444>
- Logunov D.V. 1995. The genus *Mogrus* (Araneae: Salticidae) of Central Asia. *European Journal of Entomology* 92 (3): 589–604.
- Maddison W.P. 2015. A phylogenetic classification of jumping spiders (Araneae: Salticidae). *Journal of Arachnology* 43 (3): 231–292. <https://doi.org/10.1636/arac-43-03-231-292>
- Oger P. 2024. *Les Araignées de Belgique et de France*. Available from <https://arachno.piwigo.com> [accessed 28 Jan. 2026].
- Petrunkévitch A. 1928. Systema Aranearum. *Transactions of the Connecticut Academy of Arts and Sciences* 29: 1–270.
- Prószyński, J. 1976. Studium systematyczno-zoogeograficzne nad rodziną Salticidae (Aranei) Regionów Palearktycznego i Nearktycznego. *Wyższa Szkoła Pedagogiczna Siedlcach* 6: 1–260.
- Prószyński J. 2016. Monograph of Salticidae (Araneae) of the World 1995–2016. Part II: Global Species Database of Salticidae (Araneae). Available from <https://salticidae.pl/salticidae.php?adres=publication.php?id=3500> [accessed 28 Jan. 2026].

Seton M., Müller R.D., Zahirovic S., Gaina C., Torsvik T., Shephard G., Talsma A., Gurnis M., Turner M., Maus S. & Chandler M. 2012. Global continental and ocean basin reconstructions since 200 Ma. *Earth-Science Reviews* 113: 212–270. <https://doi.org/10.1016/j.earscirev.2012.03.002>

Tripathi R., Kadam G., Asha T.J. & Sudhikumar A.V. 2023. First record of *Sparbambus* Zhang, Woon & Li, 2006 from India, with description of a new species (Araneae: Salticidae). *Zootaxa* 5352 (3): 447–450. <https://doi.org/10.11646/zootaxa.5352.3.10>

Wesołowska W. 1996. New data on the jumping spiders of Turkmenistan (Aranei, Salticidae). *Arthropoda Selecta* 5 (1/2): 17–53.

Wesołowska W. 2003. A redescription of *Mogrus mathisi* (Berland et Millot, 1941) and its synonyms (Araneae: Salticidae). *Genus* 14: 425–430.

Wesołowska W. & van Harten A. 1994. *The jumping spiders (Salticidae, Araneae) of Yemen*. Yemeni-German Plant Protection Project, Sana'a.

Wesołowska W. & van Harten A. 2010. Order Araneae, family Salticidae. *Arthropod Fauna of the UAE* 3: 27–69.

World Spider Catalog. 2026. World Spider Catalog. Version 26. Natural History Museum Bern. Available from <http://wsc.nmbe.ch> [accessed 28 Jan. 2026]. <https://doi.org/10.24436/2>

Printed versions of all papers are deposited in the libraries of two of the institutes that are members of the *EJT* consortium: Muséum national d'Histoire naturelle, Paris, France and Royal Museum for Central Africa, Tervuren, Belgium. The other members of the consortium are: Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Meise Botanic Garden, Meise, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic; The Steinhardt Museum of Natural History, Tel Aviv, Israël.