

Received: 19 November 2025 • Accepted: 16 February 2026 • Published: 21 April 2026

Topic editor: Tony Robillard • Section editor: Gavin Broad • Desk editor: Pepe Fernández

### Research article

[urn:lsid:zoobank.org:pub:A6AC8AD2-DC97-4CBC-B5D5-0E1FC44045CA](https://zoobank.org/pub:A6AC8AD2-DC97-4CBC-B5D5-0E1FC44045CA)

## Review of the genus *Acantholabus* Heinrich (Hymenoptera: Ichneumonidae)

Xia YANG<sup>1</sup>  , Alexey RESHCHIKOV<sup>2,\*</sup>  , Namiki KIKUCHI<sup>3</sup>  ,  
Matthias RIEDEL<sup>4</sup>   & Zhi-Pang HUANG<sup>5</sup>  

<sup>1,5</sup>Institute of Eastern-Himalaya Biodiversity Research, Dali University, Yunnan 671003, P.R. China.

<sup>2</sup>Insect Biodiversity and Biogeography Lab, School of Biological Sciences, University of Hong Kong, Kadoorie Biological Sciences Building, Pokfulam Road, Hong Kong SAR, P.R. China.

<sup>3</sup>Toyohashi Museum of Natural History, 1-238 Oana, Oiwa, Toyohashi, Aichi 441-3147, Japan.

<sup>3</sup>Systematic Zoology Laboratory, Department of Biological Sciences, Graduate School of Science, Tokyo Metropolitan University, 1-1 Minamiosawa, Hachioji-shi, Tokyo, 192-0397, Japan.

<sup>4</sup>Blumenlage 22 C, D-29683 Bad Fallingb., Germany.

<sup>5</sup>Center for Interdisciplinary Sciences, Dali University, Dali, Yunnan 671003, China.

<sup>5</sup>Collaborative Innovation Center for Biodiversity and Conservation in the Three Parallel Rivers Region of China, Dali, Yunnan 671003, China.

\*Corresponding author: [alexeyre@hku.hk](mailto:alexeyre@hku.hk)

<sup>1</sup>Email: [2723154053@qq.com](mailto:2723154053@qq.com)

<sup>3</sup>Email: [namikikikuchi@gmail.com](mailto:namikikikuchi@gmail.com)

<sup>4</sup>Email: [mamaflo.riedel@t-online.de](mailto:mamaflo.riedel@t-online.de)

<sup>5</sup>Email: [huangzp@eastern-himalaya.cn](mailto:huangzp@eastern-himalaya.cn)

**Abstract.** The genus *Acantholabus* Heinrich, 1974 is reviewed. The three previously known species are re-described, and a new species, *Acantholabus yunlingensis* Reshchikov & Kikuchi sp. nov., from Northwestern Yunnan, is described and illustrated. An identification key to the species of the genus is also provided.


**Keywords.** Darwin wasps, Eastern Himalaya, Three Parallel Rivers, Ichneumoninae, Platylabini.

Yang X., Reshchikov A., Kikuchi N., Riedel M. & Huang Z.-P. 2026. Review of the genus *Acantholabus* Heinrich (Hymenoptera: Ichneumonidae). *European Journal of Taxonomy* 1052: 1–20.

<https://doi.org/10.5852/ejt.2026.1052.3263>

### Introduction

*Acantholabus* Heinrich, 1974 is a small endemic Himalayan genus in the ichneumonine tribe Platylabini Berthoumieu, 1904 (Hymenoptera, Ichneumonidae, Ichneumoninae), comprising three previously known species described from Kachin State of Myanmar (Heinrich 1974) (Fig. 1A). Prior to this study, there



**Fig. 1.** A. *Acantholabus* distribution map, [yellow] = *A. yunlingensis* Reshchikov & Kikuchi sp. nov., [red] = *A. furciliatus* Heinrich, 1974, *A. malaisei* Heinrich, 1974, *A. regressus* Heinrich, 1974, available from ©Google Maps: <https://www.google.com/maps/d/u/2/edit?mid=1L35S1cDTajUBSnR64heXj83fcetdPck&usp=sharing>. B. Aerial photo of the type locality of *A. yunlingensis*, Heilong spring valley in the Cang Mountain foothills (2300 m a.s.l.) near Dali University. C. Aerial photo of the sampling site in the Chongjiang River valley at Mt Laojun. D. Malaise trap at the type locality of *A. yunlingensis*, Heilong spring in the Cang Mountain foothills (2300 m a.s.l.) near Dali University. E. Aerial photo of the sampling site in the Chongjiang River valley at Mt Laojun with Malaise trap.

were 76 Oriental species of Platylabini (Yu *et al.* 2016; Riedel 2017, 2023a), most of which are known from the Himalayan region (Heinrich 1974; Riedel 2017).

Within the Platylabini, *Acantholabus* resembles *Dentilabus* Heinrich, 1974, these genera sharing characteristics such as large and broad thyridia (each wider than the space between them), as well as elongate propodeal spiracles (Tereshkin 2009). Both *Acantholabus* and *Dentilabus* possess long propodeal apophyses, but the former can be distinguished from the latter by weak longitudinal impression between thyridia (distinct in *Dentilabus*), and flat horizontal part of scutellum, which is usually convex in *Dentilabus* (for details see remarks for the genus diagnosis).

Unfortunately, nothing is known about the biology of the genus. However, all species of *Acantholabus* (including the species described here) are restricted to the Eastern Himalaya and are found between 2100–2400 m above sea level (Fig. 1A). Other Platylabini are known as larval parasitoids of Drepanidae Meyrick, 1895 and Geometridae Leach, 1815 (Bradley 1978; van Veen 1981; Short *et al.* 2002; Riedel 2008; Riedel *et al.* 2013; Shaw *et al.* 2015).

The present publication describes a new species of *Acantholabus*, *Acantholabus yunlingensis* Reshchikov & Kikuchi sp. nov. from the Yun Mountains Range, which forms a natural boundary between the Mekong and upper flow of the Yangtze Rivers, and is the part of the Three Parallel Rivers Region of Yunnan.

## Material and methods

This work is based on material collected with Malaise traps during the “Darwin Wasps of the Eastern Himalaya” project. We operated 63 Malaise traps in northwestern Yunnan on a permanent basis from 2018 till 2023. We examined samples from 52 trap-years; despite the highly representative material, we found only three females of the new species of *Acantholabus*. These specimens were found in the samples from a site with deciduous broad-leaf forest (dominant species *Alnus nepalensis* D. Don and *Pinus yunnanensis* Franch.) at Heilong spring, in the foothills of Cang Mountain (2300 m a.s.l.) near Dali University (Fig. 1B, D), and a site at a farmland with *P. yunnanensis* along the Chongjiang River valley, on the foothills of Laojun Mountain (Yulong Naxi Autonomous County, Lijiang) (Fig. 1C, E).

Specimens were examined using a Motic SMZ171 stereo microscope. Images were acquired digitally using the CA005 microscopic imaging system and processed with Adobe Photoshop. Distribution maps were produced using Google Earth Pro.

Morphological terminology and wing venation nomenclature follow Broad *et al.* (2018) and aligned with the Hymenoptera Anatomy Ontology (HAO) (Yoder *et al.* 2010). Abbreviations and morphological terms used in the text are as follows:

- OD = the longest diameter of a posterior lateral ocellus
- OOL = the shortest distance between a posterior lateral ocellus and a compound eye
- POL = the shortest distance between the posterior lateral ocelli
- S1–S6 = refer to the metasomal sternites
- T1–T7 = refer to the metasomal tergites 1–7

The specimens examined are deposited in the following institutions (curators in parentheses):

- IEHBR = Institute of Eastern-Himalaya Biodiversity Research, Dali (Xian-Fu Li)
- ISAS = Kunming Institute of Zoology, Kunming (Kaiqing Li)
- NHRS = Swedish Museum of Natural History, Stockholm (Hege Vårdal)
- ZSM = The Bavarian State Collection of Zoology, Munich, Germany (NA)

## Results

### Taxonomy

Class Insecta Linnaeus, 1758  
Order Hymenoptera Linnaeus, 1758  
Superfamily Ichneumonoidea Latreille, 1802  
Family Ichneumonidae Latreille, 1802  
Subfamily Ichneumoninae Latreille, 1802  
Tribe Platylabini Berthoumieu, 1904  
  
Genus *Acantholabus* Heinrich, 1974

*Acantholabus* Heinrich, 1974: 157–158. Type species: *Acantholabus malaisei* Heinrich, 1974 by original designation.

### Diagnosis

Female antenna slender, bristle-shaped, with elongate basal flagellar segments, not widened beyond middle, with a white ring. Male antenna with distinct, elongate tyloids. Eye rather large. Gena short. Vertex abruptly sloping just behind ocelli. Temples short, sharply and roundly narrowed behind the eyes. Occipital carina with dorsal impression which does not reach level of eyes or hind ocelli. Clypeus strongly convex, weakly separated from middle field of face. Mandibles gradually narrowed, dorsally and ventrally carinated. Pronotum without carina interrupting transverse furrow. Upper margin of pronotum not thickened. Pronotal base gradually and uniformly curved. Mesoscutum rather strongly convex, almost as long as broad, very densely punctured. Notauli visible only at anterior third of mesoscutum. Mesoscutum matt. Subtegular ridge thick and sharp. Sternaulus weakly impressed at front half. Scutellum uniformly raised, laterally and apically carinate with high, sharp carinae, its horizontal part flat. Hind margin of metanotum with teeth. Horizontal part of propodeum in middle distinctly shorter than area petiolaris. Lateral propodeal carina indistinct, if present, situated at middle of area superomedia. Area dentipara short, with longer or short but distinct denticular apophysis. Juxtacoxal carina distinct. Propodeal spiracle relatively large, elongate, oval. Legs rather long and slender. Hind coxa elongate. Fore wing with areolet rather large, quadrangular, and symmetrical. Vein *R* straight and long. Vein *Icu-a* interstitial. T1 petiole thick, strongly flattened at base. Gastrocoeli weakly impressed and elongate. Thyridia large and broad, distant from base of T2, with rather narrow interval between them with weak impression. Hypopygium relatively large, roughly triangular in outline, without distinct longitudinal fold. Ovipositor sheath slightly projecting behind apex of metasoma.

### Remarks

This genus resembles *Dentilabus*, but can be distinguished by the following characters: male flagellum with distinct tyloids (usually absent in *Dentilabus*); horizontal part of scutellum flat (usually convex in *Dentilabus*); T2 with weak median longitudinal impression between thyridia. *Dentilabus iyoensis* Kikuchi & Konishi, 2018 is an exception, as it shares these characteristics with *Acantholabus* (Kikuchi & Konishi 2018). The generic position of this species should be reconsidered in future studies.

*Acantholabus furciliatus* Heinrich, 1974  
Figs 2–3

*Acantholabus furciliatus* Heinrich, 1974: 160.

### Diagnosis

Can be easily distinguished from all the other species of *Acantholabus* by this combination of characters: (1) propodeal apophysis distinct, as long as basally wide in profile (Fig. 3D); (2) scutellum with apically


tooth-like lateral protrusions (Fig. 3D); (3) T1 with dorso-lateral carina (Fig. 3D); (4) body mostly dark coloured (Fig. 2); (5) T1–2 with broad ivory apical margins (Figs 2, 3C–D).

### Type material


#### Holotype

MYANMAR • ♀; Kachin, Kan Paik Ti; 2100 m a.s.l.; 9 Jun. 1934; R. Malaise leg.; ZSM, Hym. 836.

“[White label] N.E. BURMA / Kambaiti 2000 m / 9.6.1934 Malaise // [White label] *Acantholabus* / *furciliatus* / det. Heinrich Hein // [Red label] Typus Nr. Hym. / 836 / Zoologische / Staatssammlung / München”.



**Fig. 2.** *Acantholabus furciliatus* Heinrich, 1974, holotype, ♀ (ZSM Hym. 836). A. Habitus, in lateral view. B. Habitus, in dorsal view.



**Fig. 3.** *Acantholabus furciliatus* Heinrich, 1974, holotype, ♀ (ZSM Hym. 836). **A.** Face. **B.** Head in dorsal view. **C.** Metasoma in dorsal view. **D.** Mesosoma in dorso-lateral view.

## Re-description

### Female

**HEAD.** Body length 8 mm. Left flagellum with 8 basal flagellomeres, right flagellum missing. First flagellomere  $7.8\times$  as long as broad at apex, all flagellomeres longer than wide (Fig. 2). Head rather transverse,  $3.65\times$  as long as broad, strongly narrowed behind eyes, (Fig. 3B). POL:OD:OOL = 1.2:1:0.9. Vertex granulate. Eye large. Frons granulate, dull. Face 1.8 times as broad as high (Fig. 3A), granulate, dull. Clypeus moderately convex,  $2.1\times$  as broad as high, granulate, its apical margin sharp, weakly rounded. Mandible moderately narrowed from base to apex, with dorsal and ventral carinae, its ventral tooth slightly shorter than dorsal tooth. Malar space  $0.8\times$  as long as basal mandibular width.

**MESOSOMA.** Notaulus weakly impressed at frontal 0.2 of mesoscutum. Mesoscutum finely punctulate, slightly longer than wide, dull. Mesopleuron including speculum and metapleuron finely granulate, dull. Sternaulus slightly impressed at frontal half of mesopleuron. Scutellum pyramidal, punctulate, lateral carina complete, with dentiform spine posteriorly (Fig. 3D). Propodeum finely rugose-punctate (Fig. 3D). Area basalis trapezoid, much wider than long. Area superomedia heart-shaped, slightly wider than long, with some longitudinal rugae; lateral propodeal carina indistinct. Posterolateral edge of area dentipara with long apophysis ( $2\times$  as long as basally wide) (Fig. 3D). Juxtacoxal carina present. Hind coxa  $1.9\times$  as long as wide, granulate, without scopa. Hind femur  $4.7\times$  as long as wide. Fore wing with areolet quadrangular, pointed anteriorly, *2m-cu* reaching its middle. Vein *1cu-a* interstitial to *M&RS*.

**METASOMA.** Amblypygous (Fig. 2). T1 with lateral carina; petiole wider than high,  $1.9\times$  as long as postpetiole. Postpetiole without median field, granulate. S2–4 with median folds. Gastrocoelus triangularly impressed, granulate (Fig. 3D). Thyridium oblique, ca  $1.3\times$  as wide as interval between thyridia (Fig. 3D). Ovipositor sheath reaching slightly beyond metasomal apex.

**COLOUR.** Black (Fig. 2). Flagellum black, apical half of 7<sup>th</sup> flagellar segment and 8<sup>th</sup> segment ivory dorsally. Head black; palps, mandible except teeth, labrum, clypeus and face except central longitudinal black stripe, frontal orbit, stripe on gena and triangular spot of vertex ivory. Mesosoma black; frontal and upper margins of pronotum, subtegular ridge, central spot of scutellum, apophysis and area posteroexterna of propodeum yellow. Tergites blackish; apical half of postpetiole and apical band of T2 yellow. Apical bands of T6 and T7 mainly ivory. Fore and mid coxae ivory (mid coxa with some reddish suffusion); legs otherwise reddish; fore and mid tarsi black; hind femur infusate in apical 0.2; hind tibia blackish, with subbasal reddish-brown ring; hind metatarsus and 2<sup>nd</sup> hind tarsomere ivory (following tarsomeres missing). Wings almost hyaline, pterostigma brownish.

### Male

Unknown.

### Host

Unknown.

### Distribution

Myanmar.

*Acantholabus malaisei* Heinrich, 1974

Figs 4–5

*Acantholabus malaisei* Heinrich, 1974: 158–160.

### Diagnosis

Can be easily distinguished from all the other species of *Acantholabus* by this combination of characters: (1) propodeal apophysis elongate,  $2.8\times$  as long as broad in profile, and downcurved (Figs 4, 5D, 7E);

(2) scutellum apically without tooth-like lateral protrusions (Fig. 5D); (3) T1 with lateral carina (Fig. 5G); (4) body mostly dark coloured (Figs 4–5); (5) T1–2 without broad white apical margins (Fig. 5G).

### Type material

#### Holotype

MYANMAR • ♀; Kachin, Kan Paik Ti; 2100 m a.s.l.; 20 May 1934; R. Malaise leg.; NHRS, NHRS-HEVA000018484.

“[White label] N.E. BURMA / Kambaiti 7000 ft / 20.V.1934 / R. MALAISE // [White label] *Acantholabus / malaisei* / ♀ Hir. / det. G. Heinrich // [Red label] Holotype // [Red label] 442 // [White label] NHRS-HEVA / 000018484”.

### Re-description

#### Female

**HEAD.** Antenna with 38 flagellomeres, slender, slightly broadened beyond middle, not flattened ventrally, with white ring dorsally (Fig. 4). Basal flagellomeres elongate, first flagellomere  $6.8\times$  as long as broad at apex. Eye rather large (Fig. 5A–B). Face  $1.6\times$  as broad as high (Fig. 5A), without punctures, shagreened, weakly protruding in profile (Fig. 5B). Space between scapes with impression (Fig. 5A). Clypeus separate from face with slight impression, rounded, slightly convex in profile (Fig. 5B),  $1.9\times$  as broad as high (Fig. 5A), sculptured as face. Its apical margin sharp. Labrum visible. Mandible strongly narrowed from base to its apex, sharply carinated dorsally and ventrally (Fig. 5A–B). Malar space  $0.9\times$  as long as basal mandibular width. Gena short, shagreened (Fig. 5B). Head rather transverse, strongly narrowed behind eyes,  $0.3\times$  as long as broad (Fig. 5C). Vertex sloping down immediately behind ocelli (Fig. 5C), shagreened. POL:OD:OOL = 1.2:1:0.9. Occipital carina not impressed medially, lower end erased, not reaching base of mandible or hypostomal carina (Fig. 5B).


**MESOSOMA.** Pronotal collar short. Mesoscutum rather strongly convex, slightly longer than broad, densely punctate with shagreened microsculpture, matt. Notauli weakly impressed in anterior third. Subtegmental ridge defined. Speculum with wrinkled sculpture. Mesopleural fovea consisting of a short, deep horizontal groove. Sternaulus rather vestigial, visible at anterior part. Mesopleurae densely punctured, wrinkled, matt. Scutellum high, elevated above postscutellum, carinated laterally and apically by high sharp carina (Fig. 5D). Hind margin of metanotum with triangular projections opposite lateral longitudinal carinae. Propodeum tetrahedral in dorsal profile, horizontal part equal to area petiolaris (Fig. 5F). Propodeal carinae developed. Area basalis rather long. Area superomedia hexagonal, approximately as long as broad. Lateral propodeal carina present, joining area superomedia before its middle. Apophysis long (Fig. 5F). Juxtacoxal carina distinct. Propodeal spiracles elongate oval. Legs long and robust. Hind coxa  $1.6\times$  times as long as broad. Hind femur  $4.2\times$  as long as wide. Fore wing with areolet large, quadrangular, receiving vein 2m-cu at its middle. Pterostigma broad. Vein 1cu-a interstitial to M&RS. Hind wing with distal abscissa of Cu straight, Cu about  $1.7\times$  as long as cu-a, these veins almost opposite (Fig. 4).

**METASOMA.** T1 with distinct lateral carina, petiole short,  $1.6\times$  as long as postpetiole (in profile) (Figs 4, 5G–H). Gastrocoeli weakly impressed and elongate. Thyridia large and broad, distant from base of T2, with rather narrow interval between them with weak impression (Fig. 5G). Ovipositor sheath slightly projecting behind apex of metasoma (Fig. 4).

**COLOUR.** Body mostly blackish-brown (Fig. 4), except: palpi, face and clypeus laterally (Fig. 5A), spots at POL (Fig. 5C), collar, pronotal base and subtegmental ridge (Fig. 5B), scutellum posteriorly (Fig. 5D), and propodeal apophysis (Fig. 5F) yellow; legs dorsally dark-red (Fig. 4); legs ventrally (Fig. 4), T1 postpetiole, and hind margin of T2 (Fig. 5G) orange-brown; flagellomeres 8–10, and T6–7 ivory (Fig. 4).



**Fig. 4.** *Acantholabus malaisei* Heinrich, 1974, holotype, ♀ (NHRS, NHRS-HEVA000018484). Habitus in lateral view.



**Fig. 5.** *Acantholabus malaisei* Heinrich, 1974, holotype, ♀ (NHRS, NHRS-HEVA000018484). **A.** Face. **B.** Head in lateral view. **C.** Head in dorsal view. **D.** Mesoscutum in dorsal view. **E.** Propodeum and T1–2 in dorso-lateral view. **F.** Propodeum in lateral view. **G.** Metasoma in dorso-lateral view. **H.** Metasoma in lateral view.

### Male

Antenna with 38 flagellomeres with distinct, elongate tyloids (on 8–20 flagellomeres), with white ring on flagellomeres 11–15. Face and clypeus mostly white, except only one black longitudinal band in their middle. Middle and hind coxae without black markings, fore and mid coxae white spotted. T5–6 with membranous, white hind margins, T7 with white apical marking. Hind tarsus yellowish. The rest as in female.

### Host

Unknown.

### Distribution

Myanmar.

### *Acantholabus regressus* Heinrich, 1974

Figs 6–7

*Acantholabus regressus* Heinrich, 1974: 161.

### Diagnosis

This species can be easily distinguished from all the other *Acantholabus* species by this combination of characters: (1) propodeum with short tooth-like protrusions not forming apophysis (Fig. 7E); (2) scutellum without tooth-like lateral protrusions (Fig. 7D); (3) T1 with lateral carina; (4) body mostly dark (Fig. 6); (5) T1–2 without broad white apical margins (Fig. 7G).

### Type material

#### Holotype

MYANMAR • ♀; Kachin, Kan Paik Ti; 2100 m a.s.l.; 8 Jun. 1934; R. Malaise leg.; NHRS, NHRS-HEVA000018485.

“[White label] N.E. BURMA / Kambaiti 7000 ft / 8/6 R. MALAISE // [White label] *Acantholabus / regressus* / 1970 Hein. / det. G. Heinrich // [Red label] Holotype // [Red label] 331 // [White label] NHRS-HEVA / 000018485”.

### Re-description

#### Female

HEAD. Antenna with 31 flagellomeres, slender, weakly broadened beyond middle, not flattened ventrally, with white ring dorsally (Fig. 6). Basal flagellomeres elongate, first flagellomere  $4.9\times$  as long as broad at apex. Eye rather large (Fig. 7A–B). Face  $1.3\times$  as broad as high (Fig. 7A), flat, with sparse punctures, setose, shining, not protruding in profile (Fig. 7B). Space between scapes with impression (Fig. 7A). Clypeus separated from face by distinct impression, inverted trapezoid shape (Fig. 7A), slightly convex in profile (Fig. 7B),  $1.6\times$  as broad as high (Fig. 7A), sculptured as face. Its apical margin sharp. Labrum visible. Mandible strongly narrowed from base to its apex. Malar space elongate,  $1.5\times$  as long as basal mandibular width (Fig. 7A). Gena elongate (Fig. 7B). Head subquadrate, roundly narrowed behind eyes,  $0.4\times$  as long as broad (Fig. 7C). Vertex sloping down behind ocelli (Fig. 7B), shagreened (Fig. 7C). POL:OD:OOL = 1.5:1:1.3.

MESOSOMA. Pronotal collar short (Fig. 7D). Mesoscutum rather strongly convex (Fig. 6), slightly longer than broad (Fig. 7D), densely punctate on shagreened microsculpture, matt. Notauli absent (Fig. 7D). Subtegular ridge defined. Speculum coarsely punctate, strongly reticulate. Mesopleural fovea a short, shallow horizontal groove covered with setae. Sternaulus absent. Mesopleurae punctured with deep,

scattered punctures, shagreened, matt. Scutellum high, elevated above postscutellum, with high, sharp carinae laterally and apically, carina, at apex interrupted, forming lateral teeth-like structure. Propodeal carinae developed (Fig. 7E). Area basalis subquadrate. Area superomedia pentagonal, approximately as long as broad (Fig. 7E). Apophysis distinct. Propodeal spiracles elongate oval. Legs long and slender. Hind coxa elongate,  $1.5\times$  as long as broad. Hind femur  $4.6\times$  as long as wide. Ratio of length of hind tarsomeres  $3:1.5:1:0.6:0.9$ . Claws elongate, slightly curved. Fore wing with areolet large, quadrangular, receiving vein  $2m-cu$  at its middle (Fig. 7H). Vein  $1cu-a$  interstitial to  $M\&RS$ . Hind wing with distal abscissa of  $Cu$  straight,  $Cu$  about 3.5 times as long as  $cu-a$ , these veins almost opposite (Fig. 7H).

**METASOMA.** T1 with lateral carina, petiole short,  $1.6\times$  as long as postpetiole (in profile) (Fig. 6). Gastrocoeli weakly impressed and elongate. Thyridia large and broad, distant from base of T2, with rather narrow interval between them with weak impression (Fig. 7G).

**COLOUR.** Body mostly black (Fig. 6), except: scapes ventrally, face and clypeus laterally, gena behind eye margin, upper part of propleuron, tegula, dorso-anterior corner of mesopleuron, fore and middle coxae and trochanters, mesoscutellum dorsally, hind margins of T4–7 yellow; flagellomeres 11–16 white; fore and middle legs, hind tarsus reddish-yellow; hind leg (except coxa) reddish; pterostigma and mesosoma brown (Figs 6–7).

#### Male

Unknown.

#### Host


Unknown.

#### Distribution

Myanmar.



**Fig. 6.** *Acantholabus regressus* Heinrich, 1974, holotype, ♀ (NHRS, NHRS-HEVA000018485). Habitus in lateral view.



**Fig. 7.** *Acantholabus regressus* Heinrich, 1974, holotype, ♀ (NHRS, NHRS-HEVA000018485). **A.** Face, **B.** Head in lateral view. **C.** Head in dorsal view. **D.** Mesoscutum in dorsal view. **E.** Propodeum in dorso-lateral view. **F.** Mesopleuron in lateral view. **G.** Metasoma in dorsal view. **H.** Wings.

*Acantholabus yunlingensis* Reshchikov & Kikuchi sp. nov.  
[urn:lsid:zoobank.org:act:F380526E-27DB-4005-88FA-5841E69E4B0F](https://doi.org/10.3896/eb.urn:lsid:zoobank.org:act:F380526E-27DB-4005-88FA-5841E69E4B0F)

Figs 8–9

### Diagnosis

Can be easily distinguished from all the other *Acantholabus* species by the following combination of characters: (1) propodeal apophysis small and straight (Fig. 9E); (2) scutellum without tooth-like lateral protrusions (Fig. 9H); (3) T1 without dorso-lateral carina; (4) body mostly brown-yellow with dark charcoal mesoscutum (Fig. 8); (5) T1–2 without white colouration (Figs 8, 9C, I).

### Etymology

The species epithet is an adjective, and refers to the Yun Mountains (Chinese 雲嶺 [Yun Ling], literally “Cloudy Peaks”), where this new species was discovered.

### Type material

#### Holotype

CHINA • ♀; Yunnan, Dali, Mt Cang, Heilong spring; 25°40′02.0″ N, 100°08′51.4″ E; 2300 m a.s.l.; 29 Sep.–6 Oct. 2019; Malaise trap; Xia Yang leg.; IEHBR, DLU 15117.

#### Paratypes

CHINA • 1 ♀; Yunnan, Lijiang district, Mt Laojun; 26°45′50.6″ N, 99°40′04.3″ E; 2400 m a.s.l.; 27 Aug.–17 Sep. 2022; Malaise trap; Xia Yang leg.; IEHBR, DLU 17923 • 1 ♀; same data as for preceding; ISAS, DLU 17958.

### Description

#### Female


**HEAD.** Antenna with 33 flagellomeres, slender, slightly broadened beyond middle, not flattened ventrally, with white ring dorsally (Fig. 8). Basal flagellomeres elongate, first flagellomere 5.5 × as long as broad at apex. Eye rather large (Fig. 9A–B, D). Face 1.8 × as broad as high (Fig. 9A), without punctures, shagreened, weakly protruding in profile (Fig. 9B). Space between scapes with impression (Fig. 9A). Clypeus separated from face by weak impression, rounded, slightly convex in profile (Fig. 9B), 1.9 × as broad as high (Fig. 9A), sculptured as face. Its apical margin sharp. Labrum visible (Fig. 9A). Mandible strongly narrowed from base to apex, sharply carinated dorsally and ventrally (Fig. 9A–B). Malar space 0.9 × as long as basal mandibular width. Gena short, shagreened (Fig. 9B). Head rather transverse, sharply narrowed behind eyes, 0.3 × as long as broad (Fig. 9D). Vertex sloping down right after ocelli (Fig. 9D), shagreened. POL:OD:OOL = 1.2:1:1.1. Occipital carina angularly impressed medially, lower end interrupted, not reaching base of mandible or hypostomal carina (Fig. 9B).

**MESOSOMA.** Pronotal collar short (Fig. 9B). Mesoscutum rather strongly convex, slightly longer than broad, densely punctate on shagreened microsculpture, matt (Fig. 9H). Notauli shallowly impressed at anterior third (Fig. 9H). Subtegular ridge defined. Speculum shagreened (Fig. 9G). Mesopleural pit consisting of short, deep horizontal groove (Fig. 9G). Sternaulus rather vestigial, visible at anterior part. Mesopleuron not punctured, shagreened, matt (Fig. 9G). Scutellum high, elevated above postscutellum, carinated laterally but interrupted at apex (Fig. 9H). Hind margin of metanotum with triangular projections opposite lateral longitudinal carinae. Propodeum tetrahedral in dorsal profile, its horizontal parts equal to area petiolaris (Fig. 9E). Propodeal carinae developed (Fig. 9F). Area basalis rather long. Area superomedia hexagonal, approximately as long as broad (Fig. 9B). Lateral propodeal carina present, joining area superomedia before its middle. Apophysis short but distinct (Fig. 9E). Juxtacoxal carina distinct. Propodeal spiracles elongate oval. Legs long and slender. Hind coxa elongate, 1.5 × as long as

broad. Hind femur  $4.2\times$  as long as wide. Basal portion of hind tibia slender, apical portion expanding evenly in profile. Ratio of length of hind tarsomeres (from base to apex)  $2.1 : 1.2 : 1 : 0.5 : 0.6$ . Claws elongate, weakly curved. Fore wing with areolet large, quadrangular, receiving vein  $2m-cu$  distal to its middle (Fig. 9K). Pterostigma broad. Vein  $1cu-a$  interstitial to  $M\&RS$ . Hind wing with distal abscissa of  $Cu$  straight,  $Cu$  about 3.5 times as long as  $cu-a$ , these veins almost opposite (Fig. 9K).



**Fig. 8.** *Acantholabus yunlingensis* Reshchikov & Kikuchi sp. nov., holotype, ♀ (IEHBR, DLU 15117). Habitus in lateral view.



**Fig. 9.** *Acantholabus yunlingensis* Reshchikov & Kikuchi sp. nov., holotype, ♀ (IEHBR, DLU 15117). **A.** Face. **B.** Head in lateral view. **C.** T1 dorsal view. **D.** Head in dorsal view. **E.** Propodeum in lateral view. **F.** Propodeum in dorsal view. **G.** Mesopleuron. **H.** Mesoscutum in dorsal view. **I.** T2 in dorsal view. **J.** T4–7 and ovipositor in lateral view. **K.** Wings.

**METASOMA.** T1 without lateral carina, petiole short (Fig. 9C),  $1.6\times$  as long as postpetiole (in profile) (Fig. 8). Gastrocoeli weakly impressed and elongate. Thyridia large and broad, distant from base of T2, with rather narrow interval between them with weak impression (Fig. 9I). Hypopygium triangular ventrally, without distinct longitudinal fold (Fig. 9J). Ovipositor sheath slightly projecting behind apex of metasoma (Fig. 9J).

**COLOUR.** Body mostly brown-yellow (Fig. 8), except: antenna (excluding scape ventrally and flagellomeres 7–11 dorsally), face medially (Fig. 9A), vertex (except lateral spots at eye orbit) (Fig. 9D), gena, pronotum dorsally, mesoscutum entirely, mesopleuron apically, T2–6 (except hind margin of T6), and hind tibia dark charcoal; flagellomeres 7–11 dorsally, face (except medial narrow band), fore coxa, pronotum ventrally and dorso-posteriorly, subtegular ridge, lower hind part of mesopleuron, hind margin of T6, T7 entirely, hind margin of S5, and hind margin of hypopygium ivory (Fig. 9J).

**VARIATIONS.** Female paratypes from Mt Laojun have darker metasoma with T2 mostly dark bean coloured (the hex code #330000) (Supp. file 1) and slightly longer apophysis (ratio of its dorsal length to its basal length  $1.2\times$  vs  $0.9\times$  in the holotype) (Supp. file 2).

**Male**

Unknown

**Host**

Unknown.

**Distribution**

China: Yunan.

**Key to species of *Acantholabus* Heinrich, 1974**

1. Propodeum with short tooth-like protrusions (Fig. 7E). Temple long,  $0.7\times$  as long as width of eye (Fig. 7B–C). Hind coxa black (Fig. 6). T5–7 with narrow white hind margins (Fig. 7G) ..... *A. regressus* Heinrich, 1974  
 – Propodeum with elongate apophysis (Figs 3D, 5F, 9E). Temple short,  $0.3\text{--}0.4\times$  as long as width of eye (Figs 5B–C, 9B–C). Hind coxa reddish-brown (Figs 2, 3D, 4, 8). T6–7 with broad white hind margins (Figs 2, 3C, 5G–H) ..... 2
2. T1–2 with broad ivory apical margins (Figs 2, 3C). Scutellum with tooth-like lateral protrusions (Fig. 3D). Area superomedia short ..... *A. furciliatus* Heinrich, 1974  
 – T1–2 without yellowish colouration (Figs 4, 5G–H, 8, 9C, I). Scutellum without tooth-like lateral protrusions (Fig. 5D). Area superomedia elongate (Fig. 5E) ..... 3
3. Propodeum with large curved apophysis, slightly elongate (Fig. 5E–F). Sternaulus distinctly impressed (Fig. 4). T1 with lateral carina (Fig. 5G–H), petiole elongate,  $1.9\times$  as long as postpetiole (in profile). Body mostly dark (Fig. 4). Scape ventrally brownish-black (Fig. 5A). Face and clypeus black medially and yellowish laterally (Fig. 5A). Pronotum mostly black, with narrow ventral and dorsal margins (Fig. 5B) ..... *A. malaisei* Heinrich, 1974  
 – Propodeum with small straight apophysis, short (Fig. 9E). Sternaulus vestigial (Fig. 9G). T1 without lateral carina, petiole short (Fig. 9C),  $1.6\times$  as long as postpetiole (in profile). Body mostly pale (Fig. 8). Scape ventrally reddish-yellow (Fig. 9A). Face and clypeus mostly yellowish-white, (narrowly black medially) (Fig. 9A). Pronotum mostly white, with small dorsal-lateral pale reddish stripe (Fig. 9B) ..... *A. yunlingensis* Reshchikov & Kikuchi sp. nov.

## Discussion

The new species is morphologically close to *A. malaisei*, the type species of the genus *Acantholabus*, in scutellar and propodeal structure (see Key). The genus has an Eastern Himalayan distribution (Fig. 1A) and all species were found between 2100–2400 m above sea level. Three species from Kachin State in northern Myanmar were collected by René Edmond Malaise (Malaise 1944). At both localities the new species was collected in autumn, Malaise's specimens between late April and early June (Heinrich 1974). *Pinus yunnanensis* is the dominant tree species at both localities and is broadly distributed across the Yun Range. The new species of *Acantholabus* seems to be rather rare, since we only found three specimens from over 50 years of Malaise trap samples. The Ichneumoninae collections in the Natural History Museum, London (NHMUK) and Florida State Collection of Arthropods, Gainesville (FSCA) were checked and no specimens of *Acantholabus* were found.

The biology of the genus *Acantholabus* is unknown; therefore, it is plausible to surmise that this genus also parasitizes larval Lepidoptera Linnaeus, 1758 (potentially Geometridae), given the habitat and the biology of other Platylabini. Considering the Himalayan diversity of the family Pinaceae Spreng. ex F.Rudolphi (Farjon 2017), we can also assume a high species diversity of geometrid moths based on coniferous plant diversity (Smetacek 2008; Ashton 2016; Sanyal *et al.* 2017) and a corresponding diversity of the members of Platylabini in the region. However, our discovery is only the fourth record of a member of the tribe from Yunnan Province, the others are *Ectopoides flavoverticilis* Riedel, 2017, *Linyctus simulator* Heinrich, 1974, and *Pachyjoppa tibialis* Cameron, 1901 (Riedel 2017). Currently, only 29 species of the entire subfamily Ichneumoninae are known from Yunnan (Riedel 2023b; Yang *et al.* 2024). We hope that our new Darwin wasp discovery will encourage further studies of the insect fauna in the unique biodiversity hotspot of the Three Parallel Rivers Region of Yunnan.

## Acknowledgements

The authors are grateful to Filippo Di Giovanni (Italy) for his help with literature, Hege Vårdal (NHRS) for her great help with type specimens examination, Davide Dal Pos (University of Central Florida, USA) and Gavin Broad (NHMUK) for checking the Ichneumoninae collections in the FSCA and the NHMUK, respectively, useful discussions and manuscript reviews, Wen Xiao (IEHBR) for his tremendous support of the project, Fang Zhang and Ruxue Li (Lijiang Laojun Mountain Biodiversity Conservation Centre) for their great help during our fieldwork on the Laojun Mountain, Alexandra Viertler (Natural History Museum Basel) and Tony Hunter (National Museums Liverpool) for manuscript reviews. This study was supported by the Yunnan Intelligence Union Program (202203AM140015), and the Yunnan High Level Personnel Training Support Program (YNWR-QNBJ-2019-262).

## References

- Ashton L.A., Nakamura A., Burwell C.J., Tang Y., Cao M., Whitaker T., Sun Z., Huang H. & Kitching R.L. 2016. Elevational sensitivity in an Asian 'hotspot': moth diversity across elevational gradients in tropical, sub-tropical and sub-alpine China. *Scientific Reports* 6: 26513. <https://doi.org/10.1038/srep26513>
- Butler L. 1993. Parasitoids associated with the macrolepidoptera community at Coopers Rock State Forest, West Virginia: a baseline study. *Proceedings of the Entomological Society of Washington* 95 (3): 504–510. Available from <https://www.biodiversitylibrary.org/page/16151238> [accessed 20 Mar. 2026].
- Bradley G.A. 1978. *Parasites of Forest Lepidoptera in Canada. Part 2. Subfamily Ichneumoninae Stenopneusticae (Hymenoptera: Ichneumonidae)*. Department of Fisheries and the Environment, Canadian Forestry Service, Forestry Technical Report 26.
- Broad G.R., Shaw M.R. & Fitton M.G. 2018. Ichneumonid wasps (Hymenoptera: Ichneumonidae): their classification and biology. *Handbooks for the Identification of British Insects* 7 (12): 1–418. <https://doi.org/10.1079/9781800625471.0000>

- Farjon A. 2017. *A Handbook of the World's Conifers (2<sup>nd</sup> edition). Vol 2*. Koninklijke Brill NV, Leiden. <https://doi.org/10.1163/9789004324510>
- Heinrich G.H. 1974. Burmesische Ichneumoninae IX. Tribus Platylabini. *Annales Zoologici* 32: 103–197.
- Kikuchi N. & Konishi K. 2018. Two new species of *Dentilabus* Heinrich (Hymenoptera: Ichneumonidae: Ichneumoninae) from Japan and Korea, with redefinition of the genus. *Zootaxa* 4524: 87–96. <https://doi.org/10.11646/zootaxa.4524.1.6>
- Malaise R. 1944. Entomological results from the Swedish Expedition 1934 to Burma and British India (Hymenoptera: Tenthredinoidea). Collected by René Malaise. The Tenthredinoidea of South-Eastern Asia. Subfamily II. Selandriinae. *Arkiv för Zoologi, Stockholm* 35: 1–58.
- Riedel M. 2008. Revision der westpalaearktischen Platylabini 1. Die Gattung *Platylabus* Wesmael, 1845 (Hymenoptera, Ichneumonidae, Ichneumoninae). *Spixiana* 31 (1): 105–172.
- Riedel M. 2017. Contribution to the Ichneumoninae (Hymenoptera, Ichneumonidae) of Southeastern Asia: 4. Platylabini, Eurylabini, and Oedicephalini. *Linzer biologische Beiträge* 49 (2): 1275–1307.
- Riedel M. 2023a. Contribution to the knowledge of the Ichneumoninae (Hymenoptera, Ichneumonidae) from Maritime Southeast Asia. *Zootaxa* 5363: 1–94. <https://doi.org/10.11646/zootaxa.5363.1.1>
- Riedel M. 2023b. Notes on Ichneumoninae (Hymenoptera: Ichneumonidae) from Southern China, with descriptions of one new genus and twelve new species. *Contributions to Entomology* 73: 223–248. <https://doi.org/10.3897/contr.entomol.73.e107542>
- Riedel M., Schmidt K. & Zmudzinski F. 2013. Beiträge zur Kenntnis der badischen Schlupfwespenfauna (Hymenoptera, Ichneumonidae) 11. Nachträge und Berichtigungen. *Carolinéa – Beiträge zur naturkundlichen Forschung in Südwestdeutschland* 71: 25–53.
- Roman A. 1939. Nordische Ichneumoniden – und einige andere. *Entomologisk Tidskrift* 60: 176–205.
- Sanyal A.K., Dey P., Uniyal V.P., Chandra K. & Raha A. 2017. Geometridae Stephens, 1829 from different altitudes in Western Himalayan Protected Areas of Uttarakhand, India (Lepidoptera: Geometridae). *SHILAP* 45 (177): 143–163. <https://doi.org/10.57065/shilap.978>
- Shaw M.R., Kan P. & Kan-van Limburg Stirum B. 2015. Emergence behaviour of adult *Trogus lapidator* (Fabricius) (Hymenoptera, Ichneumonidae, Ichneumoninae, Heresiarchini) from pupa of its host *Papilio machaon* L. (Lepidoptera, Papilionidae), with a comparative overview of emergence of Ichneumonidae from Lepidoptera pupae in Europe. *Journal of Hymenoptera Research* 47: 65–85. <https://doi.org/10.3897/JHR.47.6508>
- Short M.W., Schmidt S. & Lukacs Z. 2002. Parasitisation rates of some parasitoids (Hymenoptera: Ichneumonidae) of the autumn gum moth (Lepidoptera: Geometridae). *The Australian Entomologist* 29: 69–72. Available from <https://www.biodiversitylibrary.org/page/62372613> [accessed 20 Mar. 2026].
- Smetacek P. 2008. Moths recorded from different elevations in Nainital District, Kumaon Himalaya, India. *Bionotes* 10: 5–15.
- Tereshkin A.M. 2009. Illustrated key to the tribes of subfamilia Ichneumoninae and genera of the tribe Platylabini of world fauna (Hymenoptera, Ichneumonidae). *Linzer biologische Beiträge* 41 (2): 1317–1608.
- van Veen J.C. 1981. The biology of *Poecilostictus cothurnatus* (Hymenoptera: Ichneumonidae) an endoparasite of *Bupalus piniarius* (Lepidoptera: Geometridae). *Annales Entomologici Fennici* 47: 77–93.
- Yang X., Huang Z.P., Cui L.W., Dal Pos D., Reshchikov A. & Xiao W. 2024. A new record of *Coelichneumon rufefemoratus* (Cameron, 1903) (Hymenoptera: Ichneumonidae) from Eastern

Himalayas, with a checklist of all the species of the genus from the Oriental region. *Journal of Insect Biodiversity* 50 (2): 32–50. <https://doi.org/10.12976/jib/2024.50.2.1>

Yoder M.J., Mikó I., Seltmann K.C., Bertone M.A. & Deans A.R. 2010. A gross anatomy ontology for Hymenoptera. *PLoS ONE* 5: e15991 (8 pages). <https://doi.org/10.1371/journal.pone.0015991>

Yu D.S., van Achterberg K. & Horstmann K. 2016. *World Ichneumonoidea 2015. Taxonomy, biology, morphology and distribution*. [Flash drive]. Taxapad®, Vancouver, Canada.

Printed versions of all papers are deposited in the libraries of two of the institutes that are members of the *EJT* consortium: Muséum national d’Histoire naturelle, Paris, France and Royal Museum for Central Africa, Tervuren, Belgium. The other members of the consortium are: Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Meise Botanic Garden, Meise, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic; The Steinhardt Museum of Natural History, Tel Aviv, Israël.

## Supplementary files

**Supp. file 1.** *Acantholabus yunlingensis* Reshchikov & Kikuchi sp. nov., paratype, ♀ (IEHBR, DLU 15117), metasoma dorsal view. <https://doi.org/10.5852/ejt.2026.1052.3263.14423>

**Supp. file 2.** *Acantholabus yunlingensis* Reshchikov & Kikuchi sp. nov., paratype, ♀ (IEHBR, DLU 15117), mesosoma lateral view. <https://doi.org/10.5852/ejt.2026.1052.3263.14425>