

This work is licensed under a Creative Commons Attribution 3.0 License.

Research article

<urn:lsid:zoobank.org:pub:26A6D035-B2D9-48B2-9350-AEB9D6299B22>

Defining species boundaries in the *Merodon avidus* complex (Diptera, Syrphidae) using integrative taxonomy, with the description of a new species

Jelena AČANSKI^{1,*}, Ante VUJIĆ², Mihajla DJAN^{3,7}, Dragana OBREHT VIDAKOVIĆ^{4,7},
Gunilla STÅHLS⁵ & Snežana RADENKOVIĆ^{6,7}

¹University of Novi Sad, BioSense Institute - Research Institute for Information Technologies in Biosystems, Trg Dr Zorana Đindjića 1, 21000 Novi Sad, Serbia.

^{2,7}Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad,
Trg Dositeja Obradovića 2, 21000 Novi Sad, Serbia.

⁵Zoology Unit, Finnish Museum of Natural History, PO Box 17, 00014 University of Helsinki, Finland.

*Corresponding author: acanskijelena@gmail.com

²Email: ante.vujic@dbe.uns.ac.rs

³Email: mihajla.djan@dbe.uns.ac.rs

⁴Email: dragana.obreht@dbe.uns.ac.rs

⁵Email: gunilla.stahls@helsinki.fi

⁶Email: snezana.radenkovic@dbe.uns.ac.rs

¹<urn:lsid:zoobank.org:author:F02841B0-AC6B-4FC4-A64A-D13A3355B079>

²<urn:lsid:zoobank.org:author:A20D5863-CF18-4BF7-BB68-0DA75D34B7A8>

³<urn:lsid:zoobank.org:author:BE29D9A9-CC69-416C-8F85-4B8AA99A6C71>

⁴<urn:lsid:zoobank.org:author:75E52F0E-EEDC-4EF1-907B-9202DBDEAC9F>

⁵<urn:lsid:zoobank.org:author:D19B1262-503A-48E2-91AE-F99D0F2EA3A3>

⁶<urn:lsid:zoobank.org:author:26DF35D9-55FA-4485-8E8C-C1F90EFE1036>

Abstract. Several recent studies have detected and described complexes of cryptic and sibling species in the genus *Merodon* (Diptera, Syrphidae). One representative of these complexes is the *Merodon avidus* complex that contains four sibling species, which have proven difficult to distinguish using traditional morphological characters. In the present study, we use two geometric morphometric approaches, as well as molecular characters of the 5'-end of the mtDNA COI gene, to delimit sibling taxa. Analyses based on these data were used to strengthen species boundaries within the complex, and to validate the status of a previously-recognized cryptic taxon from Lesvos Island (Greece), here described as *Merodon megavidus* Vujić & Radenković sp. nov. Geometric morphometric results of both wing and surstyli shape confirm the present classification for three sibling species – *M. avidus* (Rossi, 1790), *M. moenium* Wiedemann in Meigen, 1822 and *M. ibericus* Vujić, 2015 – and, importantly, clearly discriminate the newly-described taxon *Merodon megavidus* sp. nov. In addition to our geometric morphometric results, supporting characters were obtained from molecular analyses of mtDNA COI sequences, which clearly differentiated *M. megavidus* sp. nov. from the other members of the *M. avidus* complex. Molecular analyses revealed that the earliest divergence of *M. ibericus* occurred around 800 ky BP, while the most recent separation happened between *M. avidus* and *M. moenium* around 87 ky BP.

Keywords. *Merodon avidus* complex, COI, geometric morphometry, wing, surstylus.

Ačanski J., Vujić A., Djan M., Vidaković D.O., Ståhls G. & Radenković S. 2016. Defining species boundaries in the *Merodon avidus* complex (Diptera, Syrphidae) using integrative taxonomy, with the description of a new species. *European Journal of Taxonomy* 237: 1–25. <http://dx.doi.org/10.5852/ejt.2016.237>

Introduction

The genus *Merodon* Meigen, 1803 (Diptera: Syrphidae: Merodontini) has become the largest European hoverfly genus due to several recent studies describing many new taxa (Marcos-García *et al.* 2007; Popov 2010; Radenković *et al.* 2011; Vujić *et al.* 2007, 2012, 2013a, 2013b, 2015). A considerable number (37 of 57 species in southeastern Europe) of the taxa are morphologically and/or genetically cryptic, with restricted distributional ranges in particular mountain ranges or islands (Vujić *et al.* 2016). Larvae of *Merodon* species are phytophagous on the underground storage organs of plants (Amaryllidaceae, Iridaceae and Hyacinthaceae) (Hurkmans 1993; Andrić *et al.* 2014). Larvae of *Merodon avidus* (Rossi, 1790) were recently found in the bulbs of *Ornithogalum umbellatum* L. (Hyacinthaceae) (Andrić *et al.* 2014). This host plant is an important foodstuff for adults, as well as for larval development and is widespread in Europe, Southwestern Asia and North Africa. Distribution of this host plant is wider than, but overlaps with, the range of the *M. avidus* complex (Andrić *et al.* 2015; Popović *et al.* 2015).

Taxa of the *Merodon avidus* complex have been the subject of many studies in the last decade due to perceived taxonomic difficulties (Milankov *et al.* 2001, 2009; Ståhls *et al.* 2009). The *M. avidus* complex is characterized by a considerable morphological variability, especially in the coloration of the antennae, thorax, abdomen and legs (Popović *et al.* 2015). This colour variability has been explained by the differential availability of trophic resources during the larval stage (Hurkmans 1993). Popović *et al.* (2015) provided justifications for the identification of three species from the *M. avidus* complex based on new diagnostic morphological characters, records of the seasonal activity and geographical distribution of bi-voltine adults, nuclear allozyme and mtDNA COI sequence analyses, and descriptions of the ecological preferences of the taxa.

However, difficulties in distinguishing the species of this complex based on morphological characters remain, despite the numerous studies on the subject. Spring generations of *Merodon avidus* are very similar to those of *M. moenium* Wiedemann *in* Meigen, 1822 based on external morphology, and, so, are easily confused using existing diagnostic features (e.g., Milankov *et al.* 2001). Furthermore, it is important to note that species of the *M. avidus* complex are not distinguishable by traditional visual identification of the structures of male genitalia under a stereo microscope. Additionally, in their analysis of COI barcodes of all *Merodon* species from Lesvos Island (Greece), Ståhls *et al.* (2009) revealed the presence of one cryptic taxon (*Merodon* sp. nova 2, herein described as *M. megavidus* sp. nov.) within the *M. avidus* complex, but did not present morphological diagnostic characters to enable characterization of the new taxon.

In the present study, two different geometric morphometric approaches were applied to quantify wing and surstylus shape variability among all hitherto described species of the *Merodon avidus* complex, including the new fourth taxon *M. megavidus* sp. nov. Insect wing shape is highly heritable and constitutes an important character for separating species (Birdsall *et al.* 2000). Geometric morphometric analysis of the wing shape has been successfully used in taxonomic studies of multiple hoverfly taxa (Francuski *et al.* 2009; Vujić *et al.* 2013b; Nedeljković *et al.* 2013, 2015). The structure and/or shape of parts of the male genitalia are very informative and thus useful for Syrphidae taxonomy and systematics (e.g., Hippa & Ståhls 2005). Differences in male genitalia structure (particularly the surstyli and aedeagal parts) detected in morphological taxonomic studies of hoverflies have revealed them to be an

important mechanism of isolation between species (Rotheray & Gilbert 2011). However, differences in genitalia structure between closely-related species can be very small, as shown for many hoverfly genera (Dušek & Laska 1964; Hippa 1990; Nedeljković *et al.* 2013, 2015; Vujić *et al.* 2013b, 2015). Geometric morphometric analysis of surstylos shape can reveal subtle shape differences that are not detectable or quantifiable by traditional visual examination (Mutanen & Pretorius 2007). This approach has only recently been applied to the taxonomy of the Syrphidae, having been successfully implemented in the genus *Chrysotoxum* Meigen, 1803 to distinguish four species (Nedeljković *et al.* 2013, 2015). The male genitalia of *Merodon* species are sclerotized and rigid and thus stable structures, so they are well suited for geometric morphometric analysis.

Our study had three objectives: (1) to further clarify the species borders of all taxa within the *M. avidus* complex using integrative taxonomy (geometric morphometrics of wings and male surstylus shape and mtDNA COI sequences); (2) to estimate times of divergence among investigated taxa; and (3) to provide descriptions and diagnostic characters of the new species. A typological (or morphological) species concept is applied in this study, integrating all available data.

Material and methods

Studied material

A total of 444 specimens belonging to the *Merodon avidus* complex from Bulgaria, Croatia, Greece, Italy, Montenegro, Morocco, Serbia, Spain and Turkey was analysed (Fig. 1; Appendix 1). Specimens of *M. avidus* were sampled during spring, summer and autumn. All specimens were identified by Ante Vujić and Snežana Radenković, and labelled using unique codes.




Fig. 1. Map of population sampling locations of the *Merodon avidus* complex from the Western Palaearctic.

Specimens detailed in the Results are in the collections of the following institutions:

FSUNS = Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia

MZF = Finnish Museum of Natural History, Helsinki, Finland

WML = World Museum Liverpool, UK

Numbers given alongside the abbreviations FSUNS, MZH, and WML in the text refer to unique identifiers from the specimen database stored in the internal electronic database of the Faculty of Sciences, University of Novi Sad.

For the description of the new species, the terminology follows McAlpine (1981) for non-genitalic morphology and Marcos-García *et al.* (2007) for morphology of the male terminalia.

Wing morphometry

Wings of all 444 available specimens were analysed using a landmark-based geometric morphometric approach (Appendix 1). Population analysis was carried out for 418 specimens of 23 populations (Appendix 1, marked with *). Populations with small sample sizes were excluded from the analysis to avoid statistical errors. The left wing of each specimen was removed using micro-scissors and mounted in Hoyer's medium on a microscope slide. Eleven homologous landmarks were digitized at vein intersections or terminations that could be reliably identified and best represented wing shape using TpsDig 2.05 software (Rohlf 2006). Generalized least-squares Procrustes superimposition was first applied to the landmark data to remove non-shape variation in terms of location, scale and orientation, and also to superimpose the objects in a common coordinate system (Rohlf & Slice 1990; Zelditch *et al.* 2004). For the wing shape analysis, partial warp scores were calculated (Zelditch *et al.* 2004). Procrustes superimposition and partial warps were computed using the free IMP software CoordGen7.14 and CVAGen 7.14a (Sheets 2012). MorphoJ v.2.0. software (Klingenberg 2011) was used for visualization of thin-plate spline deformation. Surstylus morphometry

Shape analysis of the posterior part of the left surstylus was carried out on 125 specimens of the *M. avidus* complex using a semi-landmark geometric morphometric approach (Appendix 2). The posterior part of the left surstylus (Fig. 2A: psl) was removed using a scalpel and placed on its side in glycerol on a microscope slide, and a coverslip was placed on top of the surstylus to immobilize it. Due to a lack of homologous anatomical loci, 30 semi-landmarks were digitized using the 'resample curve by length' option in TpsDig 2.05 software (Rohlf 2006). CoordGen 7.14 with an integrated Semiland module was used for semi-landmark superimposition using a distance-minimizing protocol, which minimizes the shape differences due to the arbitrary nature of semi-landmark positions along the curve (Bookstein 1997; Zelditch *et al.* 2004).

Statistical analyses

Principal component analysis (PCA) was used to test variability of wing and surstylus shape without *a priori* defining groups. Multivariate analysis of variance (MANOVA) and Fisher LSD post-hoc tests were used to test the variability connected with shape differences between species. Additionally, canonical variates analysis (CVA) and discriminant function analysis (DA) were used to test significance in wing and surstylus shape differences, to produce distance matrices and to graphically present results. The phenetic relationships among taxa were determined by UPGMA analysis based on squared Mahalanobis distances computed from the DA applied to wing variables. All statistical analyses were conducted using Statistica for Windows (Statsoft Inc. 2015: version 12).

Molecular analysis

The 5'-end of the cytochrome c oxidase 1 (COI) gene ('barcode' region) was amplified for three specimens of *M. megavidus* sp. nov. from Lesvos Island, Greece. DNA was extracted from a leg of each

specimen using a slightly modified SDS Extraction Protocol (Chen *et al.* 2010). Genomic DNA vouchers are conserved at the Faculty of Sciences, Department of Biology and Ecology, University of Novi Sad (AU827, AU885, AU886) and the Finnish Museum of Natural History, Helsinki (MZB S532). DNA barcodes were amplified with the forward primer LCO (5'-GCTCAACAAATCATAAAGATATTGG-3') and the reverse primer HCO (5'-TAAACTTCAGGGTGACCAAAAAATCA-3') (Folmer *et al.* 1994). PCR amplifications were carried out in 20 µl reaction volumes using the following mix of components: 1 × PCR buffer (Thermo Scientific), 2.5 mM MgCl₂, 0.1 mM of each nucleotide, 1 U Taq polymerase (Thermo Scientific), 2 pmol of each primer and 50 ng template DNA. PCR conditions were: initial denaturation for 2 min at 95°C; 30 s denaturation at 94°, 30 s annealing at 49°C, 2 min extension at 72°C/30 cycles; and the final extension for 8 min at 72°C. Obtained amplicons were purified using the Exo-Sap purification protocol (Thermo Scientific). Forward sequencing using the PCR primer was conducted at the Sequencing Service Laboratory of the Finnish Institute for Molecular Medicine, Helsinki, Finland (FIMM). In order to confirm the taxonomic status of *M. megavidus* sp. nov. specimens, we created a dataset consisting of 22 DNA barcode sequences, 18 of which represented GenBank-accessioned sequences of *M. avidus*, *M. moenium* and *M. ibericus* Vujić, 2015 (6 sequences per species, accession numbers indicated in Fig. 6, and one sequence for *M. megavidus* sp. nov. (generated for the Ståhls *et al.* 2009 study), in addition to three newly-generated *M. megavidus* sp. nov. DNA barcode sequences in this study). Sequences of *Eumerus sulcitibius* Rondani, 1868 (Acc. No. KT157875) and *Platynochaetus setosus* (Fabricius, 1794) (Acc. No. KM224512) (both Merodontini) were used as outgroups. Sequence alignment was conducted using the Clustal W algorithm (Thompson *et al.* 1994), as implemented in BioEdit (Hall 1999), and final modifications were done by hand. The total length of the dataset after alignment and trimming was 634 nucleotides. We constructed a Neighbour-joining (NJ) tree, and phylogenetic trees using Maximum Likelihood (ML) and Maximum Parsimony (MP) analyses. The NJ and ML trees were constructed using MEGA version 6 (Tamura *et al.* 2013) using a Tamura-3 parameter model of nucleotide substitution with Gamma correction for among-site variation in substitution rates (estimated in the same software). MP analysis was done using NONA (Goloboff 1999), with the aid of Winclada (Nixon 2002), using the heuristic search algorithm and 1000 random addition replicates (mult x 1000); holding 100 trees per round (hold/100), max trees set to 100 000; and applying TBR branch swapping. Bootstrap values were calculated for each tree using 1000 replicates. The genetic relationships among species were also tested using a Median-Joining (MJ) network (Bandelt *et al.* 1999) generated with Network v4.6.1.3 (available from <http://www.fluxus-engineering.com/sharenet.htm>) by applying the default settings ($\epsilon = 0$ and the variable sites weighted *equally* = 10), with additional post-processing with the maximum parsimony (MP) option.

Pairwise Φ_{ST} values among species of the *Merodon avidus* complex were calculated, together with an exact test of population differentiation, using ARLEQUIN 3.5.1.2 (Excoffier & Lischer 2010). Furthermore, we calculated the genetic distance between defined species using MEGA version 6 (Tamura *et al.* 2013) in order to estimate times of divergence between genetic clusters based on uncorrected p-distances divided by the pairwise evolutionary rate/MYR as described in Pröhl *et al.* (2010). We used the pairwise sequence divergence of the COI gene, relative to the mutational rate of 2.3% per million years, as estimated for various arthropod taxa (Brower 1994).

Results

Description of new species

Family Syrphidae Latreille, 1802
Subfamily Eristalinae Newman 1834
Tribe Merodontini (Edwards, 1915)
Genus *Merodon* Meigen, 1803

Merodon megavidus Vujić & Radenković sp. nov.

[urn:lsid:zoobank.org:act:B2016D80-A7ED-4958-AE91-EE5E6A8C7C51](https://doi.org/10.1545/ejt.2016.0237)

Figs 2C–D, 3–5

Diagnosis

Medium-to large-sized species (13–18 mm); black mesoscutum with four white microtrichose longitudinal stripes; tapering orange and black abdomen with white, transverse, microtrichose bands on tergites 2–4 (exceptionally without bands on tergite 2); tarsi reddish-orange dorsally; hind femur medium wide and slightly curved (Fig. 3C–D), with very short pile posteroventrally. *Merodon megavidus* Vujić & Radenković sp. nov. belongs to the *avidus* complex (male genitalia in all species identical in shape, as on Fig. 2). *Merodon megavidus* sp. nov. can be separated from the other members of the complex by larger size, golden body pile, bright orange colour of the pale parts of legs and extremely short pile on hind femur (Fig. 3C–D). These characteristics contrast with other species from the complex, which have yellow to grayish pale body pile and longer pile on the hind femur (Fig. 3A–B).

Etymology

The name *megavidus* refers to the large size (Greek word *megas* means “large”) and great similarity with *Merodon avidus*.

Type material

Holotype

GREECE: ♂, Lesvos, Petrounta, 26 Jul. 2015, leg. A. Vujić and S. Radenković (FSUNS 10132).




Fig. 2. *Merodon megavidus* Vujić & Radenković sp. nov., male genitalia. A. Epandrium, lateral view. B. Left surstylus, anterior view. C. Hypandrium, lateral view. Abbreviations: psl = posterior surstylos lobe; asl = anterior surstylos lobe; c = cercus; ae = aedeagus; ea = ejaculatory apodeme. Scale bar = 0.5 mm.

Paratypes

GREECE: Lesvos: 1 ♀, (WML Misc 38). Agiassos: 1 ♂, 7 Jul. 2007, leg. M. Hull (WML H723); 1 ♀, 8 Jun. 2003, leg. G. Ståhls (FSUNS 04385); 1 ♀, 8 Jun. 2003, leg. G. Ståhls (MZB); 1 ♂, 16 Jun. 2004, leg. M. Hull (WML Hu-93); 3 ♂♂, 19 Jun. 2003, leg. M. Hull (WML Hu70, Hu-78, Hu-81); 1 ♂, 20 Apr. 2004, leg. M. Hull (WML Hu-92); 4 ♂♂, 20 Jun. 2004, leg. M. Hull (WML Hu-77, Hu-94-96); 5 ♂♂, 22 Jun. 2004, leg. M. Hull (WML Hu-76, Hu-97-100); 2 ♂♂, 22 Jun. 2004, leg. M. Hull (FSUSN 04390, 04391); 2 ♂♂, 22 Jun. 2009, leg. M. Hull (WML H1432, H1433); 1 ♀, 23 Jun. 1999, leg. M. Hull (FSUNS 04392); 3 ♂♂, 23 Jun. 2003, leg. M. Hull (WML Hu-69, Hu-79, Hu-80). Vatoussa: 16 ♂♂, 10 ♀♀, 26 Jul. 2015, leg. A. Vujić and S. Radenković (FSUNS 10133-10151, 10159, 10160, 10162, 10163, 10243, 10253, 10256); 1 ♂, 1–4 Jun. 2012, leg. Nakas (FSUNS Ć94). Plomari: 2 ♂♂, 14 Jul. 2004, (FSUNS 02325, 03966); 1 ♂, 14. Jul. 2004, leg. H. Dahm (S532).




Fig. 3. Hind leg, lateral view. **A–B.** *Merodon avidus* (Rossi, 1790). **A.** ♂. **B.** ♀. — **C–D.** *M. megavidus* Vujić & Radenković sp. nov. **C.** ♂. **D.** ♀. Scale bar = 1 mm.

Description

Male (Figs 2C, 3, 4A, 5A)

HEAD (Fig. 4A). Antenna (Fig. 4A) orange, first flagellomere 1.8–2.0 times as long as wide, 2.0 times longer than pedicel, concave, apex acute; arista: second, third and basal part of fourth flagellomeres pale, fourth flagellomere dark brown in apical $\frac{2}{3}$ and thickened basally, 1.4 times longer than first flagellomere; with short, dense microtrichia. Face and frons black, covered with long golden pile and silver, dense microtrichia. Oral margin shiny black, except for the lateral microtrichose areas (Fig. 4A). Vertical triangle isosceles, shiny black except in front of the anterior ocellus that has pale microtrichia, covered with long orange pile except for black pile on the ocellar triangle. Ocellar triangle equilateral. Eye contiguity about 12 ommatidia long. Vertical triangle: eye contiguity: ocellar triangle = 1.5 : 0.7 : 1. Eye pile dense, white. Occiput with orange pile, along the eye margin with dense white microtrichia and posteriorly with metallic, bluish-greenish lustre.

THORAX. Mesoscutum and scutellum black with bronze lustre, covered with relatively long, dense, erect golden pile. Side of mesoscutum above wing-base with a patch of black pile. Mesoscutum with two lateral and two submedian, longitudinal, white microtrichose stripes. Proepimeron, posterior anepisternum, anteroventral and posterodorsal part of katepisternum, anepimeron, metasternum and katatergite with long golden pile and grey-green microtrichia. Wing hyaline, with dense microtrichia; veins dark brown except for light brown C, Sc and R1. Calypter pale yellow. Haltere with light brown pedicel and yellow capitulum. Legs orange, except for the black basal $\frac{3}{4}$ of the front- and mid-femora. Pile on legs golden. Hind femur (Fig. 3C) moderately thickened and curved, about 3.6 times as long as deep. Pile on hind femur very short.

ABDOMEN (Fig. 5A). Dark with white microtrichose bands, tapering, 1.4 times longer than mesonotum (including scutellum). Tergites orange and reddish except for black tergite 1 and central parts of tergites 2–3 (and 4) (Fig. 5A); orange-reddish parts of variable size on tergite 3 and 4, laterally and along microtrichose bands. Tergites 2–4 each with a pair of white microtrichose marks (exceptionally absent only on tergite 2); tergites 3–4 with wide, oblique bands (Fig. 5A). Pile on tergites golden. Sternites translucent, orange to brown towards the tip of the abdomen, covered with long yellow pile.




Fig. 4. *Merodon megavidus* Vujić & Radenković sp. nov., head, antero-lateral view. A. ♂. B. ♀. Scale bar = 1 mm.

Male genitalia (Fig. 2). Similar to all species of the *M. avidus* complex. Anterior lobe of surstylos broad and hairy (Fig. 2A–B); posterior lobe of surstylos ellipsoidal at ventral margin (Fig. 2A–B); cercus rectangular, without prominences (Fig. 2A). Hypandrium elongate and sickle-shaped, without lateral projections (Fig. 2C); lingula long (Fig. 2C).

Female (Figs 2D, 4B, 5B)

Similar to the male except for typical sexual dimorphism and for the following characteristics: first flagellomere broader and longer; frons with two wide (about 0.34 width of frons) lateral silver microtrichose longitudinal stripes; frons in the widest part about 0.25 width of head; white microtrichose longitudinal stripes on mesoscutum more visible; broad stripe of black pile between wing bases; tergites predominately red except for tergite 1 and darkened parts of tergites 2–4 in some specimens (Fig. 5B); white, microtrichose, transverse bands on tergites 3–4 (Fig. 5B); tergites 2–3 with black pile on dark parts; white microtrichose bands solely with pale pile.

Remarks

This species was mentioned as *Merodon* sp. nova 2 in Ståhls *et al.* (2009) and Ricarte *et al.* (2012). Ståhls *et al.* (2009) assumed it to be a distinct species based on the different COI barcode sequences



Fig. 5. *Merodon megavidus* Vujić & Radenković sp. nov., abdomen, dorsal view. **A.** ♂. **B.** ♀. Scale bar = 1 mm.

obtained from two specimens morphologically similar to *M. avidus* taken from Lesvos Island. In the key prepared as supporting material for Ståhls *et al.* (2009), *M. avidus* and *M. sp. nova* 2 key out together, without any morphological differences being described.

Distribution and habitat data

Lesvos Island (Greece). Maquis shrubland.

Species delimitation

Molecular data

Merodon megavidus sp. nov. clearly differs from other members of *M. avidus* complex based on our barcoding fragment of COI. All conducted phylogenetic analyses resulted in similar tree topologies (Figs 6–7; Appendix 3). Sequences from the *Merodon avidus* complex formed three separate clusters: one cluster represented *M. ibericus*, a second comprised all *M. megavidus* sp. nov. sequences with highly-significant bootstrap values (100), the third grouped sequences of *M. avidus* and *M. moenium* together. All four *M. megavidus* sp. nov. sequences were identical, defining one haplotype unique to the species. The number of mutational steps between *M. megavidus* sp. nov. and *M. avidus* is 14, with 13 and 22 mutational steps between *M. megavidus* sp. nov. and *M. moenium* and *M. ibericus*, respectively (Fig. 8).

Our UPGMA tree based on genetic distances among species revealed genetic relationships among four taxa of the *Merodon avidus* complex that support and strengthen the branch positions in wing and surstylus phenograms described below (see Fig. 9).

Significant pairwise genetic divergence (ϕ_{ST} value) was detected in each pairwise comparison between *M. megavidus* sp. nov. and each of the other species *M. ibericus*, *M. avidus* and *M. moenium* (0.420, 0.492 and 0.529, respectively). Sequence divergence (uncorrected p distance) of the COI gene was used to




Fig. 6. Maximum parsimony strict consensus tree based on DNA barcode COI sequences. Length 136 steps, Consistency Index (CI) = 93, Retention Index (RI) = 95. Filled circles denote unique changes, open circles non-unique.

Table 1. Below diagonal - pairwise p distances; above diagonal - estimated divergence times in years.

	<i>M. megavidus</i> sp. nov.	<i>M. ibericus</i>	<i>M. avidus</i>	<i>M. moenium</i>
<i>M. megavidus</i> sp. nov.	–	826.000	521.000	478.000
<i>M. ibericus</i>	0.038	–	800.000	760.000
<i>M. avidus</i>	0.024	0.037	–	87.000
<i>M. moenium</i>	0.022	0.0035	0.004	–

assess relative divergence times between the four *Merodon* taxa (Table 1), indicating an initial separation of *M. ibericus* from the rest of the complex around 800 ky BP. Divergence between *M. megavidus* sp. nov. and *M. avidus/M. moenium* occurred around 500 ky BP. The most recent separation happened between *M. avidus* and *M. moenium*, around 87 ky BP.

Geometric morphometrics - wing shape analysis

Principal component analysis conducted on 444 specimens of the *M. avidus* complex revealed six principal components (PC) that together explained 63.9% of total wing shape variability. A MANOVA with Fisher LSD post-hoc test showed that variability reflected shape changes among the investigated




Fig. 7. Maximum Likelihood tree based on a 5' fragment of COI mtDNA sequences from the *Merodon avidus* complex. Bootstrap values (1000 replicates) are shown next to the branches. The tree is drawn to scale, with branch lengths proportional to the number of substitutions per site.

species in all six PCs (MANOVA: $F_{18,1230} = 38.56$; $p < 0.00000$). Further, DA showed that all species differ highly significantly in wing shape ($p < 0.00000$), and correctly classified species with an overall classification success of 90%. All specimens of *M. megavidus* sp. nov. were correctly classified, while the lowest classification success was for *M. ibericus* (78%). Specimens belonging to *M. avidus* were correctly classified with 89% and *M. moenium* with 93% certainty. Canonical analysis produced three canonical axes (CV) related to wing shape differences (Fig. 10). CV1 separated *M. avidus* from *M. moenium* with 58% of total variation (CV1: Wilks' Lambda = 0.145440; $\chi^2 = 832.8937$; $p < 0.00000$), while CV2 separated *M. avidus* and *M. moenium* from *M. megavidus* sp. nov. and *M. ibericus* with 27% of total variation (CV2: Wilks' Lambda = 0.388320; $\chi^2 = 408.6395$; $p < 0.00000$) (Fig. 10A). *Merodon moenium* and *M. ibericus* were clearly separated according to CV3 (CV3: Wilks' Lambda = 0.698600; $\chi^2 = 154.9484$; $p < 0.00000$) (Fig. 10B). The phenogram constructed based on squared Mahalanobis distances clearly depicts that *M. avidus* and *M. moenium* have more similar wing shapes than *M. ibericus* and *M. megavidus* sp. nov. (Fig. 11). Differences in wing shape among species are depicted in Figure 12, but have been exaggerated five-fold to make them more visible.

Additionally, the phenogram constructed based on squared Mahalanobis distances of wing shape showed that all 26 analysed populations grouped according to species (Fig. 13).

Geometric morphometrics - surstylus shape

PCA of surstylus shape revealed seven PC, of which the first six were connected with shape differences among species (MANOVA: $F_{21,333} = 5.280113$; $p < 0.00000$). DA showed that all species differ highly significantly in surstylus shape ($p < 0.00000$). All specimens of *M. avidus* and *M. megavidus* sp. nov. were correctly classified (100%), while only two specimens of *M. ibericus* and *M. moenium* were misclassified (97%). CVA found three CVs connected with shape change (Fig. 14). CV1 clearly separated *M. avidus* from *M. megavidus* sp. nov., *M. moenium* and *M. ibericus* and represented 53% of total shape variability (CV1: Wilks' Lambda = 0.013392; $\chi^2 = 405.4285$; $p < 0.00000$) (Fig. 14A). The second canonical axis clearly separated *M. megavidus* sp. nov. from *M. moenium* and *M. ibericus* and was responsible for 25%




Fig. 8. Median-joining network of the mtDNA 5'-end of the COI gene. Circle sizes are proportional to haplotype frequencies. Each branch represents one mutational step; if more than one mutational step is present, it is denoted by the given number.




Fig. 9. UPGMA tree based on pairwise genetic distances for four species from the *Merodon avidus* complex.




Fig. 10. Differences in wing shape among species of the *M. avidus* complex. **A.** Scatter plot of individual scores of CV1 vs CV2. **B.** Scatter plot of individual scores of CV2 vs CV3.




Fig. 11. UPGMA phenogram constructed using the squared Mahalanobis distances of wing shape for species of the *M. avidus* complex.

of the variability (CV2: Wilks' Lambda = 0.086702; $\chi^2 = 229.8565$; $p < 0.00000$) (Fig. 14A). *Merodon moenium* and *M. ibericus* were separated by CV3, with 18% of total shape variability (CV3: Wilks' Lambda = 0.346752; $\chi^2 = 99.5597$; $p < 0.000305$) (Fig. 14B). According to the phenogram constructed based on squared Mahalanobis distances, *M. megavidus* sp. nov. has a more distinct surstylus shape, while the surstyli of *M. moenium* and *M. ibericus* are the most similar (Fig. 15A). The main shape differences among all three species lie in the posterior margin of the posterior part of the surstylus lobe (Fig. 15A).

Discussion

Differentiation and taxonomic status of the cryptic species of the *Merodon avidus* complex has been controversial for many years. An initial molecular analysis showed that *M. avidus* is a genetically and geographically structured taxon with at least two cryptic species, which were designated as *M. avidus* A and *M. avidus* B (Milankov *et al.* 2001). Further investigations based on the mtDNA COI marker expanded on this by finding two additional cryptic species from the Iberian Peninsula (*M. bicolor* Gil Collado, 1930, now *M. ibericus*) and Lesvos Island (Greece) (*M. sp. nova* 2, now *M. megavidus* sp. nov.) (Milankov *et al.* 2009, Ståhls *et al.* 2009). In Milankov *et al.* (2009), wing shape analysis could not distinguish taxa that could be differentiated by allozyme loci, even though these taxa differed in wing size. Also, they found a great deal of similarity between allopatric metapopulation pairs of *M. avidus*




Fig. 12. Thin-plate spline deformation grids showing wing shape differences between analysed species. Differences between the species have been exaggerated five-fold to make them more visible.

A and *M. avidus* B; populations of *M. avidus* A from Macedonia and the Pannonian plain overlapped with *M. avidus* B populations from Durmitor, Stara Planina and Kopaonik in terms of wing shape. Only a population of *M. avidus* A from Morinj (Montenegro) was clearly separated from other populations. Subsequent analysis showed that this substantial overlap in wing shape was a result of incorrect species identifications due to the morphological similarities between spring generations of *M. avidus* and *M. moenium* (Popović *et al.* 2015). Previously, it had been considered that *M. moenium* (*M. avidus* B) was a species distributed on mountainous regions of the Balkan Peninsula. However, it is now known that while *M. moenium* is predominantly distributed in the continental parts of Europe, it also occurs in some parts of the Mediterranean basin and the Black Sea coast, where it can occur sympatrically with *M. avidus*. Further, allozyme analysis of morphologically different spring and autumn generations from Umag (Croatia) clarified this misclassification issue by showing that both morphotypes belonged to *M. avidus* (Popović *et al.* 2015).

Our molecular analysis using the barcoding COI fragment for an additional three specimens from Lesvos Island supports the presence of a morphologically cryptic taxon (*M. megavidus* sp. nov.), previously revealed by Ståhls *et al.* (2009). All our phylogenetic analyses placed *M. megavidus* sp. nov. as a separate independent cluster from *M. ibericus* and the *M. avidus*/*M. moenium* branch (Figs 6–9). According to estimated divergence times, all diversification occurred in the Pleistocene (2.6 to 0.0117 MYA). This geological epoch was marked by repeated (at least 20) glacial and interglacial periods, which influenced speciation and the distributions of many recent taxa in Europe (Julius & Kukla 1977; Martinson *et al.* 1987; Perissoratis & Conispoliatis 2003). Recent studies have revealed many examples of insect species that have altered their ranges and/or evolved as a response to repeated isolation during glacial-interglacial cycles across three large peninsulas of Southern Europe (Hewitt 2001; Konstantinov *et al.* 2009; Dapporto 2010; Nicholls *et al.* 2010; Zhu *et al.* 2013).




Fig. 13. UPGMA phenogram constructed using squared Mahalanobis distances of wing shape for populations of species of the *M. avidus* complex.

The first mitochondrial diversification in the *M. avidus* complex took place in the Calabrian stage of the Early Pleistocene (around 800 ky BP), when *M. ibericus* diverged from a common ancestor. The Günz-Mindel interglacial corresponds to the approximate period when separation of *M. megavidius* sp. nov. from the *M. avidus/M. moenium* lineage occurred. The most recent diversification, i.e., between *M. avidus* and *M. moenium*, most likely took place at the end of the Riss-Würm interglacial or the beginning of the Würm glaciation period. The low species resolution ability of the 3' and 5' fragments of COI to separate the *M. avidus* and *M. moenium* lineages (Milankov *et al.* 2009; Popović *et al.* 2014, 2015) can be explained by recent speciation and by the founder effect associated with postglacial recolonization of northern Europe (Shikano *et al.* 2010). Both species share an identical haplotype of the COI 3'-end (Popović *et al.* 2014), although the 5'-end fragment haplotypes of DNA barcodes are not shared, they give low support in cluster analyses (Popović *et al.* 2015).

The most probable scenario for postglacial colonization of Europe is expansion of the *M. avidus* and *M. moenium* lineages from the Balkan Peninsula into Central Europe (a “grasshopper” pattern of colonization according to Hewitt 1999). The Pyrenees acted as a significant geographical barrier preventing the dispersal of *M. ibericus* to other European areas.

Our geometric morphometric analyses show that all investigated species of the *M. avidus* complex can be successfully discriminated based on wing and surstylus shape. The high overall success rate of classification indicates that wings, and especially surstylus shape, have meaningful interspecific discriminatory power. It is important to emphasize that, contrary to previous geometric morphometric studies, we found highly significant differences in wing shape between *M. avidus* and *M. moenium*. The phenogram based on squared Mahalanobis distances for wings is congruent with the UPGMA tree based on genetic distances among species. Wing shape, as a highly heritable structure, has greater importance in insect taxonomy than wing size (Birdsall *et al.* 2000), which has been confirmed by earlier studies of hoverflies in which wing shape has been successfully used for identification and delimitation of species (Vujić *et al.* 2013b; Nedeljković *et al.* 2013, 2015).

Additional evidence for species distinctiveness is provided by the highly significant differences in surstylus shape between *M. avidus*, *M. ibericus*, *M. megavidius* sp. nov. and *M. moenium*. The main differences in surstylus shape are connected to the posterior margin of the posterior part of the surstylus




Fig. 14. Differences in the posterior part of the surstylus among species of the *M. avidus* complex. **A.** Scatter plot of individual scores of CV1 vs CV2. **B.** Scatter plot of individual scores of CV2 vs CV3.

lobe, which is involved in gripping the female during copulation. It could be hypothesized that these shape differences contribute to a mechanism of sexual isolation, especially since *M. avidus* has a distinct surstylos shape compared to partly sympatric *M. moenium*. Although the exact mechanism of sexual isolation in Syrphidae is not known, traditionally the shape of male genitalia is deemed a significant mechanism of isolation between species (Rotheray & Gilbert 2011). Given that the morphology of male genital structures is considered one of the fastest evolving traits in animal groups with internal fertilization (Soto *et al.* 2013), and that insect genitalia are conspicuously variable even in closely related taxa that are otherwise morphologically very similar (Hosken & Stockley 2004), we assert that significant differences in surstylos shape is strong evidence for species delimitation. Recent studies have found that natural and sexual selection (and their interaction) caused insect genital evolution (Hasson *et al.* 2009; House *et al.* 2013), and that the degree of morphological differences in the structures of genitalia is associated with geographic distance (Soto *et al.* 2013). Despite their recent speciation and geographic proximity, *M. avidus* does not exhibit a similar surstylos shape to *M. moenium*. Other studies have found that other factors, such as specific host plant relationships Soto (2012) or sexual selection (Hosken & Stockley 2004), may promote further evolution in genital morphology.

Our population-level morphometric analysis of wing shape is in concordance with species delimitation findings. All populations clustered according to species. Furthermore, in this analysis, *M. megavidus* sp. nov. from Lesvos Island is distinct from all other species and/or populations, and it is important to underline that this taxon was clearly differentiated from population(s) of *M. avidus* also from Lesvos Island.




Fig. 15. Differences in the posterior part of the surstylos among species of the *M. avidus* complex. **A.** UPGMA phenogram constructed using squared Mahalanobis distances. **B.** Thin-plate spline deformation grids showing overall shape differences between analysed species.

Acknowledgements

We thank the curators of the several museums listed above in the “Studied material” section that facilitated visits and loaned specimens for study. We thank John O’Brien for English revision. Financial support was provided by the Serbian Ministry of Education, Science and Technological Development (projects OI173002, III43002 and OI1613001), the Provincial Secretariat for Science and Technological Development (project ‘Genetic resources of agro-ecosystems in Vojvodina and sustainable agriculture’),

the Transnational Access to Research Infrastructures activity in the FP7 of the EC (ExpeER project, TA visit ‘STEPS’), the FP7 EU project, Innosense, and National Strategic Reference Framework (NSRF)—Research Funding Program: THALES.

References

- Andrić A., Šikoparija B., Obreht D., Đan M., Preradović J., Radenković S., Pérez-Bañón C. & Vujić A. 2014. DNA barcoding applied: identification of the larva of *Merodon avidus* (Diptera: Syrphidae). *Acta Entomologica Musei Nationalis Praga* 54 (2): 741–757. <http://zoobank.org/urn:lsid:zoobank.org:pub:5CB81C03-0C5F-4220-AD2B-1C97F74C30D3>
- Andrić A., Kočić-Tubić N., Rat M. & Obreht-Vidaković D. 2015. Diversity and genetic structure of *Ornithogalum* L. (Hyacinthaceae) populations as revealed by RAPD-PCR markers. *Genetika* 47 (1): 275–288. <http://dx.doi.org/10.2298/GENS1501275A>
- Bandelt H.J., Forster P. & Röhl A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* 16 (1): 37–48. <http://dx.doi.org/10.1093/oxfordjournals.molbev.a026036>
- Birdsall K., Zimmerman E., Teeter K. & Gibson G. 2000. Genetic variation for the positioning of wing veins in *Drosophila melanogaster*. *Evolution and Development* 2: 16–24. <http://dx.doi.org/10.1046/j.1525-142x.2000.00034.x>
- Bookstein F.L. 1997. Landmark methods for forms without landmarks: morphometrics of group differences in outline shape. *Medical Image Analysis* 1 (3): 225–243. [http://dx.doi.org/10.1016/S1361-8415\(97\)85012-8](http://dx.doi.org/10.1016/S1361-8415(97)85012-8)
- Brower A.V. 1994. Rapid morphological radiation and convergence among races of the butterfly *Heliconius erato* inferred from patterns of mitochondrial DNA evolution. *Proceedings of the National Academy of Sciences* 91 (14): 6491–6495. <http://dx.doi.org/10.1073/pnas.91.14.6491>
- Chen H., Rangasamy M., Tan S.Y., Wang H. & Siegfried B.D. 2010. Evaluation of five methods for total DNA extraction from Western Corn Rootworm Beetles. *PLoS ONE* 5: e11963. <http://dx.doi.org/10.1371/journal.pone.0011963>
- Dapporto L. 2010. Speciation in Mediterranean refugia and post-glacial expansion of *Zerynthia polyxena* (Lepidoptera, Papilionidae). *Journal of Zoological Systematics and Evolutionary Research* 48 (3): 229–237. <http://dx.doi.org/10.1111/j.1439-0469.2009.00550.x>
- Dušek J. & Laska P. 1964. A contribution to distinguishing the European species of the subgenus *Syrphus* Fabricius (Diptera, Syrphidae) according to male genitalia and larvae. *Acta Societatis Entomologicae Cechosloveniae* 61 (1): 58–70.
- Excoffier L. & Lischer H. 2010. Arlequin Ver 3.5. 1.2 User Manual (2010) Computational and molecular population genetics lab (CMPG). Institute of Ecology and Evolution University of Bern, Bern.
- Folmer O., Black M., Hoeh W., Lutz R. & Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3 (5): 294–299.
- Francuski Lj., Ludoški J., Vujić A. & Milankov V. 2009. Wing geometric morphometric inferences on species delimitation and intraspecific divergent units in the *Merodon ruficornis* group (Diptera, Syrphidae) from the Balkan Peninsula. *Zoological Science* 26 (4): 301–308. <http://dx.doi.org/10.2108/zsj.26.301>
- Goloboff P. 1999. NONA (NO NAME) ver. 2. Published by the author, Tucumán, Argentina.

- Hall T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/. *Nucleic Acids Symposium Series* 41: 95–98.
- Hasson E., Soto I.M., Carreira V.P., Corio C., Soto E.M. & Betti M. 2009. Host plants, fitness and developmental instability in a guild of cactophilic species of the genus *Drosophila*. In: Eduardo B.S. (ed.) *Ecotoxicology Research Developments*: 89–109. Nova Science Publishers, New York.
- Hewitt G.M. 1999. Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society* 68 (1–2): 87–112. <http://dx.doi.org/10.1006/bijl.1999.0332>
- Hewitt G.M. 2001. Speciation, hybrid zones and phylogeography - or seeing genes in space and time. *Molecular Ecology* 10 (3): 537–549. <http://dx.doi.org/10.1046/j.1365-294x.2001.01202.x>
- Hippa H. 1990. The genus *Milesia* Latreille (Diptera : Syrphidae). *Acta Zoologica Fennica* 187: 1–226.
- Hippa H. & Ståhls G. 2005. *Morphological characters of adult Syrphidae: descriptions and phylogenetic utility*. Finnish Zoological and Botanical Publishing Board, Helsinki.
- Hosken D.J. & Stockley P. 2004. Sexual selection and genital evolution. *Trends in Ecology & Evolution* 19 (2): 87–93. <http://dx.doi.org/10.1016/j.tree.2003.11.012>
- House C.M., Lewis Z., Hodgson D.J., Wedell N., Sharma M.D., Hunt J. & Hosken D.J. 2013. Sexual and natural selection both influence male genital evolution. *PLoS ONE*: 8 (5): e63807. <http://dx.doi.org/10.1371/journal.pone.0063807>
- Hurkmans W. 1993. A monograph of *Merodon* (Diptera: Syrphidae). Part 1. *Tijdschrift voor Entomologie* 136: 147–234.
- Julius F. & Kukla J.G. 1977. Pleistocene climates in Central Europe: at least 17 interglacials after the Olduvai event. *Quaternary Research* 7 (3): 363–371.
- Klingenberg C.P. 2011. MORPHOJ: an integrated software package for geometric Morphometrics. v.2.0. [Computer software and manual]. *Molecular Ecology Resources* 11 (2): 353–357. <http://dx.doi.org/10.1111/j.1755-0998.2010.02924.x>
- Konstantinov A.S., Korotyaev B.A. & Volkovitsh M.G. 2009. Insect biodiversity in the Palearctic Region. In: Foottit R.G. & Adler P.H. (eds) *Insect Biodiversity: Science and Society*: 107–162. Wiley-Blackwell, Oxford.
- Marcos-García M.A., Vujić A. & Mengual X. 2007. Revision of Iberian species of the genus *Merodon* (Diptera: Syrphidae). *European Journal of Entomology* 104 (3): 531–572. <http://dx.doi.org/10.14411/eje.2007.073>
- Martinson D.G., Pisias N.G., Hays J.D., Imbrie J., Moore T.C. & Shackleton N.J. 1987. Age, dating and orbital theory of the Ice Ages: development of a high resolution 0–300,000 year chronostratigraphy. *Quaternary Research* 27 (1): 1–29. [http://dx.doi.org/10.1016/0033-5894\(87\)90046-9](http://dx.doi.org/10.1016/0033-5894(87)90046-9)
- McAlpine J. F. 1981. Morphology and terminology, adults. In: McAlpine J.F., Peterson B.V., Shewell G.E., Teskey H.J., Vockeroth J.R. & Wood D.M. (eds) *Manual of Nearctic Diptera*: 9–63. Agriculture Canada, Ottawa.
- Milankov V., Vujić A. & Ludoski J. 2001. Genetic divergence among cryptic taxa of *Merodon avidus* (Rossi, 1790) (Diptera: Syrphidae). *An International Journal of Dipterological Research* 12 (1): 15–24.
- Milankov V., Ludoški J., Ståhls G., Stamenković J. & Vujić A. 2009. High molecular and phenotypic diversity in the *Merodon avidus* complex (Diptera, Syrphidae): cryptic speciation in a diverse insect taxon. *Zoological Journal of the Linnean Society* 155 (4): 819–833. <http://dx.doi.org/10.1111/j.1096-3642.2008.00462.x>

- Mutanen M. & Pretorius E. 2007. Subjective visual evaluation vs. traditional and geometric morphometrics in species delimitation: a comparison of moth genitalia. *Systematic Entomology* 32 (2): 371–386. <http://dx.doi.org/10.1111/j.1365-3113.2006.00372.x>
- Nedeljković Z., Ačanski J., Vujić A., Obreht D., Djan M., Ståhls G. & Radenković S. 2013. Taxonomy of *Chrysotoxum festivum* Linnaeus, 1758 (Diptera: Syrphidae) - an integrative approach. *Zoological Journal of the Linnean Society* 169 (1): 84–102. <http://dx.doi.org/10.1111/zoj.12052>
- Nedeljković Z., Ačanski J., Đan M., Obreht-Vidaković D., Ricarte A. & Vujić A. 2015. An integrated approach to delimiting species borders in the genus *Chrysotoxum* Meigen, 1803 (Diptera: Syrphidae), with description of two new species. *Contributions to Zoology* 84 (4): 285–304. Available from <http://www.ctoz.nl/vol84/nr04/a02> [accessed 19 Sep. 2016]
- Nicholls J.A., Preuss S., Hayward A., Melika G., Csóka G.Y., Nieves-Aldrey J., Askew R.R., Tavakoli M., Schönrogge K. & Stone G.N. 2010. Concordant phylogeography and cryptic speciation in two Western Palaearctic oak gall parasitoid species complexes. *Molecular Ecology* 19 (3): 592–609. <http://dx.doi.org/10.1111/j.1365-294X.2009.04499.x>
- Nixon K.C. 2002. WinClada ver. 1.0000. Published by the author, Ithaca, NY, USA.
- Perissoratis C. & Conispoliatis N. 2003. The impacts of sea-level changes during latest Pleistocene and Holocene times on the morphology of the Ionian and Aegean seas (SE Alpine Europe). *Marine Geology* 196 (3): 145–156. [http://dx.doi.org/10.1016/S0025-3227\(03\)00047-1](http://dx.doi.org/10.1016/S0025-3227(03)00047-1)
- Popov G.V. 2010. *Merodon alexandri* spec. nov. – a new species of hoverfly (Diptera: Syrphidae) from the northern Black Sea Region. *Studia Dipterologica* 16: 133–151.
- Popović D., Djan M., Šašić L., Šnjegota D., Obreht D. & Vujić A. 2014. Usage of different molecular markers in delimitation of cryptic taxa in *Merodon avidus* species complex (Diptera: Syrphidae). *Acta Zoologica Bulgararica* 7: 33–38.
- Popović D., Ačanski J., Djan M., Obreht D., Vujić A. & Radenković S. 2015. Sibling species delimitation and nomenclature of the *Merodon avidus* complex (Diptera: Syrphidae). *European Journal of Entomology* 112 (4): 790–809. <http://dx.doi.org/10.14411/eje.2015.100>
- Pröhl H., Ron S.R. & Ryan M.J. 2010. Ecological and genetic divergence between two lineages of Middle American túngara frogs *Physalaemus* (= *Engystomops*) *pustulosus*. *BMC Evolutionary Biology* 10 (1): 146. <http://dx.doi.org/10.1186/1471-2148-10-146>
- Radenković S., Vujić A., Ståhls G., Pérez-Bañón C., Rojo S., Petanidou T. & Šimić S. 2011. Three new cryptic species of the genus *Merodon* Meigen (Diptera: Syrphidae) from the island of Lesvos (Greece). *Zootaxa* 2735: 35–56.
- Ricarte-Sabater A., Nedeljković Z., Rotheray G.E., Lyszkowski R.M., Hancock E.G., Watt K., Hewitt S.M., Horsfield D. & Wilkinson G. 2012. Syrphidae (Diptera) from the Greek island of Lesvos, with description of two new species. *Zootaxa* 3175: 1–23.
- Rohlf F.J. 2006. *TpsDig – Digitize landmarks and outlines. Ver. 2.05*. [Computer software and manual]. Department of Ecology and Evolution, State University of New York at Stony Brook.
- Rohlf F.J. & Slice D.E. 1990. Extensions of the Procrustes method for the optimal superimposition of landmarks. *Systematic Zoology* 39 (1): 40–59. <http://dx.doi.org/10.2307/2992207>
- Rotheray G.E. & Gilbert F. 2011. *The Natural History of Hoverflies*. Forrest Text, Ceredigion.
- Sheets H.D. 2012. *IMP software series*. [Computer software and manual]. Buffalo, New York: Canisius College.

- Shikano T., Shimada Y., Herczeg G. & Merilä J. 2010. History vs. habitat type: explaining the genetic structure of European nine-spined stickleback (*Pungitius pungitius*) populations. *Molecular Ecology* 19 (6): 1147–1161. <http://dx.doi.org/10.1111/j.1365-294X.2010.04553.x>
- Soto I.M. 2012. Aedeagal divergence in sympatric populations of two sibling species of cactophilic *Drosophila* (Diptera, Drosophilidae): Evidence of character displacement? *Neotropical Entomology* 41 (3): 207–213. <http://dx.doi.org/10.1007/s13744-012-0028-x>
- Soto I.M., Carreira V.P., Soto E.M., Márquez F., Lipko P. & Hasson E. 2013. Rapid divergent evolution of male genitalia among populations of *Drosophila buzzatii*. *Evolutionary Biology* 40 (3): 395–407. <http://dx.doi.org/10.1007/s11692-013-9223-x>
- Ståhls G., Vujić A., Perez-Banon C., Radenkovic S., Rojo S. & Petanidou T. 2009. COI barcodes for identification of *Merodon* hoverflies (Diptera, Syrphidae) of Lesvos Island, Greece. *Molecular Ecology Resources* 9 (6): 1431–1438. <http://dx.doi.org/10.1111/j.1755-0998.2009.02592.x>
- StatSoft Inc. STATISTICA (data analysis software system), version 12 2015. Available from www.statsoft.com [accessed 20 Feb. 2016]
- Tamura K., Stecher G., Peterson D., Filipski A. & Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* 30: 2725–2729. <http://dx.doi.org/10.1093/molbev/mst197>
- Thompson J.D., Higgins D.G. & Gibson T.J. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weigh matrix choice. *Nucleic Acids Research* 22 (22): 4673–4680. <http://dx.doi.org/10.1093/nar/22.22.4673>
- Vujić A., Radenković S., Ståhls G., Ačanski J., Stefanović A., Veselić S., Andrić A. & Hayat R. 2012. Systematics and taxonomy of the *ruficornis* group of genus *Merodon* Meigen (Diptera: Syrphidae). *Systematic Entomology* 37 (3): 578–602. <http://dx.doi.org/10.1111/j.1365-3113.2012.00631.x>
- Vujić A., Radenković S., Likov L., Trifunov S. & Nikolić T. 2013a. Three new species of the *Merodon nigrifrons* group (Diptera: Syrphidae) from the Middle East. *Zootaxa* 3640 (3): 442–464. <http://dx.doi.org/10.11646/zootaxa.3640.3.7>
- Vujić A., Ståhls G., Ačanski J., Bartsch H., Bygberg R. & Stefanović A. 2013b. Systematics of Pipizini and taxonomy of European *Pipiza* Fallén: molecular and morphological evidence (Diptera, Syrphidae). *Zoologica Scripta* 42 (3): 288–305. <http://dx.doi.org/10.1111/zsc.12005>
- Vujić A., Radenković S., Ačanski J., Grković A., Taylor M., Šenol S.G. & Hayat R. 2015. Revision of the species of the *Merodon nanus* group (Diptera: Syrphidae) including three new species. *Zootaxa* 4006 (3): 439–462. <http://dx.doi.org/10.11646/zootaxa.4006.3.2>
- Vujić A., Perez-Banon C., Radenković S., Ståhls G., Rojo S., Petanidou T. & Šimić S. 2007. Two new species of genus *Merodon* Meigen, 1803 (Syrphidae, Diptera) from the island of Lesvos (Greece), in the eastern Mediterranean. *Annales de la Société Entomologique de France* 43 (3): 319–326. <http://dx.doi.org/10.1080/00379271.2007.10697527>
- Vujić A., Petanidou T., Tscheulin T., Cardoso P., Radenković S., Ståhls G., Baturan Ž., Mijatović G., Rojo S., Pérez-Bañón C., Devalez J., Andrić A., Jovičić S., Krašić D., Markov Z., Radišić D. & Tataris G. 2016. Biogeographical patterns of the genus *Merodon* Meigen, 1803 (Diptera: Syrphidae) in islands of the eastern Mediterranean and adjacent mainland. *Insect Conservation and Diversity*. <http://dx.doi.org/10.1111/icad.12156>
- Zelditch M.L., Swiderski D.L., Sheets H.D. & Fink W.L. 2004. *Geometric Morphometrics for Biologists: a Primer*. Elsevier Academic Press, London.

Zhu G., Liu G., Bu W. & Lis J.A. 2013. Geographic distribution and niche divergence of two stinkbugs, *Parastrachia japonensis* and *Parastrachia nagaensis*. *Journal of Insect Science* 13 (102). <http://dx.doi.org/10.1673/031.013.10201>

Manuscript received: 4 January 2016

Manuscript accepted: 23 May 2016

Published on: 14 October 2016

Topic editor: Gavin Broad

Desk editor: Kristiaan Hoedemakers

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d’Histoire naturelle, Paris, France; Botanic Garden Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands.

Appendix 1. List of specimens used for wing geometric morphometric analysis, by geographical area and species.

Species	Country	Population	δ	φ	Σ
	Bulgaria	Pirin Mts.*	7	1	8
	Croatia	Umag*	16	6	22
		Drama*	10	4	14
		Lesvos island*	20	1	21
	Greece	Olympus Mts.*	13	7	20
		Peloponnesos*	10	5	15
		Pindos Mts.*	11	9	20
<i>M. avidus</i> (Rossi, 1790)	Italy	Piemonte*	15	1	16
		Pisa*	16	4	20
	Montenegro	Durmitor Mts.	2	0	2
		Djerdap gorge	1	2	3
	Serbia	Malinik Mts.*	7	2	9
		Pčinja*	29	3	32
		Dubašnica Mts.	1	0	1
	Turkey	Lake Baffa	4	1	5
	Morocco	Atlas Mts.*	3	17	20
<i>M. ibericus</i> Vujić, 2015	Spain	Cádiz	2	0	2
		Sierra Nevada Mts.*	15	0	15
	Italy	Zuclo*	8	2	10
		Castiglione Dei Pepoli	6	0	6
	Montenegro	Durmitor Mts.*	14	4	18
		Djerdap gorge*	32	4	36
		Dubašnica Mts.*	16	2	18
<i>M. moenium</i>		Kopaonik Mts.*	12	4	16
Wiedemann in Meigen, 1822		Stara Planina Mts.*	18	5	23
	Serbia	Fruška Gora*	17	0	17
		Malinik Mts.*	10	0	10
		Tara Mts.*	21	0	21
		Vršačke Planine	6	0	6
<i>M. megavidus</i>	Greece	Lesvos island*	9	9	18
Vujić & Radenković sp. nov.					
Total			351	93	444

Appendix 2. List of specimens used for surstyli geometric morphometric analysis, by geographical area and species.

Species	Country	Σ
<i>M. avidus</i> (Rossi, 1790)	Croatia	7
	Greece	9
	Italy	13
	Serbia	13
<i>M. ibericus</i> Vujić, 2015	Turkey	5
	Morocco	16
<i>M. moenium</i> Wiedemann in Meigen, 1822	Spain	17
	Montenegro	8
<i>M. megavidus</i> Vujić & Radenković sp. nov.	Serbia	22
	Greece	15
Total		125

Appendix 3. Neighbor-joining tree based on a 5' fragment of COI mtDNA sequences from the *Merodon avidus* complex. Bootstrap values (1000 replicates) are shown next to the branches.

