



This work is licensed under a Creative Commons Attribution 3.0 License.

Research article

[urn:lsid:zoobank.org:pub:3645CA2E-F2CE-41FB-91EA-48999065936B](https://doi.org/10.5852/ejt.2017.307)

Contributions to the knowledge of the mite genus *Stigmaeus* Koch, 1836 (Acari: Stigmaeidae) of Turkey

Meryem BİNGÜL¹, Salih DOĞAN^{2,*} & Sibel DİLKARAOĞLU³

^{1,2,3} Biology Department, Arts & Sciences Faculty, Erzincan University, Erzincan, Turkey.

*Corresponding author: salihdogan@erzincan.edu.tr

¹ Email: mbingul@erzincan.edu.tr

³ Email: sdilkara@erzincan.edu.tr

¹ [urn:lsid:zoobank.org:author:304DEF15-095F-41FE-9C2F-7B8E1B70B413](https://doi.org/10.5852/ejt.2017.307)

² [urn:lsid:zoobank.org:author:2444C602-B47B-4D8F-B446-DCBC2AD354FB](https://doi.org/10.5852/ejt.2017.307)

³ [urn:lsid:zoobank.org:author:9D99E481-4E14-4B08-9EC8-F1F2807C54EF](https://doi.org/10.5852/ejt.2017.307)

Abstract. Based on the mite specimens collected within the scope of a study on Erzincan (Turkey) mite biodiversity, two species of the genus *Stigmaeus* are described and illustrated here: *S. bifurcus* sp. nov. as new to science and *S. miandoabiensis* Bagheri & Zarei, 2012 as a new record for Turkey. Some morphological abnormalities in the new species are noted. The deutonymph of *S. miandoabiensis* is described for the first time in this study. Discovery of this stage from soil and litter under *Pinus sylvestris* in Turkey adds more data to our knowledge of the species.

Keywords. Mite, taxonomy, *Stigmaeus*, new species, new record, description, Turkey.

Bingül M., Doğan S. & Doğan S. 2017. Contributions to the knowledge of the mite genus *Stigmaeus* Koch, 1836 (Acari: Stigmaeidae) of Turkey. *European Journal of Taxonomy* 307: 1–16. <https://doi.org/10.5852/ejt.2017.307>

Introduction

Stigmaeus Koch, 1836 is the most diverse genus in the family Stigmaeidae with 133 valid species (Fan & Zhang 2005; Doğan *et al.* 2015a, 2016; Dilkaraoğlu *et al.* 2016a, 2016b, Fan *et al.* 2016). To date, 36 species of the genus have been recorded from Turkey (Özkan *et al.* 1988, 1994; Erman *et al.* 2007; Doğan 2007; Doğan *et al.* 2015a, 2016; Dilkaraoğlu *et al.* 2016a, 2016b).

The descriptions of *Stigmaeus bifurcus* sp. nov. and *S. miandoabiensis* Bagheri & Zarei, 2012—originally found in Iran, newly recorded for the Turkish fauna—are given in this survey. The deutonymph of *S. miandoabiensis* is described for the first time.

Material and methods

The specimens were collected from Erzincan in 2014 and 2015. Methods used for specimen collection, extraction, material preservation and preparation were as discussed by Doğan (2006). The specimens were illustrated using a Leica DM 4000 B phase-contrast microscope. The nomenclature of the idiosomal shields follows that of Summers (1962). Dorsal setal and leg setal designations follow Kethley (1990) and Grandjean (1944), respectively. Setal counts of leg segments are given with solenidia in parentheses. All measurements are given in micrometers (μm) and refer to the length of the structure unless otherwise stated. For the new species, measurements of the holotype are given first followed in parentheses by those of paratypes as a range. In newly recorded species, mean values are given first and the range is given parenthetically. Body length measurements represent the distance between the base of the gnathosoma and the posterior part of the idiosoma; width was measured at the broadest point of the idiosoma. Setae were measured from the setal base to the tip of the seta; distances between setae were measured between the setal bases. Leg length was measured from the tip of the claws to the trochanter base. Palp length was measured from the tarsal tip to the trochanter base. Specimens examined were deposited in the collection of the Acarology Laboratory of Erzincan University, Turkey.

Abbreviations

Gnathosoma

- ω = solenidion on palptarsus
- elcp* = supracoxal setae of palp
- m* = anterior or innermost pair of subcapitular setae
- n* = posterior pair of subcapitular setae
- or*₁ = 1st pair of adoral setae
- or*₂ = 2nd pair of rostral setae

Idiosoma

- 1a* = 1st pair of intercoxal setae
- 3a* = 2nd pair of intercoxal setae
- 4a* = 3rd pair of intercoxal setae
- ag*₁ = 1st pair of aggenital setae
- ag*₂ = 2nd pair of aggenital setae
- ag*₃ = 3rd pair of aggenital setae
- ag*₄ = 4th pair of aggenital setae
- c*₁ = internal pair of humeral setae
- c*₂ = external pair of humeral setae
- d*₁ = internal pair of dorsal setae
- d*₂ = external pair of dorsal setae
- e*₁ = internal pair of lumbral setae
- e*₂ = external pair of lumbral setae
- f*₁ = internal pair of sacral setae
- g*₁ = 1st pair of genital setae
- g*₂ = 2nd pair of genital setae
- h*₁ = 1st pair of clunal setae
- h*₂ = 2nd pair of clunal setae
- h*₃ = 3rd pair of clunal setae
- pob* = post-ocular bodies

- ps_1 = 1st pair of pseudanal setae
 ps_2 = 2nd pair of pseudanal setae
 ps_3 = 3rd pair of pseudanal setae
 sce = external pair of scapular setae
 sci = internal pair of scapular setae
 vi = internal pair of vertical setae
 ve = external pair of vertical setae

Legs

- ω = solenidion on tarsi
 φ = solenidion on tibia I
 $\varphi\rho$ = proximal solenidion on tibiae
 κ = solenidion on genu I
 $elcp$ = supracoxal setae of leg I

Institutional abbreviation

ALEU = Acarology Laboratory of Erzincan University, Turkey

Results

Subclass Acari Leach, 1817
Superorder Acariformes Zakhvatkin, 1952
Order Trombidiformes Reuter, 1909
Suborder Prostigmata Kramer, 1877
Superfamily Raphignathoidea Kramer, 1877
Family Stigmaeidae Oudemans, 1931

Genus *Stigmaeus* Koch, 1836

Type species

Stigmaeus cruentus Koch, 1836, by original designation.

Description

The genus *Stigmaeus* Koch, 1836 (Acari: Stigmaeidae) can be defined by the following characters: idiosoma oval; chelicerae separate; palptibial claw subequal to or slightly shorter than palptarsus; accessory claw seta-like or spine-like, terminal eupathidion on palptarsus basally fused and split into two or three long prongs; counts of setae and solenidia from palptrochanter to palptarsus: 0, 3, 1–2, 2 + 1 claw + 1 accessory claw, 4 + 1 ω + 1 subterminal spine-like eupathidion + 2 or 3 eupathidia (basally fused); subcapitulum with two pairs of subcapitular setae; prodorsum typically with a large shield, bearing three pairs of setae and a pair of platelets bearing setae *sce*; eyes present or absent, postocular bodies (*pob*) present or absent; dorsal hysterosomal area typically with one or two shields surrounded by three to five pairs of platelets, setae d_1 and d_2 never on same shield; humeral shields dorso- or ventrolateral, with setae c_2 ; intercalary shields entire or divided, with a pair of setae (f_1); suranal shield entire or divided, with two or three pairs of setae; coxisternal shields present, divided along midline; ventral opisthosoma with three to five pairs of aggenital setae; genital and anal valves fused or contiguous, with one to three pairs of genital setae and three pairs of pseudanal setae (Fan & Zhang 2005; Dönel & Doğan 2011; Doğan *et al.* 2015a, 2016).

Stigmaeus bifurcus sp. nov.

[urn:lsid:zoobank.org:act:2EB6B605-F3AA-4142-8692-BD150A8A85D0](https://doi.org/10.2307/2466605)

Figs 1–2; Table 1

Etymology

The name of this new species, *bifurcus*, refers to the shape of the eupathidion on the palptarsus.

Type material

Holotype

TURKEY: ♀, from soil under *Ulmus* sp., Erzincan, alt. 1275 m, 39°39'39.8" N, 39°29'28.5" E, 25 Oct. 2015.

Paratypes

TURKEY: 8 ♀♀, same data as holotype; 1 ♀, from soil under *Rosa canina*, same locality as holotype, 15 Mar. 2015.

Description

Female

IDIOSOMA. Elongated. Length of body (including gnathosoma): 396 (350–384). Width of body: 163 (129–157).

GNATHOSOMA (Fig. 2B). Gnathosoma punctate and 68 (65–70) long, chelicerae separate and punctate, 78 (74–81) long. Palp punctate and 91 (74–89) long. Counts of setae and solenidia from palptrochanter to palptarsus: 0, 3, 1, 2 + 1 claw + 1 seta-like accessory claw, 4 + 1 solenidion + 1 subterminal seta-like eupathidion + 1 bifurcate eupathidion. Palptibial claw slightly longer than palptarsus. Palp supracoxal setae (*elcp*) pudgy. Subcapitulum with two pairs of adoral setae (*or*_{1,2}) and two pairs of subcapitular setae (*m*, *n*). Dimensions and distance between subcapitular setae, *m* 14 (12–14), *n* 26 (22–25), *m–m* 21 (19–22), *n–n* 24 (23–26), *m–n* 12 (10–12).

DORSUM (Figs 1A, C–D, 2A). Integument striate except punctate peritremal region, propodosomal shield with reticulations, its posterior margin concave and with tubercles. Propodosomal shield bearing three pairs of setae (*vi*, *ve*, *sci*) and apodemal marking, eyes and post-ocular bodies absent. Setae *sce* on minute platelets. Central shield vaguely reticulate but other shields not ornamented. Central shield bearing two pairs setae (*c*₁, *d*₁). In one paratype specimen left member of setae *c*₁ about twice as long as right member (Fig. 1C). Setae *d*₂ located on marginal shield. Setae *e*₁ on divided median zonal shields. Setae *e*₂ located on minute platelets. Setae *f*₁ on divided intercalary shield. One pair of extra, minute and non-setose shields lateral to intercalary shields. Suranal shield divided and with three pairs of setae (*h*_{1–3}) but right member of setae *h*₃ absent in one paratype (Fig. 1D). Setae *vi*, *sci*, *d*₁ and all suranal setae slightly serrated but others smooth (Fig. 2A). Lengths and distances of dorsal idiosomal setae as follows: *vi* 16 (14–17), *ve* 42 (38–47), *sci* 20 (17–20), *sce* 41 (40–44), *c*₁ 16 (15–17), *c*₂ 46 (45–49), *d*₁ 16 (16–18), *d*₂ 34 (29–35), *e*₁ 8 (17–20), *e*₂ 18 (17–19), *f*₁ 31 (34–38), *h*₁ 21 (21–26), *h*₂ 24 (21–26), *h*₃ 13 (13–15), *vi–vi* 23 (24–29), *ve–ve* 33 (31–35), *vi–ve* 20 (19–22), *sci–sci* 46 (43–48), *ve–sci* 36 (33–35), *sce–sce* 99 (86–104), *sci–sce* 27 (23–29), *c*₁–*c*₁ 31 (26–31), *c*₂–*c*₂ 145 (124–145), *c*₁–*c*₂ 63 (46–66), *d*₂–*d*₂ 100 (89–102), *c*₁–*d*₁ 43 (40–46), *c*₁–*d*₂ 41 (35–41), *d*₁–*d*₁ 28 (27–31), *d*₂–*d*₁ 43 (39–43), *e*₂–*e*₂ 96 (83–97), *d*₂–*e*₂ 62 (51–61), *d*₁–*e*₁ 47 (40–49), *d*₁–*e*₂ 51 (44–49), *e*₁–*e*₁ 32 (29–37), *e*₂–*e*₁ 34 (27–37), *f*₁–*f*₁ 46 (43–51), *e*₁–*f*₁ 26 (24–26), *e*₂–*f*₁ 42 (36–43), *f*₁–*h*₁ 40 (35–46), *f*₁–*h*₂ 40 (30–43), *f*₁–*h*₃ 41 (27–36), *h*₁–*h*₁ 23 (22–26), *h*₂–*h*₂ 52 (48–54), *h*₃–*h*₃ 70 (57–66), *h*₁–*h*₂ 16 (12–14), *h*₂–*h*₃ 9 (6–9).




Fig. 1. *Stigmaeus bifurcus* sp. nov. **A–B.** Holotype (♀). **A.** Dorsum of body. **B.** Venter of body. – **C–D.** Paratype (♀). **C.** Abnormality: left seta c_1 about 2 times as long as the right. **D.** Abnormality: seta h_3 absent on right suranal shield in one paratype. Scale bars = 100 μ m.




Fig. 2. *Stigmaeus bifurcus* sp. nov., holotype, ♀. A. Some dorsal body setae. B. Palp. C. Leg I. D. Leg II. E. Leg III. F. Leg IV. Scale bars: A–B = 40 μ m; C–F = 100 μ m.

Table 1. Differences among the *Stigmaeus* species having laterally two extra non setose dorsal shields. – = absent; + = present.

Characters	Species									
	<i>S. bifurcus</i> sp. nov.	<i>S. hashturdiensis</i>	<i>S. caria</i>	<i>S. kerman-shahensis</i>	<i>S. shabestarensis</i>	<i>S. petilus</i>	<i>S. erzincanus</i>	<i>S. additicus</i>	<i>S. ceylani</i>	
Shape of accessory claw on palp tibia	seta-like	seta-like	spine-like	spine-like	spine-like	spine-like	seta-like	spine-like	seta-like	
Shape of palp eupathidion	bifurcate	bifurcate	bifurcate	bifurcate	trident	trident	trident	?	trident	
The number of setae on palp tibia (except for claw and accessory claw)	2	2	3	3	2	2	2	2	2	
Shape of dorsal body setae	almost smooth	smooth	almost smooth	almost smooth	smooth	slightly serrated (except for setae v_1, s_{c_2}, d_2, c_2)	stout	slightly serrated	slightly spinulate	
Platelets bearing the setae s_{c_2}	+	- or +	-	+	+	+	+	-	+	
Apodemal marking	+	- or +	-	-	-	-	+	+	+	
Additional a pair of shields next to propodosomal shield	-	-	-	-	+	-	+	-	+	
Median zonal shields	divided	divided	divided	divided	divided	entire	divided	divided	divided	
Suranal shield	divided	divided	divided	divided	entire	divided	entire	entire, but recessed anteriorly	entire	
The number of setae pairs on suranal shield	3	3	3	3	3	2	3	3	3	
Aggenital shields	longitudinally divided into two parts	longitudinally divided into two parts	longitudinally divided into two parts	longitudinally divided into two parts	longitudinally divided into two parts	longitudinally divided into two parts	longitudinally divided into two parts	longitudinally divided into two parts	divide into four parts	
Shape of aggenital setae	smooth	smooth	smooth	smooth	smooth	slightly serrated	smooth	smooth	smooth	
Pattern on ventral shields	-	- or +	-	-	-	-	+	+	-	
Setal formulae of leg genua	5(+1κ)-3-0-1	5(+1κ)-4-1-2	5(+1κ)-2-0-1	5(+1κ)-3-0-2	5(+1κ)-5-2-2	5(+1κ)-3-0-1	5(+1κ)-5-2-2	5(+1κ)-5-2-2	5(+1κ)-5-2-2	
Setal formulae of leg tarsi	13(+1ω)-8(+1ω)-7(+1ω)-7(+1ω)	13(+1ω)-8(+1ω)-7(+1ω)	13(+1ω)-8(+1ω)-7(+1ω)	13(+1ω)-8(+1ω)-7(+1ω)	13(+1ω)-9(+1ω)-7(+1ω)	13(+1ω)-8(+1ω)-7(+1ω)	13(+1ω)-9(+1ω)-7(+1ω)	13(+1ω)-9(+1ω)-7(+1ω)	13(+1ω)-9(+1ω)-7(+1ω)	

*The Turkish specimens of *S. hashturdiensis* Bagheri & Maleki, 2014 have a pair of additional small platelets (Uluçay 2015) but not in the type specimens (Bagheri *et al.* 2014).

VENTER (Fig. 1B). Almost striate. Back of coxae I–II and III–IV striate with tubercles. Humeral shields without reticulations and bearing setae c_2 . Coxisternal shields divided, smooth and bearing three pairs of setae ($1a$, $3a$, $4a$). Lengths and distances of these setae: $1a$ 14 (12–13), $3a$ 14 (12–14), $4a$ 14 (11–14), $1a-1a$ 17 (17–21), $3a-3a$ 23 (20–29), $4a-4a$ 23 (16–26). Genital and anal shields contiguous, two pairs of smooth genital ($g_{1,2}$) and three pairs of slightly serrate pseudanal setae (ps_{1-3}). ps_3 shorter than other pseudanal setae. Four pairs of aggenital setae (ag_{1-4}) on long and divided aggenital shields. Lengths of these setae as follows: ag_1 10 (8–10), ag_2 10 (8–10), ag_3 10 (8–10), ag_4 11 (9–11), g_1 7 (6–9), g_2 12 (10–13), ps_1 27 (23–27), ps_2 25 (22–25), ps_3 13 (12–14).

LEGS (Figs 2C–F). Coxae I–IV with punctuation. Length: leg I 138 (136–140), leg II 106 (100–107), leg III 114 (108–115), leg IV 128 (121–129). Counts of setae and solenidia on legs I–IV: coxae 2-2-2-2; trochanters 1-1-2-1; femora 4-4-3-2; genua 5(+1 κ)-3-0-1; tibiae 5(+1 ϕ +1 $\phi\phi$)-5(+1 $\phi\phi$)-5(+1 $\phi\phi$)-5(+1 $\phi\phi$); tarsi 13(+1 ω)-8(+1 ω)-7(+1 ω)-7(+1 ω). All tarsi with solenidia. Lengths of solenidia: I ω 11 (8–12), II ω 8 (7–9), III ω 6 (3–5), IV ω 6 (3–5).

Male and immature stages

Unknown.

Remarks

This new species resembles *Stigmaeus kermanshahiensis* Khanjani *et al.*, 2012 and *S. caria* Khanjani *et al.*, 2012 in that the eyes and post-ocular bodies are absent, suranal shield divided, one pair of extra, minute and non-setose shields lateral to the intercalary shields, and the palp tarsus bears a bifurcate eupathidium. However, it can be separated from *S. kermanshahiensis* by the following characters: one seta on genu IV (vs two setae in *S. kermanshahiensis*), palptibia with two setae (vs three setae in *S. kermanshahiensis*), and apodemal marking present (absent in *S. kermanshahiensis*). *Stigmaeus bifurcus* sp. nov. can also be distinguished from *S. caria* by the following characters: three setae on genu II (vs two setae in *S. caria*), palptibia with two setae (vs three setae in *S. caria*), *sce* on minute platelets (on striate integument in *S. caria*), and apodemal marking present (absent in *S. caria*). The new species is also similar to *S. hashtrudiensis* Bagheri & Maleki, 2014; however, it differs from the latter in its apodemal marking and different numbers of setae on genua II–IV (genua 5(+1 κ)-3-0-1 in the new species vs 5(+1 κ)-4-1-2 in *S. hashtrudiensis*). See Table 1 for a comparative presentation of characters among different *Stigmaeus* species.

Stigmaeus miandoabiensis Bagheri & Zarei, 2012

Figs 3–5

Material examined

TURKEY: 2 ♀♀, 1 DN from soil and litter under *Pinus sylvestris*, Erzincan, alt. 1210 m, 39°45'00.8" N, 39°29'17.8" E, 23 Oct. 2014; 1 ♀ from soil under *Rosa canina*, Erzincan, 1275 m, 39°39'39.8" N, 39°29'28.5" E, 15 Mar. 2015.

Description

Female (Figs 3, 5) (n=3)

IDIOSOMA. Oval. Length of body (including gnathosoma): 350 (335–371). Width of body: 120 (114–128).

GNATHOSOMA (Fig. 5B). Gnathosoma punctate and 68 (67–71) long, chelicerae separate and punctate, 79 (74–82) long. Palp punctate and 73 (69–79) long. Counts of setae and solenidia from palptrochanter to palptarsus: 0, 3, 1, 2 + 1 claw + 1 seta like accessory claw, 4 + 1 solenidion + 1 subterminal seta-like eupathidion + 1 bifurcate eupathidion. Palptibial claw about subequally as long as palptarsus. Palpal




Fig. 3. *Stigmaeus miandoabiensis* Bagheri & Zarei, 2012 (♀). A. Dorsum of body. B. Venter of body. C. Leg I. D. Leg II. E. Leg III. F. Leg IV. Scale bars = 100 µm.




Fig. 4. *Stigmaeus miandoabiensis* Bagheri & Zarei, 2012 (deutonymph). **A.** Dorsum of body. **B.** Venter of body. **C.** Leg I. **D.** Leg II. **E.** Leg III. **F.** Leg IV. Scale bars = 100 μ m.

supracoxal setae (*elcp*) pudgy. Subcapitulum with two pairs of adoral setae ($or_{1,2}$) and two pairs of subcapitular setae (m, n). Dimensions and distance between subcapitular setae: m 11 (11–12), n 16 (16–17), $m-m$ 21 (18–24), $n-n$ 23 (23–24), $m-n$ 11 (8–12).

DORSUM (Figs 3A, 5A). Integument striate except punctate peritremal region. Propodosomal shield with reticulations, its anterior and posterior margin with tubercles. Propodosomal shield bearing three pairs of setae (vi, ve, sci) and apodemal marking, eyes and post-ocular bodies absent. Setae *sce* on minute platelets. Central shield bearing two pairs of setae (c_1, d_1). Setae d_2 located on marginal shield. Setae e_1 on divided median zonal shields. Setae e_2 located on minute platelets. Setae f_1 on divided intercalary shield. Suranal shield divided and with three pairs of setae (h_{1-3}). Setae vi, sci, c_1, d_1 and e_1 slightly serrated but others smooth (Fig. 5A). Lengths and distances of dorsal idiosomal setae as follows: vi 13 (12–16), ve 41 (40–41), sci 18 (17–20), sce 40 (39–40), c_1 15 (14–18), c_2 45 (43–46), d_1 15 (14–16), d_2 25 (18–31), e_1 18 (15–22), e_2 17 (16–17), f_1 31 (28–34), h_1 21 (20–22), h_2 20 (20–21), h_3 15 (15–15), $vi-vi$ 25 (24–26), $ve-ve$ 33 (32–35), $vi-ve$ 22 (20–23), $sci-sci$ 45 (44–46), $ve-sci$ 34 (32–35), $sce-sce$ 86 (82–91), $sci-sce$ 23 (20–25), c_1-c_1 28 (27–29), c_2-c_2 115 (107–124), c_1-c_2 49 (48–51), d_2-d_2 90 (82–99), c_1-d_1 40 (39–42), c_1-d_2 38 (34–43), d_1-d_1 26 (24–28), d_2-d_1 36 (32–40), e_2-e_2 76 (70–86), d_2-e_2 54 (52–57), d_1-e_1 43 (40–45), d_1-e_2 42 (40–46), e_1-e_1 31 (29–35), e_2-e_1 24 (22–28), f_1-f_1 44 (41–48), e_1-f_1 24 (24–25), e_2-f_1 37 (34–41), f_1-h_1 32 (28–36), f_1-h_2 28 (26–30), f_1-h_3 27 (27–27), h_1-h_1 21 (20–23), h_2-h_2 46 (44–48), h_3-h_3 55 (53–57), h_1-h_2 14 (13–14), h_2-h_3 7 (6–8).




Fig. 5. *Stigmaeus miandoabiensis* Bagheri & Zarei, 2012. **A.** Some dorsal body setae. **B.** Palp (♀). Scale bars = 40 μ m.

VENTER (Fig. 3B). Almost striate. Back of coxae II and III–IV striate with tubercles. Humeral shields without ornamentation, bearing setae c_2 . Coxisternal shields divided, without ornamentation, bearing three pairs of setae ($1a$, $3a$, $4a$). Lengths and distances of these setae: $1a$ 12 (11–13), $3a$ 12 (11–13), $4a$ 11 (10–13), $1a-1a$ 19 (17–20), $3a-3a$ 23 (18–26), $4a-4a$ 21 (18–23). Genital and anal shields contiguous, two pairs of genital ($g_{1,2}$) and three pairs of pseudanal setae (ps_{1-3}), ps_3 shorter than other pseudanal setae. Four pairs of aggenital setae (ag_{1-4}) on long and divided aggenital shields. Lengths of these setae as follows: ag_1 9 (8–9), ag_2 9 (8–9), ag_3 9 (8–9), ag_4 10 (9–10), g_1 8 (7–9), g_2 9 (7–10), ps_1 24 (23–25), ps_2 21 (19–23), ps_3 11 (11–11).

LEGS (Figs 3C–F). Lengths: leg I 136 (133–142), leg II 100 (97–105), leg III 103 (102–106), leg IV 125 (118–132). Counts of setae on legs I–IV: coxae 2-2-2-2, trochanters 1-1-2-1, femora 4-4-3-2, genua 5(+1 κ)-3-0-1, tibiae 5(+1 ϕ +1 $\phi\rho$)-5(+1 $\phi\rho$)-5(+1 $\phi\rho$)-5(+1 $\phi\rho$), tarsi 13(+1 ω)-8(+1 ω)-7(+1 ω)-7(+1 ω). Lengths of solenidia: I ω 10 (10–10), II ω 8 (8–8), III ω 4 (4–5), IV ω 5 (4–6).

Deutonymph (DN) (Figs 4–5) (n=1)

IDIOSOMA. Length of body (including gnathosoma): 334. Width of body: 136.

GNATHOSOMA. Lengths: gnathosoma 60, chelicerae 69, palp 68. Number of setae on palp segments as in female. Dimensions and distance between subcapitular setae: m 12, n 15, $m-m$ 18, $n-n$ 23, $m-n$ 10.

DORSUM (Fig. 4A). As in female except suranal shields with two pairs of setae. Lengths and distances of dorsal idiosomal setae as follows: vi 13, ve 36, sci 16, sce 33, c_1 13, c_2 41, d_1 13, d_2 21, e_1 14, e_2 14, f_1 26, h_1 7, h_2 20, $vi-vi$ 26, $ve-ve$ 29, $vi-ve$ 17, $sci-sci$ 39, $ve-sci$ 31, $sce-sce$ 96, $sci-sce$ 29, c_1-c_1 27, c_2-c_2 130, c_1-c_2 52, d_2-d_2 99, c_1-d_1 37, c_1-d_2 41, d_1-d_1 23, d_2-d_1 42, e_2-e_2 94, d_2-e_2 55, d_1-e_1 41, d_1-e_2 52, e_1-e_1 30, e_2-e_1 30, f_1-f_1 40, e_1-f_1 23, e_2-f_1 37, f_1-h_1 34, f_1-h_2 29, h_1-h_1 19, h_2-h_2 41, h_1-h_2 11.

VENTER (Fig. 4B). Ventral view similar to that of female but genital shield and its setae absent, aggenital shields bearing three pairs of aggenital setae (ag_{1-3}). Lengths of ventral setae as follows: $1a$ 10, $3a$ 14, $4a$ 10, $1a-1a$ 20, $3a-3a$ 33, $4a-4a$ 33. ag_1 7, ag_2 7, ag_3 7, ps_1 18, ps_2 16, ps_3 12.

LEGS (Figs 4C–F). Lengths: leg I 122, leg II 94, leg III 105, leg IV 115. Counts of setae on legs I–IV: coxae 2-2-2-2, trochanters 0-1-2-0, femora 4-4-3-2, genua 5(+1 κ)-2-0-0, tibiae 5(+1 ϕ +1 $\phi\rho$)-5(+1 $\phi\rho$)-5(+1 $\phi\rho$)-5(+1 $\phi\rho$), tarsi 13(+1 ω)-8(+1 ω)-7(+1 ω)-7(+1 ω). Lengths of solenidia: I ω 12, II ω 7, III ω 4, IV ω 3.

Male and immature stages

Unknown.

Distribution

Iran (Bagheri & Zarei 2012; Hajizadeh *et al.* 2013; Beyzavi *et al.* 2013; Navaei-Bonab *et al.* 2015; Rahmati *et al.* 2015).

Remarks

Stigmaeus miandoabiensis was described from Iran by Bagheri & Zarei (2012). Hajizadeh *et al.* (2013) and Beyzavi *et al.* (2013) compiled checklists of Iranian prostigmatic mites, which equally included *S. miandoabiensis*. Later, Navaei-Bonab *et al.* (2015) and Rahmati *et al.* (2015) also mentioned this species briefly.

Stigmaeus miandoabiensis is very close to *S. planus* Kuznetsov, 1978, but it can be distinguished from the latter by e_2 situated on minute platelets (on striate integument in *S. planus*), ag_1 – ag_4 on the same shield ($ag_{1,2}$ on one shield and $ag_{3,4}$ on another shield in *S. planus*), and longer ve .

Table 2. The number of raphignathoid mite species known from Turkey.

Superfamily Raphignathoidea (8 families, 25 genera, 192 species)
Family Barbutiidae (1 genus, 2 species)
Genus <i>Barbutia</i> (2 species)
Family Caligonellidae (3 genera, 10 species)
Genus <i>Caligonella</i> (1 species)
Genus <i>Molothrognathus</i> (5 species)
Genus <i>Neognathus</i> (4 species)
Family Camerobiidae (3 genera, 30 species)
Genus <i>Camerobia</i> (1 species)
Genus <i>Neophyllobius</i> (18 species)
Genus <i>Tycherobius</i> (11 species)
Family Cryptognathidae (3 genera, 18 species)
Genus <i>Cryptognathus</i> (5 species)
Genus <i>Cryptofavognathus</i> (2 species)
Genus <i>Favognathus</i> (11 species)
Family Eupalopsellidae (2 genera, 5 species)
Genus <i>Eupalopsellus</i> (4 species)
Genus <i>Eupalopsis</i> (1 species)
Family Homocaligidae (1 genus, 1 species)
Genus <i>Homocaligus</i> (1 species)
Family Raphignathidae (1 genus, 23 species)
Genus <i>Raphignathus</i> (23 species)
Family Stigmaeidae (11 genera, 103 species)
Genus <i>Agistemus</i> (3 species)
Genus <i>Cheylostigmaeus</i> (5 species)
Genus <i>Eryngiopus</i> (4 species)
Genus <i>Eustigmaeus</i> (25 species)
Genus <i>Ledermuelleriopsis</i> (11 species)
Genus <i>Mediolata</i> (8 species)
Genus <i>Prostigmaeus</i> (1 species)
Genus <i>Stigmaeus</i> (38 species)
Genus <i>Storchia</i> (4 species)
Genus <i>Villersia</i> (1 species)
Genus <i>Zetzellia</i> (3 species)

This species is a new record for Turkish fauna, and its deutonymph was identified for the first time in this study. Generally, the Turkish specimens are similar to the type specimens except for the fact that the anterior and posterior regions of the propodosomal shield are covered with tubercles.

Discussion

The superfamily Raphignathoidea belongs to the suborder Prostigmata. They are worldwide in distribution, abundant in most geographical regions, and are even found in the Antarctic region (Fan & Zhang 2005). Most raphignathoid mites are known from the drier edaphic microhabitats such as litter, soil, bark and moss (Doğan *et al.* 2014). Raphignathoidea comprises 11 families, eight of which have been recorded from Turkey: Barbutiidae, Caligonellidae, Camerobiidae, Cryptognathidae, Eupalopsellidae, Homocaligidae, Raphignathidae and Stigmaeidae (Doğan 2007; Erman *et al.* 2007; Doğan *et al.* 2014). Until now, 192 raphignathoid mite species (including the two species in this paper) in 25 genera have been found in Turkey (Table 2).

Stigmaeidae, a family within the superfamily Raphignathoidea, is a large cosmopolitan group of genera distinguished by the position of the dorsal shields, number of subcapitular setae, size of the palptibial claw, shape of the terminal eupathidia on the palptarsus, cheliceral base fused or free, and presence of coxisternal shields (Dönel & Doğan 2011). Currently, it consists of 33 genera (excluding *Erynglpusopsis* Tseng, 1982 accepted as synonym of *Eryngiopus* Summers, 1964 by Doğan *et al.* [2015b]) and more than 575 species (Fan 2005; Dönel & Doğan 2011; Zhang *et al.* 2011; Fan & Flechtmann 2015; Doğan *et al.* 2015a, 2015c; Fan & Ueckermann 2016; Fan *et al.* 2016). The genus *Stigmaeus* is one of the oldest and most diverse genera in the Stigmaeidae (Fan & Zhang 2005) with 133 valid species (Fan & Zhang 2005; Doğan *et al.* 2015a, 2016; Dilkaraoğlu *et al.* 2016a, 2016b, Fan *et al.* 2016). To date, 36 species of the genus *Stigmaeus* have been reported from Turkey (Özkan *et al.* 1988, 1994; Erman *et al.* 2007; Doğan 2007; Doğan *et al.* 2015a, 2016; Dilkaraoğlu *et al.* 2016a, 2016b). In this paper, we described one new species and reported a new record of *Stigmaeus* from Turkey. This raises the total number of *Stigmaeus* species for Turkey to 38.

Some variations and asymmetric morphological abnormalities in the species are noted. In both *Stigmaeus* species, the shape of some dorsal body setae vary. Some setae are slightly serrated, whereas others are smooth (Figs 2A, 5A). Among the paratypes of *Stigmaeus bifurcus* sp. nov., in one specimen the left seta c_1 is about two times as long as the right (Fig. 1C). Seta h_3 is absent on right suranal shield in another paratype of the new species (Fig. 1D).

Acknowledgements

We would like to thank the Erzincan University, Coordinatorship of Scientific Research Projects, which financially supported this study (research project number FEN-E-140613–0018). This study is a part of the first author's MSc thesis.

References

- Bagheri M. & Zarei E. 2012. *Stigmaeus miandoabiensis* sp. nov. (Acari: Trombidiformes: Stigmaeidae), with redescription of *S. siculus* (Berlese, 1883) from Iran. *Systematic and Applied Acarology* 17: 441–447. <https://doi.org/10.11158/saa.17.4.13>
- Bagheri M., Rahimi G., Maleki N., Gharekhani G. & Saber M. 2014. *Stigmaeus hashtrudiensis*, a new species of the genus *Stigmaeus* (Acari: Trombidiformes: Stigmaeidae) from Northwest Iran. *Persian Journal of Acarology* 3: 121–128.
- Beyzavi G., Ueckermann E.A., Faraji F. & Ostovan H. 2013. A catalog of Iranian prostigmatic mites of super families Raphignathoidea and Tetranychoida (Acari). *Persian Journal of Acarology* 2: 389–474.
- Dilkaraoğlu S., Doğan S., Erman O., Sevsay S. & Adil S. 2016a. Some morphological variations and abnormalities in females of *Stigmaeus longipilis* (Canestrini) (Acari, Stigmaeidae). *Turkish Bulletin of Entomology* 6: 149–159. [In Turkish] <https://doi.org/10.16969/teb.06297>
- Dilkaraoğlu S., Doğan S., Erman O., Sevsay S. & Adil S. 2016b. Stigmaeid mites (Acari: Raphignathoidea: Stigmaeidae) of Harşit Valley and Örumcek Forests (Turkey). *Erzincan University Journal of Science and Technology* 9: 10–72. <https://doi.org/10.18185/eufbed.18373>
- Doğan S. 2006. Contributions to the knowledge of the raphignathoid mites of Turkey (Acari Raphignathoidea) with description of a new species. *International Journal of Acarology* 32: 371–375. <https://doi.org/10.1080/01647950608684484>
- Doğan S. 2007. Checklist of raphignathoid mites (Acari: Raphignathoidea) of Turkey. *Zootaxa* 1454: 1–26.

- Doğan S., Dilkaraoğlu S., Aksoy H., Aykut M. & Fan Q.-H. 2014. New occurrence of the hygrobiotic mite family Homocaligidae (Acari, Raphignathoidea) in Turkey: *Homocaligus crassipus* Fan. *Systematic & Applied Acarology* 19: 447–461. <https://doi.org/10.11158/saa.19.4.8>
- Doğan S., Bingül M., Dilkaraoğlu S. & Fan Q.-H. 2015a. Description of a new species of the genus *Stigmaeus* Koch (Acari: Stigmaeidae) from Turkey, with a list of described species in the world. *International Journal of Acarology* 41: 290–299. <https://doi.org/10.1080/01647954.2015.1028441>
- Doğan S., Dilkaraoğlu S., Fan Q.-H., Erman O., Sevsay S. & Adil S. 2015b. Description of a new species of the genus *Eryngiopus* Summers (Acari: Stigmaeidae) from Turkey. *Systematic and Applied Acarology* 20: 431–440. <https://doi.org/10.11158/saa.20.4.8>
- Doğan S., Dilkaraoğlu S., Fan Q.-H., Sevsay S., Erman O. & Adil S. 2015c. Description a species of the genus *Cheylostigmaeus* Willmann (Acari: Stigmaeidae) from Ekşisu Marsh, Turkey. *Systematic and Applied Acarology* 20: 797–808. <https://doi.org/10.11158/saa.20.7.8>
- Doğan S., Dilkaraoğlu S., Erman O., Faraji F., Bingül M., Zeytun E. & Ersin F. 2016. Redescription of *Stigmaeus solidus* Kuznetzov (Acari, Stigmaeidae) based on the Turkish and Dutch specimens. *Turkish Bulletin of Entomology* 6: 33–42. [In Turkish] <https://doi.org/10.16969/teb.41380>
- Dönel G. & Doğan S. 2011. The stigmaeid mites (Acari: Stigmaeidae) of Kelkit Valley (Turkey). *Zootaxa* 2942: 1–56.
- Erman O., Özkan M., Ayyıldız N. & Doğan S. 2007. Checklist of the mites (Arachnida: Acari) of Turkey. Second supplement. *Zootaxa* 1532: 1–21.
- Fan Q.-H. 2005. Synopsis of the described Actinedida of the world, family Stigmaeidae. In: Hallan J. (ed.) *Synopsis of the Described Arachnida of the World*. Available from <http://bug.tamu.edu/research/collection/hallan/acari/Family/Stigmaeidae.txt> [accessed 17 May 2015]
- Fan Q.-H. & Flechtmann C.H.W. 2015. Stigmaeidae. In: Carrillo D., de Moraes G.J. & Peña J. (eds) *Prospects for Biological Control of Plant Feeding Mites and Other Harmful Organisms*: 185–206. Progress in Biological Control 19, Springer International Publishing, Switzerland. https://doi.org/10.1007/978-3-319-15042-0_7
- Fan Q.-H. & Ueckermann E.A. 2016. Resurrection of the genus *Nonocaligus* Habeeb with redefinition of *Nonocaligus* and *Mullederia* Wood (Acari: Stigmaeidae). *Systematic and Applied Acarology* 21: 1447–1449. <https://doi.org/10.11158/saa.21.11.1>
- Fan Q.-H. & Zhang Z.-Q. 2005. *Fauna of New Zealand*. Vol. 52: *Raphignathoidea (Acari: Prostigmata)*. Manaaki Whenua Press, Lincoln.
- Fan Q.-H., Flechtmann C.H.W. & De Moraes D.J. 2016. Annotated catalogue of Stigmaeidae (Acari: Prostigmata), with a pictorial key to genera. *Zootaxa* 4176: 1–199. <https://doi.org/10.11646/zootaxa.4176.1.1>
- Grandjean F. 1944. Observations sur les acariens de la famille des Stigmaeidae. *Archives des Sciences physiques et naturelles* 26: 103–131.
- Hajizadeh J., Khanjani M., Faraji F. & Ueckermann E.A. 2013. Stigmaeid mites of Guilan Province of Iran with description of a new species and a checklist for Iranian stigmaeid mites (Prostigmata: Stigmaeidae). *International Journal of Acarology* 39: 571–579. <https://doi.org/10.1080/01647954.2013.850533>
- Kethley J. 1990. Acarina: Prostigmata (Actinedida). In: Dindal D.L. (ed.) *Soil Biology Guide*: 667–756. John Wiley & Sons, New York.

- Khanjani M., Pishehvar S., Mirmoayedi A.-N. & Khanjani M. 2012. Two new eyeless mite species of the genus *Stigmaeus* Koch (Acari: Stigmaeidae) from western provinces of Iran and description of the male *Stigmaeus pilatus* Kuznetsov. *International Journal of Acarology* 38: 504–513. <https://doi.org/10.1080/01647954.2012.703690>
- Kuznetsov N.N. 1978. Revision of the genus *Stigmaeus* (Acariformes, Stigmaeidae). *Zoologicheskii Zhurnal* 57: 682–694.
- Navaei-Bonab R., Kazazi M., Bagheri M., Zarei E. & Ueckermann E.A. 2015. Stigmaeid and pseudocheyleid mite fauna (Acari: Prostigmata) in three northwestern provinces of Iran with description of male and redescription of female of *Mediolata belfieldi* Momen. *Persian Journal of Acarology* 4: 373–398.
- Özkan M., Ayyıldız N. & Soysal Z. 1988. Türkiye Akar Faunası. *DOĞA TU Zooloji Dergisi* 12: 75–85. [In Turkish]
- Özkan M., Ayyıldız N. & Erman O. 1994. Check list of the Acari of Turkey. First supplement. *Euraac News Letter* 7: 4–12.
- Rahmati M., Kheradmand K., Jafari S. & Bagheri M. 2015. Fauna of Stigmaeidae and Cryptognathidae (Acari: Trombidiformes) of Lorestan province, with two new records for Iran fauna. *Journal of Crop Protection* 4: 409–418.
- Summers F.M. 1962. The genus *Stigmaeus* (Acarina: Stigmaeidae). *Hilgardia* 33 (10): 491–537.
- Uluçay İ. 2015. Two new records of the genus *Stigmaeus* (Acari: Trombidiformes: Stigmaeidae) from Turkey. *Persian Journal of Acarology* 4: 287–295.
- Zhang Z.-Q., Fan Q.-H., Pesic V., Smit H., Bochkov A.V., Khaustov A.A., Baker A., Wohltmann A., Wen T.-H., Amrine J.W., Beron P., Lin J., Gabrys G. & Husband R. 2011. Order Trombidiformes Reuter, 1909. In: Zhang Z.-Q. (ed.) *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness*. *Zootaxa* 3148: 129–138.

Manuscript received: 13 June 2016

Manuscript accepted: 23 August 2016

Published on: 3 April 2017

Topic editor: Gavin Broad

Desk editor: Chloe Chester

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'Histoire naturelle, Paris, France; Botanic Garden Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands.