



Research article

[urn:lsid:zoobank.org:pub:58C075B0-7409-41B7-A6F4-900A5A6BFECE](https://zoobank.org/pub/58C075B0-7409-41B7-A6F4-900A5A6BFECE)

***Microstomum* (Platyhelminthes, Macrostomorpha, Microstomidae) from the Swedish west coast: two new species and a population description**

Sarah ATHERTON^{1,*} & Ulf JONDELIUS²

^{1,2}Naturhistoriska riksmuseet, Box 50007, 104 05, Stockholm, Sweden.

*Corresponding author: Sarah.Atherton@nrm.se

²Email: Ulf.Jondelius@nrm.se

¹[urn:lsid:zoobank.org:author:1F597997-CD78-4F36-A82B-977B14DCAA6C](https://zoobank.org/author/1F597997-CD78-4F36-A82B-977B14DCAA6C)

²[urn:lsid:zoobank.org:author:7F116C0B-A518-45D6-B62D-0C3B459D5F70](https://zoobank.org/author/7F116C0B-A518-45D6-B62D-0C3B459D5F70)

Abstract. Two new species of marine Platyhelminthes, *Microstomum laurae* sp. nov. and *Microstomum edmondi* sp. nov. (Macrostomida: Microstomidae) are described from the west coast of Sweden. *Microstomum laurae* sp. nov. is distinguished by the following combination of characters: rounded anterior and posterior ends; presence of approximately 20 adhesive papillae on the posterior rim; paired lateral red eyespots located level with the brain; preoral gut extending anterior to brain and very small sensory pits. *Microstomum edmondi* sp. nov. is a protandrous hermaphrodite with a single ovary, single testis and male copulatory organ with stylet. It is characterized by a conical pointed anterior end, a blunt posterior end with numerous adhesive papillae along the rim, and large ciliary pits. The stylet is shaped as a narrow funnel with a short, arched tip. In addition, the first records of fully mature specimens of *Microstomum rubromaculatum* von Graff, 1882 from Fiskebäckskil and a phylogenetic analysis of *Microstomum* Schmidt, 1848 based on the mitochondrial cytochrome oxidase I (COI) gene are presented.

Keywords. Macrostomorpha, Macrostomida, flatworm, meiofauna, turbellaria.

Atherton S. & Jondelius U. 2018. *Microstomum* (Platyhelminthes, Macrostomorpha, Microstomidae) from the Swedish west coast: two new species and a population description. *European Journal of Taxonomy* 398: 1–18. <https://doi.org/10.5852/ejt.2018.398>

Introduction

Macrostomorpha Doe, 1986 is a group of free-living flatworms that may be found in aquatic and semi-aquatic habitats all over the world (Rieger 2001). Their relatively small size, typically only 1–2 mm (although animals as large as 5 mm have been recorded) and paucity of distinguishing morphological characters makes taxonomic identification within the group difficult (Janssen *et al.* 2015).

Species identification of Macrostomorpha is largely dependent on the morphology of the reproductive system, and identification of species that predominately reproduce asexually, such as most species of *Microstomum* Schmidt, 1848, is particularly challenging. It is unsurprising, then, that while limited

attention has been paid to species diversity within Macrostomorpha (e.g., Luo *et al.* 2011; Adami *et al.* 2012; Schockaert 2014; Sun *et al.* 2014; Janssen *et al.* 2015; Fang *et al.* 2016), current taxonomic research within the genus *Microstomum* is particularly scant (but see Rogozin 2015).

Species of *Microstomum* primarily reproduce through asexual fission, but their life cycle may also include short periods of sexual reproduction (Bauchhenss 1971; Heitkamp 1982). During that time, individual zooids will develop both male and female sexual structures, including for the male complex: single or bilateral testes, vas deferentia, a male copulatory apparatus with a seminal vesicle, stylet and antrum masculinum, and a male gonopore; and for the female complex: a single ovary, female antrum and female gonopore. *Microstomum* sexual maturity may be present for as little as two weeks in a year (Bauchhenss 1971; Faubel 1974) and therefore descriptions of the reproductive organs exist for only 15 of the 31 currently accepted nominal species.

Several members of *Microstomum* were encountered while investigating macrostomorph diversity in western Sweden. Herein, two new species of marine *Microstomum* are described, and a population of *Microstomum rubromaculatum* von Graff, 1882 from Fiskebäckskil is redescribed based upon the first records of fully sexual specimens. Additionally, we present a hypothesis regarding the phylogenetic position of the three species based on the mitochondrial cytochrome oxidase I (COI) gene and make COI barcode sequences available.

Material and methods

Sediments and aquatic vegetation were collected by hand from sites around the Sven Lovén Centre Kristineberg in Fiskebäckskil, Sweden on 17–20 Aug. 2015 and the Sven Lovén Centre Tjärnö in Strömstad, Sweden on 17–19 Jun. 2016. Exact dates and coordinates are listed in Table 1. Samples were transported back to the laboratory and marine flatworms were extracted within 48 hours following a standard MgCl₂ anesthetization-decantation technique (Martens 1984). Animals were manually isolated under a Nikon SMZ 1500 stereo microscope, transferred to a glass slide and identified using a Nikon Eclipse 80i compound microscope equipped with DIC (differential interference contrast). Light micrographs and digital videos were captured with a Cannon EOS 5D Mark III digital camera. Measurements were taken with an ocular micrometer. Following documentation, individual specimens were fixed in 95% ethanol and transported to the Naturhistoriska riksmuseet in Stockholm for DNA extraction and analysis.

DNA was extracted from whole animals using the DNeasy Blood & Tissue Kit (Qiagen, Valencia, CA) following the manufacturer's instructions. Amplification was performed via PCR using 0.2 ml PuReTaq Ready-To-Go PCR beads (GE Healthcare). A ~700 base-pair region of the COI gene was targeted using Mac_COIF+Mac_COIR (Janssen *et al.* 2015) or new (Mic_COIF 5'-GTT TGA GGA GGT TTG ATA GGC-3'; Mic_COIR 5'-ATC ACC CCC CTC CGT AGG AT-3') PCR and sequencing primers, and amplified with the following program: 5 minutes hotstart at 94°C; followed by 40 cycles of 30 seconds at 94°C, 90 seconds at 50°C, and 60 seconds at 72°C; and a final extension time of 10 minutes at 72°C. Products were viewed on a 0.8% agarose gel, purified using ExoSAP-IT enzymes (Exonuclease and Shrimp Alkaline Phosphatase, GE Healthcare), and sent to Macrogen (Macrogen Europe, Netherlands) for commercial sequencing.

Thirty-five COI sequences of *Microstomum* and seven outgroup sequences were used in the phylogenetic analysis (Table 1). The dataset included all COI sequences of *Microstomum* publically available in GenBank along with new sequences of the three species presented here. Outgroups were selected based on their position in the phylogenetic hypothesis of Janssen *et al.* (2015).

Sequence assembly was performed in MEGA v. 6.06 (Darriba *et al.* 2012) and trace files were manually edited. Sequences were aligned as amino acids using the standard flatworm mitochondrial genetic code

Table 1. List of specimens used in this study. The table lists specimens used in this study along with corresponding collection location, coordinates and date, as well as GenBank accession number, where available. * from Janssen *et al.* 2015; ** from Telford *et al.* 2000

Species	GenBank #	Location	Coordinates	Date
<i>Microstomum edmondi</i>	MF185700-3	Munkedal, Sweden	58°27'3" N, 11°41'10" E	15 Aug. 2015
	MF185704-11	Fiskebäckskil, Sweden	58°14'52" N, 11°27'05" E	17 Aug. 2015
<i>Microstomum laurae</i>	MF185712	Saltö, Sweden	58°52'41" N, 11°06'56" E	19 Jun. 2016
	MF185713	Saltö, Sweden	58°52'29" N, 11°08'41" E	17 Jun. 2016
<i>Microstomum lineare</i>	AJ405979**	United Kingdom		
	KP730567*	Lapptrasket, Finland	60°03'07" N, 23°40'19" E	6 Aug. 2008
	MF185697-9	Tensjön, Sweden	61°39'35" N, 15°14'52" E	2 Jul. 2016
<i>Microstomum papillosum</i>	KP730570*	Koenigshafen, Germany	55°02'24" N, 8°23'52" E	8 Mar. 2007
<i>Microstomum rubromaculatum</i>	MF185684-96	Fiskebäckskil, Sweden	58°14'59" N, 11°26'45" E	20 Aug. 2015
<i>Microstomum</i> sp. "B"	KP730580*	Mangrove Bay, Egypt	25°52'15" N, 34°25'04" E	11 Jan. 2009
<i>Microstomum</i> sp. "D"	KP730576*	Pianosa, Italy	42°34'29" N, 10°03'59" E	30 Apr. 2010
<i>Myozonaria fissipara</i>	KP730562*, KP730575*, KP730577*	Sant Andrea Bay, Italy	42°48'31" N, 10°08'30" E	26–30 Apr. 2010
	KP730574*	Pianosa, Italy	42°34'29" N, 10°03'59" E	30 Apr. 2010
<i>Myozonaria bistylifera</i>	KP730573*	Fetovaia Bay, Italy	42°43'36" N, 10°09'33" E	26 Apr. 2010
	KP730584*	Sant Andrea Bay, Italy	42°48'31" N, 10°08'30" E	26 Apr. 2010
<i>Myozonariinae</i>	KP730569*, KP730572*	Pianosa, Italy	42°34'29" N, 10°03'59" E	30 Apr. 2010

and then back-translated to nucleotides. The general time-reversible model with gamma distribution and proportion invariant sites was determined to be the best model of sequence evolution with three substitution schemes by jModelTest2 (Tamura *et al.* 2011) based upon the Akaike information criterion (Akaike 1974). Maximum Likelihood Analysis (ML) was performed in RaxmlGUI v. 1.5b1 (Silvestro & Michalak 2012) with 1000 fast bootstrap replicates.

Patristic distances were calculated using the TN93-model (Tamura & Nei 1993) with rate variation among sites and gamma distribution. Alignment gaps and ambiguous sites were not considered.

Results

Taxonomy

Order Macrostromida Karling, 1940
Family Microstromidae Luther, 1907
Genus *Microstromum* Schmidt, 1848

Microstromum laurae sp. nov.

[urn:lsid:zoobank.org:act:49378EBE-7046-49D3-BC3C-DD783580A90E](https://doi.org/10.2307/49378EBE-7046-49D3-BC3C-DD783580A90E)

Fig. 1

Diagnosis

Strap-shaped *Microstromum* with body length of 760 μm (two vegetative zooids) and rounded anterior and posterior ends. Posterior rim with approximately twenty adhesive papillae. Paired red eyespots 43 μm long and located in the lateral margins at level of the brain. Rhabdites concentrated in the anterior end above the pharynx. Nematocysts present. Preoral gut extending anterior to brain. Reproductive system unknown. GenBank accession number for partial COI sequence MF185712-3.

Etymology

This species is dedicated to Laura R. Atherton, mother of the first author.

Material examined

Holotype

SWEDEN: vegetative specimen, Strömstad, Saltö, 58°52'29" N, 11°08'41" E, 10 cm, 17 Jun. 2016, marine, eulittoral sand, S. Atherton leg. (SMNH-Type-8903, Genbank accession MF185712).

Additional material

SWEDEN: vegetative specimen, Strömstad, Saltö, 58°52'41" N, 11°06'56" E, 10 cm, 19 Jun. 2016, marine, eulittoral sand, S. Atherton leg. (SMNH-Type-8904, Genbank accession MF185713).

Description

Microstromum with a total body length of 760 μm and two vegetative zooids. Body strap-shaped with very slight constrictions between zooids and at the level of the ciliary pits. Ratio of body width:length 1:5 in slightly compressed animal. Anterior and posterior ends rounded.

Ciliary pits very small and shallow (Fig. 1A), below eyespots, 185 μm from anterior. Paired red eyespots approximately 43 μm long and present in the lateral margins of the body level with brain (Fig. 1A–B).

Many small (max. diameter 10 μm) orange lipid droplets derived from food scattered across body, heavily concentrated around pharynx and anterior end (Fig. 1A–B). Body otherwise colorless or reflective of intestine.

Epidermis uniformly covered with cilia. Nematocysts present. Rhabdite bundles, 20–30 μm long, scattered about the body, particularly concentrated in the anterior end above the pharynx (Fig. 1B). Approximately twenty adhesive papillae, 8–10 μm long, on the rim of the posterior end (Fig. 1C).

Mouth slit-like and 55 μm long. Pharynx encompassing the second fifth of the body. Preoral gut extending well anterior to brain. Intestine evenly filled with orange droplets.

Reproductive system unknown.



Fig. 1. *Microstomum laurae* sp. nov. **A.** Composite image of whole body. **B.** Anterior end. **C.** Posterior end. Abbreviations: a = auricles (ciliary pits); ap = adhesive papillae; b = brain; e = eyespots; i = intestine; m = mouth; n = nematocyst; ph = pharynx; r = rhabdite. Arrowheads indicate orange lipid droplets.

Remarks

Eight of the currently recognized *Microstomum* species (*M. bioculatum* Faubel, 1984, *M. gabriellae* Marcus, 1950, *M. giganteum* Hallez, 1878, *M. groenlandicum* Levinsen, 1879, *M. lineare* Ørsted, 1843, *M. melanophthalmum* Steinbock, 1933, *M. paràdii* Graff, 1913 and *M. spiriferum* Westblad, 1953) possess pigmented eyespots. The eyespots of *M. melanophthalmum* and *M. paràdii* are black with distinct lenses, while the eyespots of *M. laurae* sp. nov. are red. The eyespots in *M. bioculatum*, *M. gabriellae*, *M. giganteum*, *M. lineare* and *M. melanophthalmum* are situated far in the front of the animal, whereas they are level with the brain in *M. laurae* sp. nov. Finally, the eyespot of *M. groenlandicum* is unpaired, red and located medially above the brain.

Microstomum laurae sp. nov. is most similar to *M. spiriferum*, being alike in general body size and shape as well as their small ciliary pits and large bundles of rhabdites. However, *M. spiriferum* can be differentiated based on the dorsal richly yellowish pigment cells and absence of adhesive papillae (Westblad 1953). *M. laurae* sp. nov., on the other hand, is colorless or sometimes colored by its gut contents; there are no pigment cells apart from the eyespots, and it possesses a distinct line of adhesive tubes along the posterior rim. Though both species are found on the Swedish west coast, *M. spiriferum* was described from sublittoral habitats between 15 and 60 m depth, while *M. laurae* sp. nov. was collected from eulittoral sand at around 10 cm depth.

Microstomum edmondi sp. nov.

[urn:lsid:zoobank.org:act:E26B0ECB-C9DE-4FBB-A7D2-178A3C79F20C](https://doi.org/10.21203/rs.3.rs-2660932/v1)

Figs 2–3

Diagnosis

Microstomum with animal/zooid body length of 1700/750 µm. Conical pointed anterior end; blunt posterior end with numerous adhesive papillae along rim. Ciliary pits large, bottle shaped. Pigmented eyes absent. Dense field of cilia clearly covering epidermis. Preoral gut extending anteriorly to brain. Mouth distinctly encircled by glands. Protandrous hermaphrodite. Male reproductive system with single large testis. Vesicula seminalis circular to elliptical, 98 µm long, and containing the ends of numerous prostate glands in the distal part. Stylet approximately 67 µm long; shaped as an elongate, narrow funnel, slightly curved in one plane with a short, arched tip. Female reproductive system with single ovary and gonopore. Eggs develop caudally. GenBank accession number for partial COI sequences MF185700-11.

Etymology

This species is dedicated to Edmond T. Atherton, father of the first author.

Material examined

Holotype

SWEDEN: ♂, Fiskebäckskil, Kristineberg Sven Lovén Center for Marine Research, 58°14'52" N, 11°27'05" E, 25 cm, 17 Aug. 2015, marine, eulittoral sand, S. Atherton leg. (SMNH-Type-8890, Genbank accession MF185704).

Additional material

SWEDEN: 3 ♀♀, 1 ♂, 1 vegetative, Munkedal, 58°27'31" N, 11°41'10" E, 15 cm, 15 Aug. 2015, S. Atherton and Y. Jondelius leg. (SMNH-Type-8898–8902, Genbank accession MF185700-3); 2 ♂♂, 5 vegetative, same data as holotype (SMNH-Type-8891–8897, Genbank accession MF185705-11).

Description

Microstomum with vegetative chains up to four zooids long, typically two. Maximum animal/zooid length 1700/750 μm . Body strap-shaped with constrictions between zooids; ratio of body width:length approximately 1:4 in slightly compressed animal with two zooids. Anterior end conical pointed from level of ciliary pits. Posterior end blunt with row of 5–8- μm -long adhesive papillae along rim (Figs 2C, 3)

Pigmented eyes absent. Ciliary pits large and distinct; each with a wide 43 μm pore that abruptly narrows to a 15 μm wide tube; total length 55 μm (Figs 2A, 3A). Epidermis noticeably covered with dense field of cilia, 10–15 μm long (Fig. 2B). Nematocysts present. Rhabdite bundles to 45 μm long, occurring primarily in the posterior end of the animal (Fig. 2D).




Fig. 2. *Microstomum edmondi* sp. nov. **A.** Entire body. **B.** Anterior end with focus on glands around mouth. **C.** Posterior end with focus on adhesive papillae. **D.** Anterior end of male specimen. **E.** Anterior end of female specimen. Abbreviations: a = auricles (ciliary pits); ap = adhesive papillae; b = brain; eg = egg; i = intestine; m = mouth; o = ovary; ph = pharynx; r = rhabdite bundle; st = male stylet; t = testis; vs = vesicula seminalis.

Mouth distinctly encircled by glands (Figs 2B, 3A). Pharynx spherical, 150 µm in diameter connected to yellow-brown or red-brown intestine. Body colorless. Preoral gut extending well anteriorly to brain.

Protandrous hermaphrodite. Male reproductive system with single large testis connected by short vas deferens to male copulatory apparatus (Fig. 3A). Vesicula seminalis circular to elliptical, 98 µm long, and containing the ends of numerous prostate glands in the distal part (Fig. 2D). Stylet 67 µm long, shaped as a tube very slightly curved in one plane and narrowing to a short, arched tip; width at base approximately 17 µm and terminal opening 4 µm (Figs 2D, 3C). Male pore not seen.

Female reproductive system including single mediolateral ovary and ventral female gonopore (Figs 2E, 3B). Eggs develop caudally. Very small testis posterior to ovary present in some animals (Fig. 2E).

Remarks

Almost all sexually mature specimens of *Microstomum edmondi* sp. nov. displayed only male or female sexual organs. Only one individual (Fig. 2E) contained both an ovary with a single egg and what appeared to be a small testis, roughly a quarter of the size of the testes of the other male specimens. This individual otherwise lacked any discernible male copulatory apparatus (vesicula seminalis or stylet). Furthermore, animals with male reproductive anatomy were generally composed of two zooids with the sexual organs in the posterior zooid only, and animals with female anatomy were always solitary. Previous life cycle studies on *M. papillosum* Graff, 1882 and *M. spiculifer* Faubel, 1974 found that male genital organs first occur in asexually produced zooids that are otherwise well-developed, and sexual development then finishes in solitary individuals (Faubel 1974, 1976; Hellwig 1987). Thus, *M. edmondi* sp. nov. has a protandrous hermaphroditic development. Protandrous development also occurs in other species of *Microstomum*, including *M. bispiralis* Stirewalt, 1937, *M. lineare* (Müller, 1773), *M. papillosum* and *M. spiculifer* (Bauchhenss 1971; Faubel 1974, 1976; Heitkamp 1982; Hellwig 1987).

In general, *M. edmondi* sp. nov. can be easily distinguished by the shape of the male stylet. Of the 15 currently accepted species of *Microstomum* for which sexual anatomy is known, six (*M. bispiralis*, *M. giganteum* Hallez, 1878, *M. groenlandicum* Levinsen, 1879, *M. jenseni* Riedel, 1932, *M. lineare* and *M. spiriferum* Westblad, 1953) have spiraled or coiled stylets, two (*M. dermatophthalmum* Riedel, 1932 and *M. spiculifer*) have straight stylets (*M. dermatophthalmum* with a thicker distal end) and five (*M. crildensis* Faubel, 1984, *M. ornatum* Uljanin, 1870, *M. papillosum*, *M. septentrionale* Sabussow, 1900 and *M. trichotum* Marcus, 1950) have distinctly and continuously curved or crescent shaped stylets. Furthermore, the stylet of *M. edmondi* sp. nov. clearly differs from that of *M. melanophthalmum* Steinböck, 1933 by its size (67 vs 30 µm, respectively), as well as the lack of very wide, almost flat proximal rims and mid-way 90° bend.

Of the species of *Microstomum* for which sexual anatomy is known, *M. edmondi* sp. nov. is most similar to *M. hamatum* Westblad, 1953. The male reproductive system for both includes a large seminal vesicle and a 60–70-µm-long stylet with similar shape. Both species additionally include individuals with only a single large testis, although animals with smaller, paired testes were also found in *M. hamatum* (Westblad 1953). The stylet of *M. edmondi* sp. nov., however, can be distinguished by the narrower base, more gradual distal tapering and a small arched tip. *Microstomum hamatum* has a much broader funnel-shaped stylet ending in an 180° curve that forms a large hook. Other morphological differences include the pointed anterior end, large ciliary pits and lack of dark gray pigmentation in *M. edmondi* sp. nov. Finally, *M. edmondi* sp. nov. was collected from shallow, fairly clean marine sediments instead of deeper black mud.

Of the species of *Microstomum* for which the sexual organs remain undocumented, only eight inhabit marine waters (*M. bioculatum*, *M. breviceps* Marcus, 1951, *M. davenporti* von Graff, 1911, *M. lucidum*




Fig. 3. *Microstomum edmondi* sp. nov., composite sketches. **A.** Entire body of male specimen. **B.** Posterior end of female specimen. **C.** Stylet. Abbreviations: a = auricles (ciliary pits); ap = adhesive papillae; b = brain; eg = egg; i = intestine; m = mouth; n = nematocysts; o = ovary; ph = pharynx; r = rhabdites; st = male stylet; t = testis; vs = vesicula seminalis.

Fuhrmann, 1896, *M. mundum* von Graff, 1905, *M. rhabdotum* Marcus, 1951, *M. rubromaculatum* von Graff, 1882, *M. ulum* Marcus, 1950). *Microstomum edmondi* sp. nov. is clearly most similar to *M. ulum* in that both have conically pointed anterior ends, large ciliary pits and large rhabdite bundles, and both lack eyespot pigmentation. Morphological differences occur in the shape of the posterior end, where a clear constriction sets apart a rounded adhesive tail plate in *M. ulum* while the posterior of *M. edmondi* sp. nov. is more paddle-like and blunt, and perhaps in the density of the locomotory cilia. Both species may be found in shallow marine sediments but are described from very distant locales, *M. ulum* being from the southwest Atlantic near the Island of São Sebastião, Brazil (Marcus 1950) while *M. edmondi* sp. nov. was described from the Swedish west coast.

Microstomum rubromaculatum von Graff, 1882

Figs 4–5

Material examined

SWEDEN: 20 live specimens, Fiskebäckskil, Kristineberg Sven Lovén Center for Marine Research, 58°14'59" N, 11°26'45" E, 20 Aug. 2015, marine, sublittoral phytal on algae, M. Curini-Galletti leg. (Genbank accession MF185684-96).

Type locality

ITALY: Gulf of Naples, Tyrrhenian Sea. Deposition not recorded.

Habitat

Marine, sublittoral phytal in algae (e.g., *Sargassum* sp.) or benthal on shells, fine sand and mud.

Distribution

Ireland: New Harbor, Galway, 1–2 m; Malahide Inlet, Dublin, 2 m (Southern 1936).

England: Wembury (Meixner 1938).

Faroe Islands: Vaagfjord, Suderø, 10 m (Steinböck 1931).

France: Concarneau (von Graff 1913).

Iceland: North of Ísafjörður, 1–2 m (Steinböck 1938).

Sweden: Gullmar Fjord, Fiskebäckskil, 1–2 m (Westblad 1953; pers. obs. by author).

Norway: Herdla (Westblad 1934; Karling 1953).

Population description

Microstomum with field of bright red pigmentation spots on each well-developed zooid; length of pigmentation stretches from just below the anterior tip to halfway to the brain, width of pigmentation somewhat variable: either predominately at the lateral margins and thinning toward the middle or, most frequently, a band that encircles the entire body (Fig. 4A). Other small orange-red droplets may be scattered within the parenchyma of some specimens, particularly around the anterior and pharynx (Fig. 4B–C). Body otherwise colorless, clear and reflective of intestine.

Vegetative chains to four zooids long; maximum animal/zooid body length 2000/1400 µm. Body width generally consistent, accepting slight constrictions between zooids and at the level of the ciliary pits;




Fig. 4. *Microstomum rubromaculatum* von Graff, 1882. **A.** Entire body. **B.** Anterior end with focus on glands around mouth. **C.** Nematocysts. **D.** Posterior end with focus on reproductive anatomy. **E.** Male stylet. Abbreviations: a = auricles (ciliary pits); am = antrum masculinum; b = brain; gl = glands; i = intestine; m = mouth; o = ovary; ph = pharynx; s = sperm; st = male stylet; t = testis; vs = vesicula seminalis. Arrowheads indicate orange lipid droplets.

slightly tapering toward the rounded anterior end. Posterior end bluntly rounded. Posterior rims of well-developed zooids with many 5–6- μm -long posterior adhesive papillae (Fig. 5).

Epidermis uniformly covered with cilia. Bundles of 3–8 nematocysts present, scattered in the parenchyma (Karling 1966); each nematocyst 4–6 μm long (Fig. 4C). According to Westblad (1953), expelled nematocysts along the lateral body margins may resemble papillae.

Mouth slit-like at rest, but able to distend to encompass very large food. Pharynx spherical to elliptical, encompassing up to the length of the second quarter of the zooid. Preoral gut extending to brain or slightly anterior. Intestine yellow-brown or tinged with red or pink; may contain ingested prey.

Male reproductive system with paired testes located anterolaterally to male copulatory apparatus and gonopore (Figs 4D, 5). Testes round, average diameter 47 μm , containing little or no sperm. Vasa deferentia connect individually to circular vesicula seminalis. Numerous prostatic glands insert anteriorly in vesicula seminalis and extend ventrally around the center of stylet (Fig. 4D–E). Stylet a single, wide spiral bent around a 90° angle, terminating in a ~10- μm -long fingerlike hook (Figs 4E, 5C–D); average length 95 μm (range 75–112 μm); width largest at the base, ~12 μm , tapering only slightly towards the distal end, ~5 μm at the base of the hook; opening subterminal. Stylet projects into a ciliated antrum masculinum (Fig. 4A, D).

Female reproductive system typical for the genus (Figs 4D, 5). Single ovary situated mid-body, ventral to intestine, leading to ciliated female antrum. Female gonopore separate. Eggs develop caudally.

Remarks

Collected specimens generally appeared morphologically similar to the type description (von Graff 1882) and to previous accounts of *M. rubromaculatum* from Fiskebäckskil. Westblad (1953) recorded *M. rubromaculatum* from Fiskebäckskil with vegetative chains up to four zooids long, yet all currently collected specimens except one were composed of either two weakly developed zooids separated by a faint fission plane or one or two well-developed zooids only. This follows other patterns found in species of *Microstomum* in which slender chains of multiple, short zooids dominate during the asexual reproductive phase of the lifecycle while larger single or double zooid animals dominate during periods of sexual reproduction (Bauchhens 1971).

The amount of eyespot pigmentation in *M. rubromaculatum* can greatly vary between individuals. Specimens from Fiskebäckskil generally agreed with the original description of Graff (1882): paired, lateral eyespots composed of an accumulation of red pigmentation that extends medially to form a ring around the anterior end. However, pigmentation spots in four of the observed specimens remained clearly distinct, a phenomenon that has been recorded in other populations of *M. rubromaculatum* (von Graff 1913; Steinböck 1931). COI sequences were identical between specimens with two distinct eyespots and those with a circular band, which indicates amount of pigmentation is not necessarily a systematically important character. Rather, accumulation of eyespot pigmentation may be more “correlated with light intensity”, as in, e.g., *Microstomum lineare* (Bauchhens 1971). Steinböck (1938) reported a single specimen of *M. rubromaculatum* from Iceland with a large central pigment spot that thinned toward the body margins. However, such a pattern was not observed in any of our specimens, nor otherwise recorded in any other population.

Red-orange droplets (Fig. 4B–C), that have not been previously documented in *M. rubromaculatum*, were observed in 18 of the 20 live specimens. The droplets ranged in size from a diameter of ~2–10 μm and were most often located anteriorly, especially around the pharynx. The droplets were most likely lipid deposits whose presence and coloration stems from ingested food. While such deposits have not

been recorded before in *Microstomum*, colored lipid droplets are known to occur in other species of Macrostomorpha (Rieger *et al.* 1991).

The distribution of *Microstomum rubromaculatum* is wide, with populations reported from the Mediterranean, the North Sea and the Baltic. Although such patterns do occur for other macrostomorphs (e.g., *Macrostomum pusillum*, *M. rubrocinctum*, *Paramalostomum dubium* – see Ax 1956; Karling 1974; Armonies 1988), including other species of *Microstomum* (e.g., *M. lineare*, *M. papillosum* – see




Fig. 5. *Microstomum rubromaculatum* von Graff, 1882, composite sketches. **A.** Ventral view of entire body. **B.** Lateral view of posterior end. **C.** Lateral view of stylet. **D.** Ventral view of stylet. Abbreviations: a = auricles (ciliary pits); am = antrum masculinum; ap = adhesive papillae; b = brain; eg = egg; fp = female pore; i = intestine; m = mouth; mp = male pore; o = ovary; ph = pharynx; s = sperm; st = male stylet; t = testis; vs = vesicula seminalis.

Steinböck 1931; Karling 1974), the distribution may still be considered surprising giving the large geographic distances and differences in salinity and temperature (Boyer & Levitus 1994). Evidence has increasingly shown that widespread taxa previously thought to represent a single species are in fact morphologically indistinct complexes. However, all our specimens were collected from a single location in west Sweden, and thus different populations of *M. rubromaculatum* could not be compared at this time. Sexually mature specimens of *M. rubromaculatum* have not been recorded from any other populations, including those inhabiting the type locality, and therefore further research may be necessary to confirm the identity of *M. rubromaculatum* from Fiskebäckskil, Sweden before sexual anatomy can be included in the description of the species as a whole.

Phylogeny

The maximum likelihood analysis found four moderately or highly supported clades of *Microstomum* (Fig. 6). *M. rubromaculatum* was sister to *M. edmondi* sp. nov. with moderate support and further formed




Fig. 6. Phylogenetic relationships of *Microstomum* Schmidt, 1848 inferred from ML analysis of partial COI gene. Outgroups were selected based on the phylogenetic hypothesis presented in Janssen *et al.* (2015). Numbers at nodes represent bootstrap support. Genbank accession numbers are listed after each taxon name.

Table 2. Patristic distances between DNA sequences used in this study. Distances were calculated using the TN93-model (Tamura & Nei 1993) with rate variation among sites and gamma distribution. Alignment gaps and ambiguous sites were not considered. Accession numbers are given after species name. Specimens with identical sequences are listed together.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1. <i>Myo. fissipara</i> (KP730575, 577, 562)	–																	
2. <i>Myo. fissipara</i> (KP730574)	0.003	–																
3. <i>Myozonariinae</i> (KP730569, 572)	0.287	0.293	–															
4. <i>Myo. bistylifera</i> (KP730584, 573)	0.277	0.277	0.292	–														
5. <i>Mic. rubromaculatum</i> (MF185684–5, 87, 94)	0.841	0.841	1.024	0.962	–													
6. <i>Mic. rubromaculatum</i> (MF185686, 88, 89, 92–3, 96)	0.817	0.817	1.011	0.934	0.006	–												
7. <i>Mic. rubromaculatum</i> (MF185690)	0.869	0.869	1.059	0.978	0.020	0.014	–											
8. <i>Mic. rubromaculatum</i> (MF185691)	0.829	0.829	1.009	0.947	0.003	0.003	0.017	–										
9. <i>Mic. rubromaculatum</i> (MF185695)	0.817	0.817	1.011	0.949	0.011	0.006	0.011	0.008	–									
10. <i>Mic. papillosum</i> (KP730570)	0.837	0.850	0.828	0.842	0.253	0.253	0.279	0.248	0.263	–								
11. <i>Mic. lineare</i> (KP730567, AJ405979, MF185698)	0.605	0.604	0.850	0.748	0.224	0.215	0.239	0.219	0.224	0.209	–							
12. <i>Microstomum</i> (KP730580)	0.699	0.698	0.849	0.798	0.244	0.234	0.254	0.239	0.244	0.200	0.178	–						
13. <i>Microstomum</i> (KP730576)	0.675	0.675	0.847	0.771	0.219	0.209	0.233	0.214	0.219	0.233	0.173	0.224	–					
14. <i>Mic. lineare</i> (MF185697, 99)	0.605	0.605	0.851	0.749	0.219	0.210	0.234	0.215	0.219	0.204	0.003	0.174	0.169	–				
15. <i>Mic. edmondi</i> (MF185700)	0.843	0.843	1.079	0.847	0.224	0.214	0.238	0.219	0.224	0.264	0.225	0.287	0.165	0.220	–			
16. <i>Mic. edmondi</i> (MF185701–2, 04–9, 11)	0.831	0.831	1.062	0.835	0.219	0.209	0.234	0.214	0.219	0.258	0.220	0.282	0.160	0.215	0.003	–		
17. <i>Mic. edmondi</i> (MF185703)	0.844	0.844	1.080	0.848	0.224	0.214	0.239	0.219	0.224	0.264	0.225	0.287	0.165	0.220	0.006	0.003	–	
18. <i>Mic. edmondi</i> (MF185710)	0.854	0.854	1.093	0.886	0.233	0.224	0.248	0.228	0.233	0.263	0.224	0.292	0.169	0.219	0.008	0.011	0.014	–
19. <i>Mic. laurae</i> (MF185712–13)	0.822	0.821	0.853	0.840	0.223	0.218	0.243	0.218	0.228	0.268	0.238	0.233	0.228	0.233	0.274	0.269	0.274	0.274

a clade with an unidentified species of *Microstomum* (species “D” in Janssen *et al.* 2015; see Table 1) and *M. laurae* sp. nov. The other three species of *Microstomum* represented in the analysis (species “B” in Janssen *et al.* 2015, *M. lineare*, *M. papillosum*) individually comprised the remaining three clades. Patristic distances are presented in Table 2.

A true understanding of the evolutionary relationships within *Microstomum* would require multiple nuclear and mitochondrial gene sequences as well as a much greater species representation (Maddison 1997). However, the results of the ML analysis and patristic distances presented here clearly separate specimens of *M. edmondi* sp. nov., *M. laurae* sp. nov. and *M. rubromaculatum* into three distinct lineages representing the three species.

Discussion

This study extends the number of species of *Microstomum* known to occur within Sweden to nine. Eight species (*M. hamatum*, *M. jenseni*, *M. papillosum*, *M. rubromaculatum*, *M. septentrionale*, *M. spiriferum*,

and now *M. edmondi* sp. nov. and *M. laurae* sp. nov.) are marine and from the Swedish west coast. The ninth species, *M. lineare*, occurs primarily in fresh or brackish waters with salinity up to 6–8‰ (Karling 1974) and is, perhaps, the most widespread species of *Microstomum*, with populations reported from waters throughout Europe, Asia and North America (Karling 1974; Kolasa *et al.* 1987; but see Janssen *et al.* 2015). Additionally, the occurrence of sexually mature specimens of *M. rubromaculatum* from Fiskebäckskil, Sweden brings the number of species of *Microstomum* with known sexual anatomy to seventeen.

Acknowledgements

We would like to acknowledge the support of the staff at the Sven Lovén Centre for Marine Research in Kristineberg and Tjärnö, Sweden. Thank you also to Dr. Marco Curini-Galletti, Dr. Lukas Schärer and Ylva Jondelius for help with sample collection and identification. This research was supported by a grant from the Swedish Taxonomy Initiative (no. DHA 2014-151 4.3) to UJ.

References

- Adami M., Damborenea C. & Ronderos J. 2012. A new limnic species of *Macrostomum* (Platyhelminthes: Macrostomida) from Argentina and its muscle arrangement labeled with phalloidin. *Zoologischer Anzeiger* 251: 197–205. <https://doi.org/10.1016/j.jcz.2011.08.006>
- Akaike H. 1974. A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19: 716–723. <https://doi.org/10.1109/TAC.1974.1100705>
- Armonies W. 1988. Common pattern of Platyhelminthes distribution in North Sea salt marshes and in the Baltic Sea. *Archiv für Hydrobiologie* 111: 625–636.
- Ax P. 1956. Les Turbellariés des étangs côtiers du littoral méditerranéen de la France méridionale. *Vie et Milieu* supplément 5: 1–152.
- Bauchhens J. 1971. Die kleinturbellarien Frankens. Ein Beitrag zur Systematik und Ökologie der Turbellarien excl. Tricladida in Süddeutschland. *International Revue der Gesamten Hydrobiologia* 56: 609–666.
- Boyer T. & Levitus S. 1994. *Quality control and processing of historical oceanographic temperature, salinity, and oxygen data*. National Oceanic and Atmospheric Administration, National Environmental Satellite, Data, and Information Service 81: 1–72. US Department of Commerce, Washington DC.
- Darriba D., Taboada G., Doallo R. & Posada D. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9: 772–772. <https://doi.org/10.1038/nmeth.2109>
- Fang C., Wang L., Zang Y. & Wang A. 2016. Two new species of brackish-water *Macrostomum* (Platyhelminthes, Macrostomida) from southern China. *Zootaxa* 4170 (2): 298–310. <https://doi.org/10.11646/zootaxa.4170.2.4>
- Faubel A. 1974. Macrostomida (Turbellaria) von einem Sandstrand der Nordseeinsel Sylt. *Mikrofauna Meeresboden* 45: 339–370.
- Faubel A. 1976. Populationsdynamik und Lebenszyklen interstitieller Acoela und Macrostomida (Turbellaria). *Mikrofauna Meeresboden* 56: 253–359.
- Heitkamp U. 1982. Untersuchungen zur Biologie, Ökologie und Systematik limnischer Turbellarien periodischer und perennierender Kleingewässer Südniedersachsens. *Archiv für Hydrobiologie* supplement 64: 65–188.
- Hellwig M. 1987. Ökologie freilebender Plathelminthen im Grenzraum Watt-Salzwiese lenitischer Gezeitenküsten. *Microfauna Marina* 3: 157–248.

- Janssen T., Vizoso D., Schulte G., Littlewood D., Waeschenbach A. & Schärer L. 2015. The first multi-gene phylogeny of the Macrostomorpha sheds light on the evolution of sexual and asexual reproduction in basal Platyhelminthes. *Molecular Phylogenetics and Evolution* 92: 82–107.
<https://doi.org/10.1016/j.ympev.2015.06.004>
- Karling T. 1934. Ein Beitrag zur Kenntnis der Nemertinen des finnischen Meerbusens. *Memoranda Societatis pro Fauna et Flora Fennica* 10: 76–90.
- Karling T. 1966. On nematocysts and similar structure in turbellarians. *Acta Zoologica Fennica* 116: 3–28.
- Karling T. 1974. Turbellarian fauna of the Baltic proper: identification, ecology and biogeography. *Fauna Fennica* 27: 1–101.
- Kolasa J., Strayer J. & Bannon-O'Donnell E. 1987. Microturbellarians from interstitial waters, streams, and springs in southeastern New York. *Journal of the North American Benthological Society* 6: 125–132. <https://doi.org/10.2307/1467222>
- Luo H., Zhao P. & Wang A. 2011. Research on taxonomic characters and one new recorded species of genus *Macrostomum* (Platyhelminthes, Macrostomida) from China. *Sichuan Journal of Zoology* 30: 39–44. [In Chinese.]
- Maddison W. 1997. Gene trees in species trees. *Systematic Biology* 46: 523–536.
<https://doi.org/10.1093/sysbio/46.3.523>
- Marcus E. 1950. Turbellaria Brasileiros. *Boletins da Faculdade de Filosofia Ciências e Letras Zoologia* 17: 5–191.
- Martens P. 1984. Comparison of three different extraction methods for Turbellaria. *Marine Ecology Progress Series* 14: 229–234. <https://doi.org/10.3354/meps014229>
- Meixner J. 1938. *Turbellaria (Strudelwürmer) I. Allgemeiner Teil. Die Tierwelt der Nord- und Ostsee.* Akademischer Verlag, Leipzig.
- Rieger R. 2001. Phylogenetic systematics of the Macrostomorpha. In: Littlewood D. & Bray R. (eds) *Interrelationships of the Platyhelminthes*: 28–38. Taylor & Francis Inc., New York.
- Rieger R., Tyler S., Smith J. & Rieger G. 1991. Platyhelminthes: Turbellaria. In: Harrison F. & Bogish B. (eds). *Microscopic Anatomy of Invertebrates*. Vol. 3: *Platyhelminthes and Nemertinea*: 7–140. Wiley-Liss. Inc., New York.
- Rogozin A. 2015. New species of Turbellaria in the fauna of the Urals. *Biology Bulletin* 42: 755–758.
<https://doi.org/10.1134/S1062359015080075>
- Schockaert E. 2014. Marine Macrostomorpha (Platyhelminthes, Rhabditophora) from the Algarve (Southern Portugal). *Zootaxa* 3872: 577–590. <https://doi.org/10.11646/zootaxa.3872.5.8>
- Silvestro I. & Michalak D. 2012. raxmlGUI: a graphical front-end for RAxML. *Organism Diversity and Evolution* 12: 335–337. <https://doi.org/10.1007/s13127-011-0056-0>
- Southern R. 1936. Turbellaria of Ireland. *Proceedings of the Royal Irish Academy* section B 43: 43–72.
- Steinböck O. 1931. Marine Turbellaria. In: Jensen S., Lundebeck W. & Mortensen T. (eds) *Zoology of the Faeroes*. A.F. Høst & Søn, Copenhagen.
- Steinböck O. 1938. Marine Turbellaria. *The Zoology of Iceland* 2 (9): 1–26.
- Sun T., Zhang L., Wang A. & Zhang Y. 2014. Three new species of freshwater *Macrostomum* (Platyhelminthes, Macrostomida) from southern China. *Zootaxa* 4012: 120–134.
<https://doi.org/10.11646/zootaxa.4012.1.6>

Tamura K. & Nei M. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* 10: 512–526.

Tamura K., Peterson D., Peterson N., Stecher G., Nei M. & Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28: 2731–2739. <https://doi.org/10.1093/molbev/msr121>

Telford M., Herniou E., Russell R. & Littlewood D. 2000. Changes in mitochondrial genetic codes as phylogenetic characters: two examples from the flatworms. *Proceedings of the National Academy of Sciences* 97: 11359–11364. <https://doi.org/10.1073/pnas.97.21.11359>

Von Graff L. 1882. *Monographie der Turbellarien: I. Rhabdocoelida*. Verlag Wilhelm Engelmann, Leipzig.

Von Graff L. 1913. *Turbellaria II. Rhabdocoelida*. Das Tierreich, eine Zusammenstellung und Kennzeichnung der rezenten Tierformen 35, Verlag Friedländer und Sohn, Berlin.

Westblad E. 1953. Marine Macrostomida (Turbellaria) from Scandinavia and England. *Arkiv för Zoologi* 4: 391–408.

Manuscript received: 3 April 2017

Manuscript accepted: 16 June 2017

Published on: 25 January 2018

Topic editor: Rudy Jocqué

Desk editor: Kristiaan Hoedemakers

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d’Histoire naturelle, Paris, France; Botanic Garden Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Real Jardín Botánico de Madrid CSIC, Spain.